



Full wwPDB X-ray Structure Validation Report ⓘ

Jun 24, 2025 – 01:56 pm BST

PDB ID : 8PAE / pdb_00008pae
Title : Structure of the ectodomain of Atypical Porcine Pestivirus E2 at 1.2A resolution
Authors : Aitkenhead, H.; Stuart, D.I.; EL Omari, K.
Deposited on : 2023-06-07
Resolution : 1.20 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0rc1
Mogul	:	1.8.4, CSD as541be (2020)
Xtriage (Phenix)	:	2.0rc1
EDS	:	3.0
buster-report	:	1.1.7 (2018)
Percentile statistics	:	20231227.v01 (using entries in the PDB archive December 27th 2023)
CCP4	:	9.0.003 (Gargrove)
Density-Fitness	:	1.0.11
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.44


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	164625	1079 (1.20-1.20)
Clashscore	180529	1183 (1.20-1.20)
Ramachandran outliers	177936	1146 (1.20-1.20)
Sidechain outliers	177891	1146 (1.20-1.20)
RSRZ outliers	164620	1078 (1.20-1.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	AA	247	8% 52% 7% 40%
2	A	2	100%

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 1527 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Genome polypeptide.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	AA	149	Total	C	N	O	S	0	15	0
			1311	810	238	252	11			

There are 35 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
AA	-2	GLU	-	expression tag	UNP A0A1B1M0D5
AA	-1	THR	-	expression tag	UNP A0A1B1M0D5
AA	212	GLY	-	expression tag	UNP A0A1B1M0D5
AA	213	THR	-	expression tag	UNP A0A1B1M0D5
AA	214	GLY	-	expression tag	UNP A0A1B1M0D5
AA	215	GLY	-	expression tag	UNP A0A1B1M0D5
AA	216	SER	-	expression tag	UNP A0A1B1M0D5
AA	217	GLY	-	expression tag	UNP A0A1B1M0D5
AA	218	GLY	-	expression tag	UNP A0A1B1M0D5
AA	219	SER	-	expression tag	UNP A0A1B1M0D5
AA	220	GLY	-	expression tag	UNP A0A1B1M0D5
AA	221	LEU	-	expression tag	UNP A0A1B1M0D5
AA	222	ASN	-	expression tag	UNP A0A1B1M0D5
AA	223	ASP	-	expression tag	UNP A0A1B1M0D5
AA	224	ILE	-	expression tag	UNP A0A1B1M0D5
AA	225	PHE	-	expression tag	UNP A0A1B1M0D5
AA	226	GLU	-	expression tag	UNP A0A1B1M0D5
AA	227	ALA	-	expression tag	UNP A0A1B1M0D5
AA	228	GLN	-	expression tag	UNP A0A1B1M0D5
AA	229	LYS	-	expression tag	UNP A0A1B1M0D5
AA	230	ILE	-	expression tag	UNP A0A1B1M0D5
AA	231	GLU	-	expression tag	UNP A0A1B1M0D5
AA	232	TRP	-	expression tag	UNP A0A1B1M0D5
AA	233	HIS	-	expression tag	UNP A0A1B1M0D5
AA	234	GLU	-	expression tag	UNP A0A1B1M0D5
AA	235	GLY	-	expression tag	UNP A0A1B1M0D5
AA	236	ARG	-	expression tag	UNP A0A1B1M0D5

Continued on next page...

Continued from previous page...


Chain	Residue	Modelled	Actual	Comment	Reference
AA	237	THR	-	expression tag	UNP A0A1B1M0D5
AA	238	LYS	-	expression tag	UNP A0A1B1M0D5
AA	239	HIS	-	expression tag	UNP A0A1B1M0D5
AA	240	HIS	-	expression tag	UNP A0A1B1M0D5
AA	241	HIS	-	expression tag	UNP A0A1B1M0D5
AA	242	HIS	-	expression tag	UNP A0A1B1M0D5
AA	243	HIS	-	expression tag	UNP A0A1B1M0D5
AA	244	HIS	-	expression tag	UNP A0A1B1M0D5

- Molecule 2 is an oligosaccharide called 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
2	A	2	Total	C	N	O	0	0	0
			28	16	2	10			

- Molecule 3 is 2-acetamido-2-deoxy-beta-D-glucopyranose (CCD ID: NAG) (formula: C₈H₁₅NO₆) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	AA	1	Total	C	N	O	0	0
			14	8	1	5		

Continued on next page...



Continued from previous page...

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	AA	1	Total	C	N	O	0	0
			14	8	1	5		

- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	AA	160	Total	O	0	0
			160	160		

- Molecule 1: Genome polyprotein



4 Data and refinement statistics

Property	Value	Source
Space group	H 3	Depositor
Cell constants a, b, c, α , β , γ	101.12Å 101.12Å 43.12Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	26.26 – 1.20 26.26 – 1.20	Depositor EDS
% Data completeness (in resolution range)	91.3 (26.26-1.20) 91.3 (26.26-1.20)	Depositor EDS
R_{merge}	0.03	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.30 (at 1.20Å)	Xtriage
Refinement program	PHENIX 1.18.2_3874	Depositor
R, R_{free}	0.150 , 0.180 0.152 , 0.179	Depositor DCC
R_{free} test set	2287 reflections (4.87%)	wwPDB-VP
Wilson B-factor (Å ²)	18.0	Xtriage
Anisotropy	0.004	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.42 , 61.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	0.016 for h,-h-k,-l	Xtriage
F_o, F_c correlation	0.98	EDS
Total number of atoms	1527	wwPDB-VP
Average B, all atoms (Å ²)	30.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.20% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: NAG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	AA	0.33	0/1335	0.67	2/1804 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	AA	0	2

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	AA	56	GLU	N-CA-C	14.28	130.34	110.35
1	AA	55	SER	CA-CB-OG	7.45	126.00	111.10

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	AA	55	SER	Peptide
1	AA	56	GLU	Sidechain

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within

the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	AA	1311	0	1263	13	0
2	A	28	0	25	0	0
3	AA	28	0	26	1	0
4	AA	160	0	0	1	0
All	All	1527	0	1314	13	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (13) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:AA:8:TYR:CE2	1:AA:36:GLU:HG3	2.21	0.75
1:AA:125[B]:GLN:OE1	4:AA:401:HOH:O	2.14	0.64
1:AA:105:THR:HB	3:AA:301:NAG:H82	1.79	0.64
1:AA:46[B]:ILE:HG21	1:AA:59:CYS:SG	2.40	0.61
1:AA:13:LEU:HD23	1:AA:32:VAL:HG21	1.86	0.57
1:AA:17:GLU:HG3	1:AA:63:LEU:HD11	1.90	0.53
1:AA:5:ARG:HB2	1:AA:58:TYR:CE1	2.46	0.50
1:AA:5:ARG:HB2	1:AA:58:TYR:HE1	1.81	0.45
1:AA:13:LEU:HD13	1:AA:24[B]:ARG:NH1	2.31	0.45
1:AA:62[B]:ARG:HD2	1:AA:105:THR:HA	1.99	0.45
1:AA:78:CYS:SG	1:AA:108[B]:CYS:HB3	2.56	0.44
1:AA:80:GLN:HB2	1:AA:116[B]:LEU:HG	2.00	0.44
1:AA:62[A]:ARG:HD3	1:AA:105:THR:HA	2.00	0.43

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	AA	160/247 (65%)	155 (97%)	5 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	AA	147/212 (69%)	147 (100%)	0	100	100

There are no protein residues with a non-rotameric sidechain to report.

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	AA	6	GLN
1	AA	47	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

2 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the

expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	NAG	A	1	1,2	14,14,15	0.36	0	17,19,21	0.52	0
2	NAG	A	2	2	14,14,15	0.18	0	17,19,21	0.63	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAG	A	1	1,2	-	0/6/23/26	0/1/1/1
2	NAG	A	2	2	-	2/6/23/26	0/1/1/1

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.


All (2) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	2	NAG	C3-C2-N2-C7
2	A	2	NAG	C1-C2-N2-C7

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.



5.6 Ligand geometry [i](#)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	NAG	AA	301	1	14,14,15	0.20	0	17,19,21	0.61	0
3	NAG	AA	302	1	14,14,15	0.26	0	17,19,21	0.40	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns.

'-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	NAG	AA	301	1	-	0/6/23/26	0/1/1/1
3	NAG	AA	302	1	-	2/6/23/26	0/1/1/1

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (2) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
3	AA	302	NAG	O5-C5-C6-O6
3	AA	302	NAG	C4-C5-C6-O6

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	AA	301	NAG	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	AA	149/247 (60%)	0.55	19 (12%) 9 9	8, 21, 63, 90	15 (10%)

All (19) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	AA	137	ILE	8.8
1	AA	37	GLY	5.5
1	AA	136	ALA	3.8
1	AA	143	THR	3.8
1	AA	0	GLY	3.2
1	AA	36	GLU	3.2
1	AA	52	ASN	3.1
1	AA	116[A]	LEU	2.9
1	AA	53	SER	2.6
1	AA	55	SER	2.6
1	AA	153	THR	2.5
1	AA	3	HIS	2.5
1	AA	151	VAL	2.5
1	AA	58	TYR	2.3
1	AA	144	MET	2.3
1	AA	39[A]	GLU	2.2
1	AA	35	ARG	2.2
1	AA	51	ASN	2.1
1	AA	119	LYS	2.1

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ


There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	NAG	A	1	14/15	-	-	26,32,36,48	0
2	NAG	A	2	14/15	-	-	65,74,88,94	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.





6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	NAG	AA	302	14/15	0.71	0.14	66,75,85,86	0
3	NAG	AA	301	14/15	0.78	0.14	50,61,69,69	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.





6.5 Other polymers ⓘ

There are no such residues in this entry.