



# Full wwPDB X-ray Structure Validation Report i

Jun 24, 2025 – 02:20 pm BST

PDB ID : 8R3X / pdb\_00008r3x  
Title : Crystal structure of aPKC Iota kinase domain with LLGL2 peptide  
Authors : Soriano, E.V.; Earl, C.P.; Briggs, D.C.; McDonald, N.Q.  
Deposited on : 2023-11-10  
Resolution : 2.59 Å (reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>  
with specific help available everywhere you see the i symbol.

The types of validation reports are described at  
<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references](#) i) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0rc1  
Mogul : 1.8.4, CSD as541be (2020)  
Xtriage (Phenix) : 2.0rc1  
EDS : 3.0  
Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)  
CCP4 : 9.0.003 (Gargrove)  
Density-Fitness : 1.0.11  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.44


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## X-RAY DIFFRACTION


The reported resolution of this entry is 2.59 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	164625	3775 (2.60-2.60)
Clashscore	180529	4181 (2.60-2.60)
Ramachandran outliers	177936	4129 (2.60-2.60)
Sidechain outliers	177891	4129 (2.60-2.60)
RSRZ outliers	164620	3775 (2.60-2.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.



## 2 Entry composition [\(i\)](#)

There are 3 unique types of molecules in this entry. The entry contains 5721 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Protein kinase C iota type.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	340	Total	C	N	O	P	S	0	1	0
			2777	1769	467	526	2	13			
1	B	339	Total	C	N	O	P	S	0	0	0
			2753	1750	463	525	2	13			

- Molecule 2 is a protein called LLGL scribble cell polarity complex component 2.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
2	C	10	Total	C	N	O			0	0	0
			91	56	22	13					
2	D	11	Total	C	N	O			0	0	0
			96	59	23	14					

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
C	659	LYS	ARG	conflict	UNP Q6P1M3
D	659	LYS	ARG	conflict	UNP Q6P1M3


- Molecule 3 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	2	Total O 2 2	0	0
3	B	1	Total O 1 1	0	0
3	C	1	Total O 1 1	0	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Protein kinase C iota type



- Molecule 1: Protein kinase C iota type



- Molecule 2: LLGL scribble cell polarity complex component 2



- Molecule 2: LLGL scribble cell polarity complex component 2



## 4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	78.42 $\text{\AA}$ 86.32 $\text{\AA}$ 114.80 $\text{\AA}$ 90.00° 90.00° 90.00°	Depositor
Resolution ( $\text{\AA}$ )	69.00 – 2.59 69.00 – 2.59	Depositor EDS
% Data completeness (in resolution range)	99.6 (69.00-2.59) 99.6 (69.00-2.59)	Depositor EDS
$R_{\text{merge}}$	(Not available)	Depositor
$R_{\text{sym}}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.18 (at 2.58 $\text{\AA}$ )	Xtriage
Refinement program	REFMAC 5.8.0419	Depositor
$R$ , $R_{\text{free}}$	0.232 , 0.269 0.232 , 0.269	Depositor DCC
$R_{\text{free}}$ test set	1236 reflections (4.98%)	wwPDB-VP
Wilson B-factor ( $\text{\AA}^2$ )	56.9	Xtriage
Anisotropy	0.305	Xtriage
Bulk solvent $k_{\text{sol}}$ (e/ $\text{\AA}^3$ ), $B_{\text{sol}}$ ( $\text{\AA}^2$ )	0.34 , 29.8	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.94	EDS
Total number of atoms	5721	wwPDB-VP
Average B, all atoms ( $\text{\AA}^2$ )	65.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.16% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality i

### 5.1 Standard geometry i

Bond lengths and bond angles in the following residue types are not validated in this section: TPO

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.58	0/2824	1.14	7/3813 (0.2%)
1	B	0.56	0/2794	1.15	6/3771 (0.2%)
2	C	0.66	0/91	1.24	0/116
2	D	0.62	0/96	1.18	0/123
All	All	0.57	0/5805	1.15	13/7823 (0.2%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	B	0	4
2	C	0	1
All	All	0	5

There are no bond length outliers.

All (13) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	293	ASP	CA-CB-CG	8.73	121.33	112.60
1	B	463	ASP	CA-CB-CG	8.45	121.05	112.60
1	A	575	ASP	CA-CB-CG	6.14	118.74	112.60
1	A	396	ASP	CA-CB-CG	5.80	118.40	112.60
1	A	408	ASP	CA-CB-CG	5.79	118.39	112.60
1	B	408	ASP	CA-CB-CG	5.57	118.17	112.60
1	A	306	PHE	CA-CB-CG	5.43	119.23	113.80
1	A	294	GLU	CB-CG-CD	5.38	121.75	112.60
1	B	575	ASP	CA-CB-CG	5.34	117.94	112.60
1	A	255	ASP	CA-CB-CG	5.22	117.82	112.60

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	537	PRO	N-CA-CB	5.20	106.10	103.19
1	B	396	ASP	CA-CB-CG	5.15	117.75	112.60
1	B	515	ASP	CA-CB-CG	5.01	117.61	112.60

There are no chirality outliers.

All (5) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	B	272	ARG	Sidechain
1	B	346	ARG	Sidechain
1	B	372	ARG	Sidechain
1	B	572	ARG	Sidechain
2	C	649	SER	Peptide

## 5.2 Too-close contacts [\(i\)](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2777	0	2688	2	0
1	B	2753	0	2662	8	0
2	C	91	0	102	0	0
2	D	96	0	104	0	0
3	A	2	0	0	0	0
3	B	1	0	0	0	0
3	C	1	0	0	0	0
All	All	5721	0	5556	10	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

All (10) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:358:TYR:HB2	1:B:446:MET:HE1	1.85	0.57
1:A:461:ASN:HD22	1:A:464:GLN:NE2	2.03	0.57

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:316:VAL:HG21	1:B:395:THR:HG22	1.87	0.55
1:B:525:VAL:HG13	1:B:530:MET:HE3	1.95	0.48
1:B:301:THR:HG23	1:B:402:GLU:HG2	1.97	0.47
1:B:385:LEU:CD1	1:B:395:THR:HG21	2.45	0.46
1:B:301:THR:O	1:B:305:VAL:HG12	2.20	0.42
1:B:287:LYS:HA	1:B:290:VAL:HG22	2.02	0.41
1:B:316:VAL:CG2	1:B:395:THR:HG22	2.49	0.41
1:A:461:ASN:HD22	1:A:464:GLN:HE21	1.68	0.40

There are no symmetry-related clashes.

### 5.3 Torsion angles [\(i\)](#)

#### 5.3.1 Protein backbone [\(i\)](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	337/356 (95%)	323 (96%)	14 (4%)	0	100 100
1	B	335/356 (94%)	324 (97%)	11 (3%)	0	100 100
2	C	8/20 (40%)	7 (88%)	1 (12%)	0	100 100
2	D	9/20 (45%)	8 (89%)	1 (11%)	0	100 100
All	All	689/752 (92%)	662 (96%)	27 (4%)	0	100 100

There are no Ramachandran outliers to report.

#### 5.3.2 Protein sidechains [\(i\)](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	304/316 (96%)	303 (100%)	1 (0%)	91	97
1	B	301/316 (95%)	297 (99%)	4 (1%)	65	84
2	C	10/20 (50%)	10 (100%)	0	100	100
2	D	10/20 (50%)	10 (100%)	0	100	100
All	All	625/672 (93%)	620 (99%)	5 (1%)	79	91

All (5) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	404	LEU
1	B	293	ASP
1	B	305	VAL
1	B	347	GLN
1	B	404	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (7) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	461	ASN
1	A	472	GLN
1	A	478	GLN
1	B	312	HIS
1	B	418	ASN
1	B	464	GLN
1	B	558	ASN

### 5.3.3 RNA [\(i\)](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [\(i\)](#)

4 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the

expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
1	TPO	B	412	1	8,10,11	0.68	0	10,14,16	0.87	1 (10%)
1	TPO	B	564	1	8,10,11	0.74	0	10,14,16	0.92	1 (10%)
1	TPO	A	564	1	8,10,11	0.81	0	10,14,16	0.93	1 (10%)
1	TPO	A	412	1	8,10,11	0.69	0	10,14,16	0.98	1 (10%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	TPO	B	412	1	-	1/9/11/13	-
1	TPO	B	564	1	-	1/9/11/13	-
1	TPO	A	564	1	-	0/9/11/13	-
1	TPO	A	412	1	-	1/9/11/13	-

There are no bond length outliers.

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed( $^{\circ}$ )	Ideal( $^{\circ}$ )
1	A	564	TPO	O-C-CA	-2.11	119.25	124.78
1	A	412	TPO	O-C-CA	-2.10	119.28	124.78
1	B	564	TPO	O-C-CA	-2.08	119.34	124.78
1	B	412	TPO	O-C-CA	-2.01	119.51	124.78

There are no chirality outliers.

All (3) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
1	A	412	TPO	O-C-CA-CB
1	B	412	TPO	O-C-CA-CB
1	B	564	TPO	O-C-CA-CB

There are no ring outliers.

No monomer is involved in short contacts.

## 5.5 Carbohydrates [\(i\)](#)

There are no oligosaccharides in this entry.

## 5.6 Ligand geometry [\(i\)](#)

There are no ligands in this entry.

## 5.7 Other polymers [\(i\)](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [\(i\)](#)

There are no chain breaks in this entry.

## 6 Fit of model and data (i)

### 6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	338/356 (94%)	0.20	5 (1%) 71 67	26, 64, 94, 111	1 (0%)
1	B	337/356 (94%)	0.12	4 (1%) 76 72	34, 62, 101, 128	0
2	C	10/20 (50%)	0.87	1 (10%) 14 12	64, 72, 83, 90	0
2	D	11/20 (55%)	0.64	1 (9%) 16 13	62, 71, 80, 82	0
All	All	696/752 (92%)	0.18	11 (1%) 70 65	26, 63, 96, 128	1 (0%)

All (11) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	588	LEU	4.1
1	A	249	LEU	3.6
2	D	657	MET	3.1
1	A	298	TRP	2.5
1	B	587	PRO	2.4
1	A	348	ARG	2.3
1	B	296	ILE	2.1
2	C	649	SER	2.1
1	B	455	ILE	2.1
1	B	466	THR	2.1
1	A	550	ASP	2.0

### 6.2 Non-standard residues in protein, DNA, RNA chains (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
1	TPO	A	412	11/12	0.87	0.11	60,63,64,65	0

*Continued on next page...*

*Continued from previous page...*

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
1	TPO	B	412	11/12	0.88	0.11	56,60,68,69	0
1	TPO	A	564	11/12	0.89	0.09	79,81,86,86	0
1	TPO	B	564	11/12	0.94	0.08	56,59,62,62	0

### 6.3 Carbohydrates [\(i\)](#)

There are no oligosaccharides in this entry.

### 6.4 Ligands [\(i\)](#)

There are no ligands in this entry.

### 6.5 Other polymers [\(i\)](#)

There are no such residues in this entry.