



# Full wwPDB X-ray Structure Validation Report ⓘ

Feb 17, 2026 – 12:46 pm GMT

PDB ID : 8RU6 / pdb\_00008ru6  
Title : Nitratidesulfovibrio vulgaris [FeFe]-hydrogenase [FeFe]-hydrogenase variant with both subunits linked by a 4 amino acid linker peptide derived from Cpl of Clostridium pasteurianum  
Authors : Bikbaev, K.; Jaenecke, J.; Winkler, M.; Span, I.  
Deposited on : 2024-01-30  
Resolution : 1.15 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0  
Mogul : 1.8.4, CSD as541be (2020)  
Xtrriage (Phenix) : 2.0  
EDS : 3.0  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)  
CCP4 : 9.0.010 (Gargrove)  
Density-Fitness : 1.0.12  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.48


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*


The reported resolution of this entry is 1.15 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	164625	1095 (1.16-1.12)
Clashscore	180529	1248 (1.16-1.12)
Ramachandran outliers	177936	1224 (1.16-1.12)
Sidechain outliers	177891	1224 (1.16-1.12)
RSRZ outliers	164620	1095 (1.16-1.12)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	B	491	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	PG4	B	502	-	-	X	-

## 2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 8204 atoms, of which 3839 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Periplasmic [Fe] hydrogenase large subunit,CpI,Periplasmic [Fe] hydrogenase small subunit.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	H	N	O	S			
1	B	484	7543	2412	3756	631	706	38	122	9	0

There are 10 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
B	482	SER	-	expression tag	UNP P07603
B	483	ALA	-	expression tag	UNP P07603
B	484	TRP	-	expression tag	UNP P07603
B	485	SER	-	expression tag	UNP P07603
B	486	HIS	-	expression tag	UNP P07603
B	487	PRO	-	expression tag	UNP P07603
B	488	GLN	-	expression tag	UNP P07603
B	489	PHE	-	expression tag	UNP P07603
B	490	GLU	-	expression tag	UNP P07603
B	491	LYS	-	expression tag	UNP P07603

- Molecule 2 is TETRAETHYLENE GLYCOL (CCD ID: PG4) (formula: C<sub>8</sub>H<sub>18</sub>O<sub>5</sub>).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	H	O		
2	B	1	26	7	15	4	1	0
2	B	1	31	8	18	5	1	0
2	B	1	12	3	7	2	1	0
2	B	1	62	16	36	10	2	1

- Molecule 3 is IRON/SULFUR CLUSTER (CCD ID: SF4) (formula: Fe<sub>4</sub>S<sub>4</sub>) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	B	1	Total Fe S 8 4 4	0	0
3	B	1	Total Fe S 8 4 4	0	0
3	B	1	Total Fe S 8 4 4	0	0

- Molecule 4 is Binuclear [FeFe], di(thiomethyl)amine, carbon monoxide, cyanide cluster (-CO form) (CCD ID: MHX) (formula: C<sub>8</sub>H<sub>5</sub>Fe<sub>2</sub>N<sub>3</sub>O<sub>4</sub>S<sub>2</sub>) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	B	1	Total C Fe H N O S 26 8 2 7 3 4 2	0	0


- Molecule 5 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	B	480	Total O 480 480	0	0

### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Periplasmic [Fe] hydrogenase large subunit,CpI,Periplasmic [Fe] hydrogenase small subunit



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	49.40Å 88.86Å 106.59Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	45.75 – 1.15 45.75 – 1.15	Depositor EDS
% Data completeness (in resolution range)	99.9 (45.75-1.15) 99.9 (45.75-1.15)	Depositor EDS
$R_{merge}$	0.10	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.48 (at 1.15Å)	Xtrriage
Refinement program	REFMAC 5.8.0425	Depositor
R, $R_{free}$	0.137 , 0.158 0.137 , 0.158	Depositor DCC
$R_{free}$ test set	8364 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	11.0	Xtrriage
Anisotropy	0.439	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.38 , 35.9	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.48$ , $\langle L^2 \rangle = 0.30$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.98	EDS
Total number of atoms	8204	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	15.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.02% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality i

### 5.1 Standard geometry i

Bond lengths and bond angles in the following residue types are not validated in this section: MHX, SF4, PG4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	B	0.69	2/3909 (0.1%)	0.99	11/5287 (0.2%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	B	0	2

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	355	ARG	NE-CZ	-6.22	1.26	1.33
1	B	286	MET	SD-CE	-5.73	1.65	1.79

All (11) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	486	HIS	CB-CG-CD2	-8.01	120.79	131.20
1	B	13	MET	CG-SD-CE	-7.89	83.54	100.90
1	B	486	HIS	CB-CG-ND1	7.07	133.31	122.70
1	B	443	HIS	CA-CB-CG	-6.90	106.90	113.80
1	B	286	MET	CG-SD-CE	-5.97	87.76	100.90
1	B	116	ASP	CA-CB-CG	5.74	118.34	112.60
1	B	393	LYS	CB-CA-C	-5.53	102.12	111.02
1	B	39	ASP	CA-CB-CG	5.37	117.97	112.60
1	B	372	PHE	CA-CB-CG	5.27	119.07	113.80
1	B	399	ASP	CA-CB-CG	5.20	117.80	112.60
1	B	92	GLU	CB-CG-CD	5.02	121.14	112.60

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	B	355	ARG	Sidechain
1	B	8	ARG	Sidechain

## 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	B	3787	3756	3738	18	2
2	B	55	76	72	10	0
3	B	24	0	0	1	0
4	B	19	7	0	1	0
5	B	480	0	0	4	1
All	All	4365	3839	3810	20	2

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (20) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:502:PG4:O1	2:B:503:PG4:C6	2.35	0.74
1:B:94:LYS:HZ2	2:B:502:PG4:H61	1.53	0.73
1:B:54:MET:HG3	5:B:602:HOH:O	1.87	0.72
1:B:94:LYS:NZ	2:B:502:PG4:H32	2.18	0.58
1:B:100:VAL:HG22	2:B:502:PG4:H72	1.88	0.56
1:B:105[B]:MET:HE3	1:B:204:ILE:HD11	1.88	0.56
1:B:391[B]:MET:HB3	5:B:615:HOH:O	2.08	0.53
1:B:397:ILE:HG13	5:B:615:HOH:O	2.08	0.52
1:B:192:GLU:OE2	1:B:364:LYS:HD2	2.10	0.51
1:B:94:LYS:HD3	2:B:503:PG4:H72	1.93	0.50
1:B:295:ILE:O	1:B:301:GLY:HA3	2.13	0.49
1:B:94:LYS:NZ	2:B:502:PG4:H61	2.26	0.48
1:B:94:LYS:HZ2	2:B:502:PG4:H32	1.78	0.48
1:B:97:ASP:OD2	2:B:503:PG4:H81	2.13	0.48

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
4:B:508:MHX:C	4:B:508:MHX:N1	2.78	0.47
1:B:94:LYS:HE2	2:B:502:PG4:H11	1.97	0.46
1:B:13:MET:HB3	1:B:13:MET:HE3	1.86	0.43
1:B:391[B]:MET:CB	5:B:615:HOH:O	2.65	0.43
1:B:180:PRO:HD2	3:B:506:SF4:S1	2.59	0.42
1:B:97:ASP:OD2	2:B:503:PG4:C8	2.68	0.41

All (2) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:218:ARG:HE	1:B:486:HIS:HD2[2_454]	1.31	0.29
1:B:455:LYS:NZ	5:B:935:HOH:O[3_454]	2.12	0.08

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	B	491/491 (100%)	484 (99%)	7 (1%)	0	<b>100</b> <b>100</b>

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	B	407/405 (100%)	406 (100%)	1 (0%)	92 76

All (1) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	294	THR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	26	HIS

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

9 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
3	SF4	B	505	1	0,12,12	-	-	-		
2	PG4	B	501	-	10,10,12	0.20	0	9,9,11	0.54	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
3	SF4	B	507	1	0,12,12	-	-	-		
2	PG4	B	502	-	12,12,12	0.24	0	11,11,11	1.24	2 (18%)
2	PG4	B	504[B]	-	12,12,12	0.15	0	11,11,11	0.49	0
2	PG4	B	503	-	4,4,12	0.23	0	3,3,11	0.56	0
3	SF4	B	506	1	0,12,12	-	-	-		
2	PG4	B	504[A]	-	12,12,12	0.14	0	11,11,11	0.55	0
4	MHX	B	508	1	12,21,21	2.89	7 (58%)	2,42,42	0.95	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	SF4	B	505	1	-	-	0/6/5/5
2	PG4	B	501	-	-	1/8/8/10	-
3	SF4	B	507	1	-	-	0/6/5/5
2	PG4	B	502	-	-	5/10/10/10	-
2	PG4	B	504[B]	-	-	3/10/10/10	-
2	PG4	B	503	-	-	0/2/2/10	-
3	SF4	B	506	1	-	-	0/6/5/5
2	PG4	B	504[A]	-	-	4/10/10/10	-
4	MHX	B	508	1	-	-	0/5/3/3

All (7) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	B	508	MHX	C2-S2	-5.41	1.75	1.85
4	B	508	MHX	O5-C5	4.76	1.25	1.17
4	B	508	MHX	C1-S1	-3.86	1.78	1.85
4	B	508	MHX	S2-FE1	2.85	2.30	2.26
4	B	508	MHX	O-C	2.51	1.20	1.14
4	B	508	MHX	S1-FE1	2.39	2.29	2.26
4	B	508	MHX	C6-FE2	-2.26	1.88	1.95

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	502	PG4	O3-C4-C3	-2.28	100.10	110.39
2	B	502	PG4	C5-O3-C4	2.16	122.63	113.29

There are no chirality outliers.

All (13) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	B	502	PG4	O2-C3-C4-O3
2	B	504[A]	PG4	O2-C3-C4-O3
2	B	504[A]	PG4	O3-C5-C6-O4
2	B	502	PG4	O1-C1-C2-O2
2	B	504[B]	PG4	O1-C1-C2-O2
2	B	504[A]	PG4	C5-C6-O4-C7
2	B	504[A]	PG4	C3-C4-O3-C5
2	B	502	PG4	C5-C6-O4-C7
2	B	504[B]	PG4	O4-C7-C8-O5
2	B	504[B]	PG4	C1-C2-O2-C3
2	B	502	PG4	O3-C5-C6-O4
2	B	501	PG4	C4-C3-O2-C2
2	B	502	PG4	C6-C5-O3-C4


There are no ring outliers.


4 monomers are involved in 12 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	502	PG4	7	0
2	B	503	PG4	4	0
3	B	506	SF4	1	0
4	B	508	MHX	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.









### 5.7 Other polymers [i](#)

There are no such residues in this entry.

### 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	B	484/491 (98%)	-0.26	11 (2%) 61 68	5, 13, 25, 38	9 (1%)

All (11) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	484	TRP	6.5
1	B	483	ALA	5.1
1	B	486	HIS	4.9
1	B	392	PRO	3.8
1	B	485	SER	3.3
1	B	451	PHE	2.9
1	B	395	LEU	2.8
1	B	391[A]	MET	2.8
1	B	400	TYR	2.6
1	B	393	LYS	2.2
1	B	394	ASP	2.2

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum,




median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
2	PG4	B	503	5/13	0.79	0.14	32,36,39,41	1
2	PG4	B	504[A]	13/13	0.80	0.13	23,32,35,38	31
2	PG4	B	504[B]	13/13	0.80	0.13	18,26,30,32	31
2	PG4	B	502	13/13	0.87	0.12	26,34,38,40	1
2	PG4	B	501	11/13	0.92	0.09	19,24,28,30	1
3	SF4	B	505	8/8	1.00	0.01	9,9,10,10	0
3	SF4	B	506	8/8	1.00	0.01	8,8,8,8	0
3	SF4	B	507	8/8	1.00	0.02	13,14,15,16	0
4	MHX	B	508	19/19	1.00	0.04	8,9,11,11	2

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




**Electron density around SF4 B 505:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)






**Electron density around SF4 B 506:**


$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)



**Electron density around SF4 B 507:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)





## 6.5 Other polymers [i](#)

There are no such residues in this entry.