

Full wwPDB EM Validation Report (i)

Jul 8, 2025 – 02:33 PM JST

PDB ID : 8WLR / pdb 00008wlr

EMDB ID : EMD-37629

Title: Cryo-EM structure of SARS-CoV-2 prototype spike protein receptor-binding

domain in complex with hippopotamus ACE2

Authors: Han, P.; Yang, R.R.; Li, S.H.

Deposited on : 2023-09-30

Resolution : 3.12 Å(reported)

This is a Full wwPDB EM Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
https://www.wwpdb.org/validation/2017/EMValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

EMDB validation analysis : FAILED

Mogul : 1.8.5 (274361), CSD as541be (2020)

MolProbity : 4-5-2 with Phenix2.0rc1

buster-report : 1.1.7 (2018)

Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)

MapQ : FAILED

Ideal geometry (proteins) : Engh & Huber (2001) Ideal geometry (DNA, RNA) : Parkinson et al. (1996)


Validation Pipeline (wwPDB-VP) : 2.44

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: $ELECTRON\ MICROSCOPY$

The reported resolution of this entry is 3.12 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive $(\# \mathrm{Entries})$	${ m EM\ structures} \ (\#{ m Entries})$
Clashscore	210492	15764
Ramachandran outliers	207382	16835
Sidechain outliers	206894	16415

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5%

Mol	Chain	Length	Quality of chain				
1	A	597	87% 12%				
2	В	1217	14% • 84%				



2 Entry composition (i)

There are 4 unique types of molecules in this entry. The entry contains 6431 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called Angiotensin-converting enzyme.

Mol	Chain	Residues	Atoms			AltConf	Trace		
1	A	597	Total 4873	C 3110	N 805	O 929	S 29	0	0

• Molecule 2 is a protein called Spike glycoprotein.

Mol	Chain	Residues	Atoms			AltConf	Trace		
9	D	195	Total	С	N	О	S	0	0
2	Б	190	1543	989	257	289	8	U	U

There are 9 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
В	682	GLY	ARG	engineered mutation	UNP P0DTC2
В	683	SER	ARG	engineered mutation	UNP P0DTC2
В	685	SER	ARG	engineered mutation	UNP P0DTC2
В	817	PRO	PHE	engineered mutation	UNP P0DTC2
В	892	PRO	ALA	engineered mutation	UNP P0DTC2
В	899	PRO	ALA	engineered mutation	UNP P0DTC2
В	942	PRO	ALA	engineered mutation	UNP P0DTC2
В	986	PRO	LYS	engineered mutation	UNP P0DTC2
В	987	PRO	VAL	engineered mutation	UNP P0DTC2

• Molecule 3 is ZINC ION (CCD ID: ZN) (formula: Zn) (labeled as "Ligand of Interest" by depositor).

Mol	Chain	Residues	Atoms	AltConf
3	A	1	Total Zn 1 1	0

• Molecule 4 is 2-acetamido-2-deoxy-beta-D-glucopyranose (CCD ID: NAG) (formula: $C_8H_{15}NO_6$) (labeled as "Ligand of Interest" by depositor).






Mol	Chain	Residues	Atoms			AltConf	
1	В	1	Total	С	N	O	0
4	Ъ	1	14	8	1	5	0



3 Residue-property plots (i)


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Angiotensin-converting enzyme



THE SECOND SECON







4 Experimental information (i)

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, Not provided	
Number of particles used	227185	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	PHASE FLIPPING AND AMPLITUDE	Depositor
	CORRECTION	
Microscope	FEI TITAN KRIOS	Depositor
Voltage (kV)	300	Depositor
Electron dose $(e^-/\text{Å}^2)$	60	Depositor
Minimum defocus (nm)	1000	Depositor
Maximum defocus (nm)	2000	Depositor
Magnification	Not provided	
Image detector	GATAN K3 BIOQUANTUM (6k x 4k)	Depositor



5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: NAG, ZN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol Chain		Bond	lengths	Bond angles		
MIOI	Chain	RMSZ	# Z > 5	RMSZ	# Z > 5	
1	A	0.14	0/5007	0.30	0/6793	
2	В	0.14	0/1587	0.31	0/2161	
All	All	0.14	0/6594	0.31	0/8954	

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4873	0	4641	46	0
2	В	1543	0	1459	16	0
3	A	1	0	0	0	0
4	В	14	0	13	1	0
All	All	6431	0	6113	61	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (61) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.



Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:160:ARG:HA	1:A:163:TRP:HD1	1.60	0.66
1:A:520:ILE:HD12	1:A:581:VAL:HG12	1.78	0.66
2:B:343:ASN:OD1	4:B:1301:NAG:N2	2.30	0.65
1:A:25:GLN:HE21	2:B:487:ASN:HD21	1.42	0.64
1:A:160:ARG:HA	1:A:163:TRP:CD1	2.35	0.60
1:A:33:PHE:HD1	1:A:77:GLN:HE21	1.51	0.59
1:A:288:LYS:NZ	1:A:431:ASP:OD2	2.37	0.57
1:A:149:ASP:O	1:A:153:GLU:HG3	2.05	0.57
2:B:339:GLY:O	2:B:343:ASN:HB2	2.05	0.57
1:A:379:ILE:O	1:A:383:MET:HG3	2.04	0.57
1:A:367:ASP:O	1:A:371:THR:HG23	2.07	0.55
1:A:474:MET:HE1	1:A:499:ASP:H	1.71	0.55
1:A:198:ASP:N	1:A:198:ASP:OD1	2.40	0.54
1:A:435:GLU:OE1	1:A:540:TYR:OH	2.27	0.53
1:A:306:ARG:O	1:A:310:GLU:HG2	2.09	0.53
1:A:257:SER:HB2	1:A:610:TRP:CE2	2.44	0.52
1:A:171:GLU:O	1:A:175:GLN:NE2	2.36	0.52
1:A:248:LEU:HD12	1:A:262:LEU:HD22	1.93	0.52
1:A:160:ARG:HG3	1:A:163:TRP:HE1	1.73	0.51
1:A:144:LEU:HA	1:A:148:LEU:HB2	1.92	0.51
1:A:183:TYR:OH	1:A:509:ASP:OD1	2.27	0.51
1:A:592:LEU:O	1:A:596:LYS:HG3	2.12	0.50
2:B:365:TYR:HD2	2:B:388:ASN:HB3	1.78	0.49
1:A:184:VAL:HG22	1:A:464:PHE:HE1	1.77	0.49
1:A:243:TYR:OH	1:A:247:LYS:NZ	2.37	0.48
2:B:388:ASN:OD1	2:B:388:ASN:N	2.41	0.47
1:A:304:THR:HB	1:A:362:THR:HG21	1.97	0.47
1:A:72:ALA:O	1:A:76:GLU:HG2	2.15	0.47
1:A:93:THR:HG22	1:A:97:GLN:HE21	1.79	0.46
2:B:366:SER:HA	2:B:369:TYR:CE2	2.50	0.46
2:B:418:ILE:HA	2:B:422:ASN:HD22	1.80	0.46
1:A:123:THR:O	1:A:127:ILE:HG22	2.16	0.46
1:A:536:GLU:N	1:A:536:GLU:OE1	2.49	0.46
1:A:232:GLU:O	1:A:235:PRO:HD2	2.16	0.45
1:A:267:LEU:HD11	1:A:487:VAL:HG21	1.99	0.44
2:B:406:GLU:OE1	2:B:495:TYR:OH	2.34	0.44
2:B:401:VAL:HG22	2:B:509:ARG:HG2	1.98	0.44
2:B:462:LYS:HE3	2:B:462:LYS:HB3	1.80	0.44
1:A:40:LEU:HA	1:A:40:LEU:HD23	1.72	0.44
1:A:458:LYS:O	1:A:462:MET:HG3	2.18	0.44
1:A:157:ASP:OD1	1:A:157:ASP:C	2.61	0.44
1:A:303:ASP:OD1	1:A:303:ASP:N	2.39	0.44

Continued on next page...



Continued from previous page...

A + 1	A4 0	Interatomic	Clash
Atom-1	Atom-2	${\rm distance}({\rm \AA})$	overlap (Å)
1:A:216:ASP:OD2	1:A:216:ASP:C	2.62	0.43
1:A:117:LEU:HD13	1:A:186:LEU:HB3	2.01	0.43
1:A:144:LEU:HB2	1:A:168:TRP:CZ3	2.53	0.43
2:B:359:SER:HA	2:B:524:VAL:HG22	2.01	0.43
1:A:346:PRO:HB3	1:A:360:MET:HG3	2.01	0.42
1:A:109:LEU:HD23	1:A:109:LEU:HA	1.91	0.42
1:A:69:LYS:HE3	1:A:69:LYS:HB2	1.85	0.42
1:A:145:GLU:HA	1:A:146:PRO:HA	1.86	0.42
2:B:350:VAL:HG12	2:B:422:ASN:HB3	2.02	0.42
1:A:269:ASP:OD1	1:A:272:GLY:N	2.50	0.41
2:B:376:THR:O	2:B:434:ILE:HA	2.21	0.41
1:A:152:MET:HE3	1:A:270:MET:HA	2.02	0.41
1:A:186:LEU:HD23	1:A:186:LEU:HA	1.92	0.41
2:B:438:SER:O	2:B:438:SER:OG	2.31	0.41
1:A:194:ASN:OD1	1:A:194:ASN:N	2.54	0.40
1:A:55:ILE:HD11	1:A:343:VAL:HG23	2.04	0.40
1:A:177:ARG:HB2	1:A:498:CYS:HB2	2.02	0.40
2:B:353:TRP:CD1	2:B:353:TRP:H	2.40	0.40
2:B:405:ASP:OD1	2:B:405:ASP:N	2.54	0.40

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Perce	entiles
1	A	595/597~(100%)	587 (99%)	8 (1%)	0	100	100
2	В	193/1217 (16%)	186 (96%)	7 (4%)	0	100	100
All	All	788/1814 (43%)	773 (98%)	15 (2%)	0	100	100

There are no Ramachandran outliers to report.



5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	525/526 (100%)	524 (100%)	1 (0%)	92 96
2	В	168/1065 (16%)	168 (100%)	0	100 100
All	All	693/1591 (44%)	692 (100%)	1 (0%)	92 97

All (1) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	431	ASP

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (13) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	25	GLN
1	A	43	GLN
1	A	77	GLN
1	A	90	GLN
1	A	97	GLN
1	A	102	GLN
1	A	122	ASN
1	A	241	HIS
1	A	531	GLN
1	A	535	HIS
2	В	414	GLN
2	В	450	ASN
2	В	501	ASN

5.3.3 RNA (i)

There are no RNA molecules in this entry.



5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no oligosaccharides in this entry.

5.6 Ligand geometry (i)

Of 2 ligands modelled in this entry, 1 is monoatomic - leaving 1 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Pos	Link	Bond lengths		Bond angles		les	
IVIOI	туре	Chain	nes	Lilik	Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	NAG	В	1301	2	14,14,15	0.37	0	17,19,21	0.75	1 (5%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	NAG	В	1301	2	-	2/6/23/26	0/1/1/1

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$\mathbf{Observed}(^{o})$	$\operatorname{Ideal}({}^{o})$
4	В	1301	NAG	C1-O5-C5	2.58	115.69	112.19

There are no chirality outliers.

All (2) torsion outliers are listed below:




Mol	Chain	Res	Type	Atoms
4	В	1301	NAG	C4-C5-C6-O6
4	В	1301	NAG	O5-C5-C6-O6

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	В	1301	NAG	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.

