



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 9, 2026 – 09:49 AM UTC

PDB ID : 8Z43 / pdb_00008z43
Title : Beta-galactosidase from Bacteroides xyloxydans (E350G, ligand-free)
Authors : Nakajima, M.; Motouchi, S.; Kobayashi, K.
Deposited on : 2024-04-16
Resolution : 1.91 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.91 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1188 (1.92-1.92)
Clashscore	190562	1209 (1.92-1.92)
Ramachandran outliers	187476	1195 (1.92-1.92)
Sidechain outliers	187428	1195 (1.92-1.92)
RSRZ outliers	180081	1188 (1.92-1.92)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	551	 5% 88% 7% . .
1	B	551	 5% 86% 8% . .

2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 8996 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Beta-galactosidase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	528	4228	2724	698	789	17	0	1	0
1	B	528	4228	2724	698	789	17	0	1	0

There are 20 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	8	MET	-	initiating methionine	UNP D6CYU7
A	350	GLY	GLU	engineered mutation	UNP D6CYU7
A	551	LEU	-	expression tag	UNP D6CYU7
A	552	GLU	-	expression tag	UNP D6CYU7
A	553	HIS	-	expression tag	UNP D6CYU7
A	554	HIS	-	expression tag	UNP D6CYU7
A	555	HIS	-	expression tag	UNP D6CYU7
A	556	HIS	-	expression tag	UNP D6CYU7
A	557	HIS	-	expression tag	UNP D6CYU7
A	558	HIS	-	expression tag	UNP D6CYU7
B	8	MET	-	initiating methionine	UNP D6CYU7
B	350	GLY	GLU	engineered mutation	UNP D6CYU7
B	551	LEU	-	expression tag	UNP D6CYU7
B	552	GLU	-	expression tag	UNP D6CYU7
B	553	HIS	-	expression tag	UNP D6CYU7
B	554	HIS	-	expression tag	UNP D6CYU7
B	555	HIS	-	expression tag	UNP D6CYU7
B	556	HIS	-	expression tag	UNP D6CYU7
B	557	HIS	-	expression tag	UNP D6CYU7
B	558	HIS	-	expression tag	UNP D6CYU7

- Molecule 2 is GLYCEROL (CCD ID: GOL) (formula: C₃H₈O₃) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	1	Total C O 6 3 3	0	0
2	A	1	Total C O 6 3 3	0	0


- Molecule 3 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	264	Total O 264 264	0	0
3	B	264	Total O 264 264	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Beta-galactosidase



- Molecule 1: Beta-galactosidase



4 Data and refinement statistics i

Property	Value	Source
Space group	I 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	144.70Å 51.06Å 180.09Å 90.00° 90.42° 90.00°	Depositor
Resolution (Å)	48.15 – 1.91 48.15 – 1.91	Depositor EDS
% Data completeness (in resolution range)	99.8 (48.15-1.91) 99.8 (48.15-1.91)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.41 (at 1.91Å)	Xtrriage
Refinement program	REFMAC 5.8.0419	Depositor
R, R_{free}	0.174 , 0.202 0.183 , 0.208	Depositor DCC
R_{free} test set	5220 reflections (5.08%)	wwPDB-VP
Wilson B-factor (Å ²)	25.6	Xtrriage
Anisotropy	0.030	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.39 , 34.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.47$, $\langle L^2 \rangle = 0.30$	Xtrriage
Estimated twinning fraction	0.024 for h,-k,-l	Xtrriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	8996	wwPDB-VP
Average B, all atoms (Å ²)	29.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.92% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GOL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.63	0/4345	1.08	9/5901 (0.2%)
1	B	0.63	0/4345	1.09	7/5901 (0.1%)
All	All	0.63	0/8690	1.08	16/11802 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1
1	B	0	2
All	All	0	3

There are no bond length outliers.

All (16) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	377	ASP	CA-CB-CG	7.72	120.32	112.60
1	B	380	ASP	CA-CB-CG	7.34	119.94	112.60
1	B	509	THR	CA-CB-OG1	-6.40	100.00	109.60
1	B	481	THR	CA-CB-OG1	-6.15	100.37	109.60
1	B	494	ASP	CA-CB-CG	6.07	118.67	112.60
1	A	521	ASP	CA-CB-CG	6.05	118.65	112.60
1	A	278	ILE	CB-CA-C	6.02	116.36	111.06
1	A	201	LYS	CB-CA-C	-5.88	101.41	110.81
1	A	405	ASP	CA-CB-CG	5.83	118.43	112.60
1	A	114	PHE	CA-CB-CG	5.81	119.61	113.80
1	B	256	LYS	CG-CD-CE	5.65	124.30	111.30
1	B	246	LYS	N-CA-CB	5.52	118.83	110.28

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	420	THR	CA-CB-OG1	-5.48	101.38	109.60
1	A	179	THR	CA-CB-OG1	-5.21	101.78	109.60
1	A	284	PHE	CA-CB-CG	5.07	118.87	113.80
1	A	380	ASP	CA-CB-CG	5.03	117.63	112.60

There are no chirality outliers.

All (3) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	406	ARG	Sidechain
1	B	240	ARG	Sidechain
1	B	508	ARG	Sidechain

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4228	0	4112	17	0
1	B	4228	0	4112	28	0
2	A	12	0	16	0	0
3	A	264	0	0	3	0
3	B	264	0	0	2	0
All	All	8996	0	8240	45	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (45) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:67:LYS:HG3	3:A:808:HOH:O	1.47	1.13
1:A:459:TYR:HE2	3:A:929:HOH:O	1.68	0.76
1:A:314:LYS:HE2	1:A:323:LEU:HD22	1.67	0.75
1:B:423:MET:HE3	1:B:428:LEU:HD22	1.79	0.64
1:B:509:THR:HB	1:B:537:VAL:HG13	1.80	0.62

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:550:ARG:C	3:B:784:HOH:O	2.44	0.60
1:A:508:ARG:HD3	1:A:538:TRP:CH2	2.38	0.59
1:B:465:LYS:H	1:B:465:LYS:HE2	1.69	0.58
1:B:296:MET:O	1:B:299:VAL:HG13	2.05	0.56
1:B:100:ASP:OD2	1:B:104:LYS:HE2	2.05	0.56
1:B:465:LYS:H	1:B:465:LYS:CE	2.19	0.56
1:A:353:MET:HE1	1:A:378:GLY:HA3	1.92	0.51
1:A:493:GLY:HA2	1:A:498:HIS:CD2	2.45	0.51
1:A:286:ASN:HA	1:A:324:ALA:O	2.11	0.51
1:B:286:ASN:HA	1:B:324:ALA:O	2.11	0.51
1:A:26:LYS:HE3	1:A:32:TRP:CZ2	2.46	0.51
1:B:508:ARG:HB2	1:B:538:TRP:CZ3	2.46	0.51
1:A:305:PRO:HB2	1:A:323:LEU:HG	1.94	0.50
1:A:294:ASP:OD2	3:A:701:HOH:O	2.20	0.49
1:B:228:LYS:O	1:B:228:LYS:HG3	2.13	0.49
1:A:517:ASN:HB3	1:A:532:THR:HG22	1.93	0.49
1:A:117:TRP:HA	1:A:121:GLU:O	2.13	0.49
1:A:433:ASP:OD1	1:A:457:LYS:HE2	2.13	0.47
1:B:166:LYS:O	1:B:170:GLU:HG3	2.15	0.46
1:A:423:MET:HE3	1:A:428:LEU:HD22	1.98	0.46
1:B:517:ASN:HB3	1:B:532:THR:HG22	1.99	0.45
1:B:508:ARG:HG2	1:B:509:THR:N	2.33	0.44
1:A:529:TYR:CD2	1:A:531:PRO:HD3	2.52	0.44
1:B:166:LYS:HE3	1:B:278:ILE:HD13	2.00	0.44
1:B:324:ALA:HB1	1:B:349:PRO:HD3	2.00	0.43
1:B:352:VAL:HG12	1:B:354:LYS:HG2	2.00	0.43
1:B:493:GLY:HA2	1:B:498:HIS:CD2	2.54	0.43
1:A:539:SER:HB2	1:A:540:PRO:CD	2.49	0.43
1:A:360:TRP:CZ2	1:A:364:GLU:HG2	2.54	0.42
1:B:118:LYS:HG2	1:B:189:ASN:HB3	2.00	0.42
1:B:310:MET:HG3	1:B:323:LEU:HD12	2.00	0.42
1:B:440:LEU:C	1:B:440:LEU:HD12	2.44	0.42
1:B:360:TRP:CZ2	1:B:364:GLU:HG2	2.54	0.42
1:B:412:LYS:HG3	1:B:432:TYR:CD2	2.54	0.42
1:B:33:GLN:HG2	1:B:402:GLN:OE1	2.20	0.42
1:B:117:TRP:HA	1:B:121:GLU:O	2.20	0.42
1:B:518:TYR:OH	1:B:533:SER:OG	2.28	0.42
1:B:208:GLU:OE2	1:B:246:LYS:HE3	2.20	0.41
1:B:246:LYS:HD2	3:B:693:HOH:O	2.21	0.40
1:B:252:ASN:O	1:B:256:LYS:HG3	2.21	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	525/551 (95%)	514 (98%)	11 (2%)	0	100	100
1	B	525/551 (95%)	513 (98%)	12 (2%)	0	100	100
All	All	1050/1102 (95%)	1027 (98%)	23 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	449/471 (95%)	442 (98%)	7 (2%)	55	44
1	B	449/471 (95%)	434 (97%)	15 (3%)	33	17
All	All	898/942 (95%)	876 (98%)	22 (2%)	42	28

All (22) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	224	LYS
1	A	278	ILE
1	A	288	TRP
1	A	323	LEU
1	A	423	MET
1	A	435	GLU
1	A	532	THR
1	B	23	ARG

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	31	SER
1	B	226	LEU
1	B	246	LYS
1	B	278	ILE
1	B	288	TRP
1	B	299	VAL
1	B	404	SER
1	B	423	MET
1	B	465	LYS
1	B	466	LYS
1	B	517	ASN
1	B	532	THR
1	B	533	SER
1	B	537	VAL

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (7) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	33	GLN
1	A	222	ASN
1	B	33	GLN
1	B	213	GLN
1	B	217	GLN
1	B	222	ASN
1	B	447	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	GOL	A	601	-	5,5,5	0.16	0	5,5,5	0.51	0
2	GOL	A	602	-	5,5,5	0.42	0	5,5,5	0.88	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	GOL	A	601	-	-	2/4/4/4	-
2	GOL	A	602	-	-	2/4/4/4	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (4) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	A	601	GOL	C1-C2-C3-O3
2	A	602	GOL	C1-C2-C3-O3
2	A	601	GOL	O2-C2-C3-O3
2	A	602	GOL	O2-C2-C3-O3


There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is

within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	528/551 (95%)	0.01	25 (4%) 36 41	12, 26, 47, 77	1 (0%)
1	B	528/551 (95%)	0.02	26 (4%) 35 39	13, 25, 51, 85	1 (0%)
All	All	1056/1102 (95%)	0.02	51 (4%) 35 40	12, 26, 48, 85	2 (0%)

All (51) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	537	VAL	6.6
1	B	520	ALA	5.6
1	B	510	PHE	5.5
1	B	528	VAL	4.9
1	B	527	PHE	4.5
1	B	536	ALA	4.4
1	B	526	ILE	4.3
1	A	380	ASP	4.1
1	A	537	VAL	4.1
1	B	511	THR	4.1
1	B	509	THR	4.0
1	B	514	LYS	3.3
1	A	510	PHE	3.2
1	B	521	ASP	3.1
1	B	532	THR	3.1
1	A	515	SER	3.1
1	A	520	ALA	3.0
1	B	518	TYR	3.0
1	B	534	TYR	2.9
1	A	509	THR	2.9
1	B	512	SER	2.8
1	B	516	ASN	2.7
1	A	518	TYR	2.7
1	B	519	ASN	2.7

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	529	TYR	2.7
1	A	533	SER	2.7
1	B	465	LYS	2.7
1	B	517	ASN	2.7
1	B	535	GLN	2.6
1	A	534	TYR	2.6
1	B	531	PRO	2.6
1	A	512	SER	2.5
1	A	528	VAL	2.5
1	A	278	ILE	2.5
1	B	515	SER	2.5
1	A	532	THR	2.4
1	A	526	ILE	2.3
1	B	382	PHE	2.3
1	A	513	GLU	2.3
1	A	516	ASN	2.3
1	A	517	ASN	2.2
1	A	511	THR	2.2
1	A	521	ASP	2.2
1	A	527	PHE	2.1
1	B	530	SER	2.1
1	A	514	LYS	2.1
1	B	522	HIS	2.1
1	A	144	GLY	2.1
1	A	531	PRO	2.1
1	A	522	HIS	2.1
1	A	406	ARG	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.


6.4 Ligands [i](#)


In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum,

median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	GOL	A	601	6/6	0.91	0.12	27,38,42,47	0
2	GOL	A	602	6/6	0.94	0.10	23,28,31,32	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.





6.5 Other polymers [i](#)

There are no such residues in this entry.