

# Integrative Structure Validation Report

October 09, 2025 - 04:40 PM PDT

*The following software was used in the production of this report:*


*IHMValidation Version 3.0*

*Python-IHM Version 2.5*

PDB ID	9A1J   pdb_00009a1j
PDB-Dev ID	PDBDEV_00000091
Structure Title	Integrative structure determination of the A3G-CRL5-Vif complex (flexible)
Structure Authors	Robyn M Kaake; Ignacia Echeverria; Seung Joong Kim; John Von Dollen; Nicholas M Chesarino; Yuqing Feng; Clinton Yu; Hai Ta; Linda Chelico; Lan Huang; John Gross; Andrej Sali; Nevan J Krogan
Deposited on	2021-08-11

*This is a PDB-IHM Structure Validation Report.*

*We welcome your comments at [helpdesk@pdb-ihm.org](mailto:helpdesk@pdb-ihm.org)*

*A user guide is available at [https://pdb-ihm.org/validation\\_help.html](https://pdb-ihm.org/validation_help.html) with specific help available everywhere you see the  symbol.*

*List of references used to build this report is available [here](#).*

## 1. Overview

### 1.1. Summary

*This entry consists of 1 model(s). A total of 12 dataset(s) were used to build this entry.*


Name	Type	Count
Crosslinking-MS data	Experimental data	2
Mass Spectrometry data	Experimental data	1
Experimental model	Starting model	6

Name	Type	Count
Comparative model	Starting model	3


## 1.2. Overall quality ?

This validation report contains model quality assessments for all structures, data quality and fit to model assessments for SAS and crosslinking-MS datasets. Data quality and fit to model assessments for other datasets and model uncertainty are under development. Number of plots is limited to 256.

### Model Quality: Excluded Volume Analysis ?



### Fit to Data Used for Modeling ?



## 2. Model Details ?

### 2.1. Ensemble information ?

This entry consists of 1 distinct ensemble(s).

### 2.2. Representation ?

This entry has 1 representation(s).

ID	Model(s)	Entity ID	Molecule name	Chain(s) [auth]	Total residues	Rigid segments	Flexible segments	Model coverage/ Starting model coverage (%)	Scale
1	1	1	CBFB	A	182	1-156	157-182	100.00 / 85.71	Multiscale: Coarse-grained: 1 - 5 residue(s) per bead
		2	Vif	B	175	6-154, 166-175	1-5, 155-165	100.00 / 90.86	Coarse-grained: 1 residue(s) per bead
		3	EloB	C	161	1-105	106-161	100.00 / 65.22	Multiscale: Coarse-grained: 1 - 5 residue(s) per bead

ID	Model(s)	Entity ID	Molecule name	Chain(s) [auth]	Total residues	Rigid segments	Flexible segments	Model coverage/ Starting model coverage (%)	Scale
		4	EloC	D	112	17-112	1-16	100.00 / 85.71	Multiscale: Coarse-grained: 1 - 5 residue(s) per bead
		5	CUL5	E	780	11-302, 308-382, 405-515, 521-568, 574-687, 695-780	1-10, 303-307, 383-404, 516-520, 569-573, 688-694	100.00 / 93.08	Coarse-grained: 1 residue(s) per bead
		6	Rbx2	F	113	27-113	1-26	100.00 / 76.99	Coarse-grained: 1 residue(s) per bead
		7	A3G	G	384	6-194, 200-243, 258-380	1-5, 195-199, 244-257, 381-384	100.00 / 92.71	Coarse-grained: 1 residue(s) per bead

### 2.3. Datasets used for modeling

There are 12 unique datasets used to build the models in this entry.

ID	Dataset type	Database name	Data access code
1	Experimental model	PDB	<a href="#">pdb_00004n9f</a>
2	Experimental model	PDB	<a href="#">pdb_00001ldj</a>
3	Experimental model	PDB	<a href="#">pdb_00002ecl</a>
4	Experimental model	PDB	<a href="#">pdb_00002ma9</a>
5	Comparative model	Zenodo	<a href="#">10.5281/zenodo.5176959</a>
6	Experimental model	PDB	<a href="#">pdb_00005k81</a>
7	Comparative model	Zenodo	<a href="#">10.5281/zenodo.5176959</a>
8	Experimental model	PDB	<a href="#">pdb_00003v4k</a>
9	Comparative model	Zenodo	<a href="#">10.5281/zenodo.5176959</a>
10	Mass Spectrometry data	PRIDE	<a href="#">PXD025391</a>
11	Crosslinking-MS data	Zenodo	<a href="#">10.5281/zenodo.5176959</a>
12	Crosslinking-MS data	Zenodo	<a href="#">10.5281/zenodo.5176959</a>

### 2.4. Methodology and software

This entry is a result of 1 distinct protocol(s).

Step number	Protocol ID	Method name	Method type	Method description	Number of computed models	Multi state modeling	Multi scale modeling
1	1	Sampling	Replica exchange monte carlo	Not available	203100	False	True

There are 4 software packages reported in this entry.

ID	Software name	Software version	Software classification	Software location
1	IMP PMI module	develop-548de65454	integrative model building	<a href="https://integrativemodeling.org">https://integrativemodeling.org</a>
2	Integrative Modeling Platform (IMP)	develop-548de65454	integrative model building	<a href="https://integrativemodeling.org">https://integrativemodeling.org</a>
3	MODELLER	9.20	comparative modeling	<a href="https://salilab.org/modeller/">https://salilab.org/modeller/</a>
4	MODELLER	9.19	comparative modeling	<a href="https://salilab.org/modeller/">https://salilab.org/modeller/</a>

### 3. Data quality ?

#### 3.2. Crosslinking-MS

At the moment, data validation is only available for crosslinking-MS data deposited as a fully *compliant* dataset in the *PRIDE Crosslinking* database. Correspondence between crosslinking-MS and entry entities is established using *pyHMMER*. Only residue pairs that passed the reported threshold are used for the analysis. The values in the report have to be interpreted in the context of the experiment (i.e. only a minor fraction of in-situ or in-vivo dataset can be used for modeling).

Crosslinking-MS dataset is not available in the *PRIDE Crosslinking* database.

#### 3.4. Mass Spectrometry ?

Validation for this section is under development.

### 4. Model quality ?

For models with atomic structures, MolProbity analysis is performed. For models with coarse-grained or multi-scale structures, excluded volume analysis is performed.

#### 4.1a. Excluded Volume Analysis ?

Excluded volume satisfaction for the models in the entry are listed below. The *Analysed* column shows the number of particle-particle or particle-atom pairs for which excluded volume was analysed.

Model ID	Analysed	Number of violations	Excluded Volume Satisfaction (%)
1	1675365	4692	99.72

### 5. Fit to Data Used for Modeling Assessment ?

#### 5.2. Crosslinking-MS ?

##### 5.2.1. Restraint types ?



This table summarizes information about crosslinker(s) used for data generation, and how crosslinking information was translated into actual modeling restraints. Restraints assigned "*by-residue*" are interpreted as between CA atoms. Restraints between coarse-grained beads are indicated as "*coarse-grained*". *Restraint group* represents a set of crosslinking restraints applied collectively in the modeling.

There are 132 crosslinking restraints combined in 132 restraint groups.

Linker	Residue 1	Atom 1	Residue 2	Atom 2	Restraint type	Distance, Å	Count
DSSO	LYS	CA	LYS	CA	upper bound	26.00	125
DSSO	LYS	CA	MET	CA	upper bound	26.00	7

#### Distograms of individual restraints

Distograms (i.e., histogram plots of distances) provide an overview of distributions of distances between residues for which chemical crosslinks were identified. The shift of the distogram relative to the threshold value may indicate a poor model. Restraints with identical thresholds are grouped into one plot. Only the best distance per restraint per model group/ensemble is plotted. Inter- and intramolecular (including self-links) restraints are also grouped into one plot. Distance for a restraint between coarse-grained beads is calculated as a minimal distance between shells; if beads intersect, the distance will be reported as 0.0. A bead with the highest available resolution for a given residue is used for the assessment.



#### 5.2.2. Satisfaction of restraints ?


*Satisfaction of restraints is calculated on a **restraint group** (a set of crosslinking restraints applied collectively in the modeling) level. Satisfaction of a restraint group depends on satisfaction of individual restraints in the group and the conditionality (all/any). A restraint group is considered satisfied, if the condition was met in at least one model of the model group/ensemble. The number of measured restraints can be smaller than the total number of restraint groups if crosslinks involve non-modeled residues. Only deposited models are used for validation right now.*

State group	State	Model group	# of Deposited models/Total	Restraint group type	Satisfied (%)	Violated (%)	Count (Total=132)
1	1	1	1/198632	All	81.91	18.09	94
				Heteromeric links/ Intermolecular	62.86	37.14	35
				Self-links/ Intramolecular	93.22	6.78	59

#### Per-model satisfaction rates in ensembles

*Every point represents one model in a model group/ensemble. Where possible, boxplots with quartile marks are also plotted.*

### Satisfaction rates in Model Group 1



## 5.4. Mass Spectrometry ?

Validation for this section is under development.

## 6. Fit to Data Used for Validation Assessment ?

Validation for this section is under development.

### Acknowledgments

*The development of integrative model validation metrics, implementation of a model validation pipeline, and creation of a validation report for integrative structures are funded by NSF awards to the [PDB-IHM team](#) (DBI-1756248, DBI-2112966, DBI-2112967, DBI-2112968, and DBI-1756250) and awards from NSF, NIH, and DOE to the [RCSB PDB](#) (DBI-2321666, R01GM157729, and DE-SC0019749). The PDB-IHM team and members of the [Sali lab](#) contributed model validation metrics and software packages.*

*Dr. Jill Trewhella, Dr. Dina Schneidman, and members of the [SASBDB](#) repository are acknowledged for their advice and support in implementing SAS validation methods. Team members from the labs of Dr. Juri Rappsilber, Dr. Alexander Leitner, Dr. Andrea Graziadei, and members of [PRIDE](#) database are acknowledged for their advice and support in implementing crosslinking-MS validation methods. We are grateful to Dr. Shruthi Viswanath for discussions about uncertainty assessment of integrative structural models.*

*Members of the [wwPDB Integrative/Hybrid Methods Task Force](#) provided recommendations and community support for the project.*