

# Integrative Structure Validation Report

October 09, 2025 - 04:48 PM PDT

*The following software was used in the production of this report:*

*IHMValidation Version 3.0*

*Python-IHM Version 2.5*


*MolProbity Version 4.5.2*

*pyHMMER Version 0.11.1*

PDB ID	9A7G   pdb_00009a7g
PDB-Dev ID	PDBDEV_00000345
Structure Title	Integrative model of ODPa-ODP2 by crosslinking MS and deep learning
Structure Authors	Stahl, K.; Brock, O.; Rappsilber, J.
Deposited on	2024-01-23

*This is a PDB-IHM Structure Validation Report.*

*We welcome your comments at [helpdesk@pdb-ihm.org](mailto:helpdesk@pdb-ihm.org)*

*A user guide is available at [https://pdb-ihm.org/validation\\_help.html](https://pdb-ihm.org/validation_help.html) with specific help available everywhere you see the  symbol.*

*List of references used to build this report is available [here](#).*

## 1. Overview

### 1.1. Summary


*This entry consists of 1 model(s). A total of 1 dataset(s) were used to build this entry.*

Name	Type	Count
Crosslinking-MS data	Experimental data	1


### 1.2. Overall quality

This validation report contains model quality assessments for all structures, data quality and fit to model assessments for SAS and crosslinking-MS datasets. Data quality and fit to model assessments for other datasets and model uncertainty are under development. Number of plots is limited to 256.


### Model Quality: MolProbity Analysis ?



### Data Quality ?



### Fit to Data Used for Modeling ?



## 2. Model Details ?

### 2.1. Ensemble information ?

This entry consists of 0 distinct ensemble(s).

### 2.2. Representation ?

This entry has 1 representation(s).

ID	Model(s)	Entity ID	Molecule name	Chain(s) [auth]	Total residues	Rigid segments	Flexible segments	Model coverage/ Starting model coverage (%)	Scale
1	1	1	ODPA_BACSU	A	371	-	1-371	100.00 / 0.00	Atomic
		2	ODP2_BACSU	B	442	-	1-442	100.00 / 0.00	Atomic

### 2.3. Datasets used for modeling ?

There is 1 unique dataset used to build the models in this entry.

ID	Dataset type	Database name	Data access code
1	Crosslinking-MS data	PRIDE	<a href="#">PXD035508</a>

## 2.4. Methodology and software

This entry is a result of 1 distinct protocol(s).

Step number	Protocol ID	Method name	Method type	Method description	Number of computed models	Multi state modeling	Multi scale modeling
1	1	AlphaLink2	AlphaLink2	Not available	1	False	False

There is 1 software package reported in this entry.

ID	Software name	Software version	Software classification	Software location
1	<a href="#">AlphaLink2</a>	1.00	model building	<a href="https://github.com/Rappsilber-Laboratory/AlphaLink2">https://github.com/Rappsilber-Laboratory/AlphaLink2</a>

## 3. Data quality

### 3.2. Crosslinking-MS

At the moment, data validation is only available for crosslinking-MS data deposited as a fully *compliant* dataset in the *PRIDE Crosslinking* database. Correspondence between crosslinking-MS and entry entities is established using *pyHMMER*. Only residue pairs that passed the reported threshold are used for the analysis. The values in the report have to be interpreted in the context of the experiment (i.e. only a minor fraction of in-situ or in-vivo dataset can be used for modeling).

#### [PXD035508](#)

Number of entities in the crosslinking-MS dataset: 810

Number of entities in the entry: 2

Matching entities:

Entity ID	Molecule name	Crosslinking-MS Entity ID	E-value	Exact match
1	ODPA_BACSU	dbseq_P21881_target	0.00	True
2	ODP2_BACSU	dbseq_P21883_target	0.00	True

Residue pairs stats:

Source	Total	In matched entities	Total matched
9A7G	7	7 (100.00%)	6 (85.71%)
PXD035508	3666	61 (1.66%)	6 (0.16%)

## 4. Model quality

For models with atomic structures, MolProbity analysis is performed. For models with coarse-grained or multi-scale structures, excluded volume analysis is performed.

### 4.1b. MolProbity Analysis

Excluded volume satisfaction for the models in the entry are listed below. The Analysed column shows the number of particle-particle or particle-atom pairs for which excluded volume was analysed.

#### Standard geometry: bond outliers

There are no bond length outliers.

#### Standard geometry: angle outliers

There are 49 bond angle outliers in this entry (0.57% of 8623 assessed bonds). A summary is provided below.

Chain	Res	Type	Atoms	Z	Observed (Å)	Ideal (Å)	Model ID (Worst)	Models (Total)
B	184	GLU	C-N-CA	10.64	140.86	121.70	1	1
B	189	GLU	C-N-CA	7.80	135.73	121.70	1	1
B	187	PRO	C-N-CA	7.00	134.29	121.70	1	1
B	186	ALA	CA-C-N	6.17	126.15	116.90	1	1
B	186	ALA	N-CA-C	6.16	128.25	111.00	1	1
B	189	GLU	CA-C-N	5.71	127.62	116.20	1	1
B	189	GLU	O-C-N	5.62	114.01	123.00	1	1
B	184	GLU	CA-C-N	5.45	127.11	116.20	1	1
B	184	GLU	O-C-N	5.40	114.36	123.00	1	1
B	109	GLN	C-N-CA	5.14	130.95	121.70	1	1
B	131	GLN	OE1-CD-NE2	5.14	117.46	122.60	1	1
A	84	GLN	OE1-CD-NE2	5.11	117.49	122.60	1	1
B	187	PRO	CA-C-N	5.07	126.34	116.20	1	1
B	187	PRO	O-C-N	4.92	115.13	123.00	1	1
B	133	GLU	C-N-CA	4.89	130.51	121.70	1	1
B	427	HIS	CB-CG-CD2	4.85	124.90	131.20	1	1
A	224	GLN	OE1-CD-NE2	4.80	117.80	122.60	1	1
B	183	GLN	OE1-CD-NE2	4.75	117.85	122.60	1	1
A	291	ASN	OD1-CG-ND2	4.70	117.90	122.60	1	1
B	109	GLN	OE1-CD-NE2	4.69	117.91	122.60	1	1
B	118	GLN	OE1-CD-NE2	4.59	118.01	122.60	1	1
B	237	HIS	CB-CG-CD2	4.58	125.24	131.20	1	1
B	40	GLN	OE1-CD-NE2	4.58	118.02	122.60	1	1
A	295	GLN	OE1-CD-NE2	4.52	118.08	122.60	1	1
B	129	GLN	OE1-CD-NE2	4.51	118.09	122.60	1	1

Chain	Res	Type	Atoms	Z	Observed (Å)	Ideal (Å)	Model ID (Worst)	Models (Total)
A	150	GLN	OE1-CD-NE2	4.48	118.12	122.60	1	1
B	110	ASP	CA-CB-CG	4.44	117.04	112.60	1	1
B	128	GLN	OE1-CD-NE2	4.41	118.19	122.60	1	1
B	190	THR	CA-CB-OG1	4.40	116.19	109.60	1	1
B	263	GLN	OE1-CD-NE2	4.39	118.21	122.60	1	1
B	116	GLN	OE1-CD-NE2	4.36	118.24	122.60	1	1
B	436	GLN	OE1-CD-NE2	4.35	118.25	122.60	1	1
B	102	GLN	OE1-CD-NE2	4.31	118.29	122.60	1	1
B	252	HIS	CB-CG-CD2	4.31	125.60	131.20	1	1
A	27	GLN	OE1-CD-NE2	4.26	118.34	122.60	1	1
A	22	GLN	OE1-CD-NE2	4.24	118.36	122.60	1	1
A	48	GLN	OE1-CD-NE2	4.21	118.39	122.60	1	1
B	85	GLN	OE1-CD-NE2	4.19	118.41	122.60	1	1
B	194	GLN	OE1-CD-NE2	4.19	118.41	122.60	1	1
B	100	GLN	OE1-CD-NE2	4.12	118.48	122.60	1	1
A	15	GLN	OE1-CD-NE2	4.12	118.48	122.60	1	1
B	288	ASN	OD1-CG-ND2	4.12	118.48	122.60	1	1
B	130	ASP	C-N-CA	4.10	129.08	121.70	1	1
B	300	GLN	OE1-CD-NE2	4.09	118.51	122.60	1	1
A	65	ARG	NE-CZ-NH2	4.06	122.86	119.20	1	1
B	188	GLN	OE1-CD-NE2	4.04	118.56	122.60	1	1
A	361	GLN	OE1-CD-NE2	4.02	118.58	122.60	1	1
A	153	GLN	OE1-CD-NE2	4.01	118.59	122.60	1	1
A	64	GLN	OE1-CD-NE2	4.01	118.59	122.60	1	1

### Too-close contacts

The following all-atom clashscore is based on a MolProbity analysis. All-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The table below contains clashscores for all atomic models in this entry.

Model ID	Clash score	Number of clashes
1	0.56	7

There are 7 clashes. The table below contains the detailed list of all clashes based on a MolProbity analysis. Bad clashes are  $\geq 0.4$  Angstrom.

Atom 1	Atom 2	Clash(Å)	Model ID (Worst)	Models (Total)
B:298:VAL:HG22	B:415:MET:HE1	0.54	1	1
B:273:VAL:HG11	B:385:ILE:HD11	0.47	1	1

Atom 1	Atom 2	Clash(Å)	Model ID (Worst)	Models (Total)
A:44:LEU:HD21	A:245:TYR:CZ	0.47	1	1
B:119:GLU:HB3	B:120:PRO:HD3	0.45	1	1
B:119:GLU:CB	B:120:PRO:HD3	0.45	1	1
B:243:GLU:O	B:429:LYS:HE3	0.43	1	1
B:94:ASP:C	B:96:LYS:H	0.41	1	1

### Torsion angles: Protein backbone ?

In the following table, Ramachandran outliers are listed. The Analysed column shows the number of residues for which the backbone conformation was analysed.

Model ID	Analysed	Favored	Allowed	Outliers
1	809	760	44	5

There are 5 unique backbone outliers. Detailed list of outliers are tabulated below.

Chain	Res	Type	Models (Total)
A	209	ILE	1
B	93	ASP	1
B	95	ALA	1
B	110	ASP	1
B	120	PRO	1

### Torsion angles : Protein sidechains ?

In the following table, sidechain rotameric outliers are listed. The Analysed column shows the number of residues for which the sidechain conformation was analysed.

Model ID	Analysed	Favored	Allowed	Outliers
1	656	627	19	10

There are 10 unique sidechain outliers. Detailed list of outliers are tabulated below.

Chain	Res	Type	Models (Total)
A	5	THR	1
B	97	THR	1
B	104	THR	1
B	111	VAL	1
B	114	GLU	1
B	119	GLU	1
B	124	THR	1
B	190	THR	1
B	413	HIS	1

Chain	Res	Type	Models (Total)
B	440	MET	1

## 5. Fit to Data Used for Modeling Assessment ?

### 5.2. Crosslinking-MS ?

#### 5.2.1. Restraint types ?


This table summarizes information about crosslinker(s) used for data generation, and how crosslinking information was translated into actual modeling restraints. Restraints assigned "by-residue" are interpreted as between CA atoms. Restraints between coarse-grained beads are indicated as "coarse-grained". *Restraint group* represents a set of crosslinking restraints applied collectively in the modeling.

There are 7 crosslinking restraints combined in 7 restraint groups.

Linker	Residue 1	Atom 1	Residue 2	Atom 2	Restraint type	Distance, Å	Count
SDA	LYS	CA	LYS	CA	upper bound	25.00	5
SDA	ALA	CA	LYS	CA	upper bound	25.00	2

#### Distograms of individual restraints

Distograms (i.e., histogram plots of distances) provide an overview of distributions of distances between residues for which chemical crosslinks were identified. The shift of the distogram relative to the threshold value may indicate a poor model. Restraints with identical thresholds are grouped into one plot. Only the best distance per restraint per model group/ensemble is plotted. Inter- and intramolecular (including self-links) restraints are also grouped into one plot. Distance for a restraint between coarse-grained beads is calculated as a minimal distance between shells; if beads intersect, the distance will be reported as 0.0. A bead with the highest available resolution for a given residue is used for the assessment.




#### 5.2.2. Satisfaction of restraints ?

Satisfaction of restraints is calculated on a *restraint group* (a set of crosslinking restraints applied collectively in the modeling) level. Satisfaction of a restraint group depends on satisfaction of individual restraints in the group and the conditionality (all/any). A restraint group is considered satisfied, if the condition was met in at least one model of the model group/ensemble. The number of measured restraints can be smaller than the total number of restraint groups if crosslinks involve non-modeled residues. Only deposited models are used for validation right now.

State group	State	Model group	# of Deposited models/Total	Restraint group type	Satisfied (%)	Violated (%)	Count (Total=7)
1	1	1	1/1	All	42.86	57.14	7
				Heteromeric links/Intermolecular	42.86	57.14	7

Per-model satisfaction rates in ensembles

Every point represents one model in a model group/ensemble. Where possible, boxplots with quartile marks are also plotted.



## 6. Fit to Data Used for Validation Assessment

Validation for this section is under development.

### Acknowledgments

The development of integrative model validation metrics, implementation of a model validation pipeline, and creation of a validation report for integrative structures are funded by NSF awards to the [PDB-IHM team](#) (DBI-1756248, DBI-2112966, DBI-2112967, DBI-2112968, and DBI-1756250) and awards from NSF, NIH, and DOE to the [RCSB PDB](#) (DBI-2321666, R01GM157729, and DE-SC0019749). The PDB-IHM team and members of the [Sali lab](#) contributed model validation metrics and software packages.

Dr. Jill Trewhella, Dr. Dina Schneidman, and members of the [SASBDB](#) repository are acknowledged for their advice and support in implementing SAS validation methods. Team members from the labs of Dr. Juri Rappsilber; Dr. Alexander Leitner; Dr. Andrea Graziadei, and members of [PRIDE](#) database are acknowledged for their advice and support in implementing crosslinking-MS validation methods. We are grateful to Dr. Shruthi Viswanath for discussions about uncertainty assessment of integrative structural models.

Members of the [wwPDB Integrative/Hybrid Methods Task Force](#) provided recommendations and community support for the project.