

Summary of integrative structure determination of Complex of the closed conformation of P. furiosus Mre11-Rad50 bound to dsDNA (PDB ID: 9A9N | pdb_00009a9n)

1. Model Composition	
1.1. Entry composition	<ul style="list-style-type: none"> - DNA double-strand break repair Rad50 ATPase: chain(s) A, D (192 residues) - DNA double-strand break repair Rad50 ATPase: chain(s) B, E (167 residues) - DNA double-strand break repair protein Mre11: chain(s) C, F (32 residues) - DNA double-strand break repair protein Mre11: chain(s) G, H (333 residues) - DNA (36-MER): chain(s) I, J (36 residues)
1.2. Datasets used for modeling	<ul style="list-style-type: none"> - NMR data, Other: bmrbig115 - Integrative model, PDB: pdb_00009a9j - De Novo model, Zenodo: 10.5281/zenodo.15065939
2. Representation	
2.1. Number of representations	1
2.2. Scale	Atomic
2.3. Number of rigid and flexible segments	0, 10
3. Restraints	
3.1. Physical principles	Information about physical principles was not provided

<p>3.2. Experimental data</p>	<ul style="list-style-type: none"> - 2 unique DerivedDistanceRestraint: Lower Upper Bound Distance: 18.0-44.5 - 2 unique DerivedDistanceRestraint: Lower Upper Bound Distance: 17.2-42.6 - 4 unique DerivedDistanceRestraint: Lower Upper Bound Distance: 17.6-43.7 - 2 unique DerivedDistanceRestraint: Lower Upper Bound Distance: 19.3-48.0 - 2 unique DerivedDistanceRestraint: Lower Upper Bound Distance: 17.3-42.8 - 2 unique DerivedDistanceRestraint: Lower Upper Bound Distance: 18.5-45.8 - 2 unique DerivedDistanceRestraint: Lower Upper Bound Distance: 19.4-48.2 - 2 unique DerivedDistanceRestraint: Lower Upper Bound Distance: 19.9-49.3 - 2 unique DerivedDistanceRestraint: Lower Upper Bound Distance: 16.5-40.8 - 4 unique DerivedDistanceRestraint: Lower Upper Bound Distance: 16.9-41.8 - 2 unique DerivedDistanceRestraint: Lower Upper Bound Distance: 17.1-42.3 - 2 unique DerivedDistanceRestraint: Lower Upper Bound Distance: 15.9-39.5 - 4 unique DerivedDistanceRestraint: Lower Upper Bound Distance: 17.2-42.5 - 6 unique DerivedDistanceRestraint: Lower Upper Bound Distance: 17.8-44.2 - 2 unique DerivedDistanceRestraint: Lower Upper Bound Distance: 16.1-39.9 - 2 unique DerivedDistanceRestraint: Lower Upper Bound Distance: 16.3-40.5 - 2 unique DerivedDistanceRestraint: Lower Upper Bound Distance: 16.1-39.8 - 2 unique DerivedDistanceRestraint: Lower Upper Bound Distance: 15.5-38.5 - 2 unique DerivedDistanceRestraint: Lower Upper Bound Distance: 15.7-39.0 - 4 unique DerivedDistanceRestraint: Lower Upper Bound Distance: 17.0-42.2 - 4 unique DerivedDistanceRestraint: Lower Upper Bound Distance: 20.2-49.9 - 4 unique DerivedDistanceRestraint: Lower Upper Bound Distance: 20.2-50.0 - 4 unique DerivedDistanceRestraint: Lower Upper Bound Distance: 18.4-45.6 - 2 unique DerivedDistanceRestraint: Lower Upper Bound Distance: 19.1-47.5 - 4 unique DerivedDistanceRestraint: Lower Upper Bound Distance: 18.7-46.3 - 4 unique DerivedDistanceRestraint: Lower Upper Bound Distance: 19.1-47.3 - 4 unique DerivedDistanceRestraint: Lower Upper Bound Distance: 19.6-48.6 - 4 unique DerivedDistanceRestraint: Lower Upper Bound Distance: 20.4-50.7 - 4 unique DerivedDistanceRestraint: Lower Upper Bound Distance: 19.4-48.0
<p>4. Validation</p>	
<p>4.2. Number of ensembles</p>	<p>0</p>

4.3. Number of models in ensembles	Not applicable
4.4. Number of deposited models	1
4.5. Model precision	Not available
4.6. Data quality	Data quality has not been assessed
4.7. Model quality: assessment of atomic segments	<ul style="list-style-type: none"> - Clashscore: 7.04 - Ramachandran outliers: 15 - Sidechain outliers: 79
4.8. Fit to data used for modeling	Fit of model to information used to compute it has not been determined
4.9. Fit to data used for validation	Fit of model to information not used to compute it has not been determined
5. Methodology and Software	
1. 5.1. Method name	Not available
5.3. Method description	<p>This model was calculated using NMR paramagnetic relaxation enhancement (PRE) data as distance restraints in the 'Guru' and 'multi-body' interface of the HADDOCK v2.4 webserver. The starting structural model of the P. furiosus Closed MR(NBD) complex (PDBIHM id 9A9J) was obtained from previous HADDOCK calculations using LRET measurements as distance restraints to dock a Mre11 dimer and two Rad50 monomers (https://doi.org/10.7554/elife.69579). Two hairpin dsDNA ensembles were used as the starting structural models for the DNA. When HADDOCK performed the docking simulations, one of the dsDNA structures from the ensemble of 20 was randomly selected for each run. During the rigid body energy minimization, 3000 structures were calculated, and the 200 best solutions based on the inter-molecular energy were selected for the semiflexible simulated annealing followed by explicit water refinement.</p>
5.5. Software	HADDOCK (version 2.4)