



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 9, 2026 – 02:22 PM EDT

PDB ID : 9CRT / pdb_00009crt
Title : Crystal structure of IgG1 FC at natural pH
Authors : Reddem, E.R.; Shapiro, L.
Deposited on : 2024-07-22
Resolution : 2.19 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.48.1


1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.19 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	164625	5791 (2.20-2.20)
Clashscore	180529	6634 (2.20-2.20)
Ramachandran outliers	177936	6560 (2.20-2.20)
Sidechain outliers	177891	6561 (2.20-2.20)
RSRZ outliers	164620	5791 (2.20-2.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	231	
1	B	231	
2	C	9	
3	D	9	

2 Entry composition [i](#)


There are 4 unique types of molecules in this entry. The entry contains 3676 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called IgG1 FC,Immunoglobulin gamma-1 heavy chain.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	207	Total 1660	C 1057	N 279	O 318	S 6	0	0	0
1	B	207	Total 1660	C 1057	N 279	O 318	S 6	0	0	0

- Molecule 2 is an oligosaccharide called beta-D-galactopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-2)-alpha-D-mannopyranose-(1-6)-[2-acetamido-2-deoxy-beta-D-glucopyranose-(1-2)-alpha-D-mannopyranose-(1-3)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-[alpha-L-fucopyranose-(1-6)]2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
2	C	9	Total 110	C 62	N 4	O 44	0	0	0

- Molecule 3 is an oligosaccharide called alpha-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-2)-alpha-D-mannopyranose-(1-6)-[2-acetamido-2-deoxy-beta-D-glucopyranose-(1-2)-alpha-D-mannopyranose-(1-3)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-[alpha-L-fucopyranose-(1-6)]2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
3	D	9	Total 110	C 62	N 4	O 44	0	0	0


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	83	Total 83	O 83	0	0
4	B	53	Total 53	O 53	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: IgG1 FC, Immunoglobulin gamma-1 heavy chain



- Molecule 1: IgG1 FC, Immunoglobulin gamma-1 heavy chain



- Molecule 2: beta-D-galactopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-2)-alpha-D-mannopyranose-(1-6)-[2-acetamido-2-deoxy-beta-D-glucopyranose-(1-2)-alpha-D-mannopyranose-(1-3)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-[alpha-L-fucopyranose-(1-6)]2-acetamido-2-deoxy-beta-D-glucopyranose



- Molecule 3: alpha-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-2)-alpha-D-mannopyranose-(1-6)-[2-acetamido-2-deoxy-beta-D-glucopyranose-(1-2)-alpha-D-mannopyr

anose-(1-3)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-[alpha-L-fucopyranose-(1-6)]2-acetamido-2-deoxy-beta-D-glucopyranose

Chain D: 22% 56% 22%

MAG1	MAG2	BMA3	MAN4	NAG5	MAN6	MAN7	MAG8	FUC9
------	------	------	------	------	------	------	------	------

4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	49.64Å 80.67Å 136.34Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	33.52 – 2.19 33.52 – 2.19	Depositor EDS
% Data completeness (in resolution range)	99.9 (33.52-2.19) 99.9 (33.52-2.19)	Depositor EDS
R_{merge}	0.15	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.53 (at 2.20Å)	Xtrriage
Refinement program	PHENIX 1.20.1_4487	Depositor
R, R_{free}	0.222 , 0.282 0.223 , 0.281	Depositor DCC
R_{free} test set	2876 reflections (9.93%)	wwPDB-VP
Wilson B-factor (Å ²)	44.8	Xtrriage
Anisotropy	0.392	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.29 , 25.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	3676	wwPDB-VP
Average B, all atoms (Å ²)	55.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.52% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: FUC, MAN, NAG, GAL, BMA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.40	0/1706	0.54	0/2324
1	B	0.42	0/1706	0.66	1/2324 (0.0%)
All	All	0.41	0/3412	0.60	1/4648 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	B	0	1

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
1	B	286	ASN	CB-CA-C	-5.31	104.46	111.74

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	B	292	ARG	Sidechain

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within

the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1660	0	1628	14	0
1	B	1660	0	1628	31	0
2	C	110	0	94	2	0
3	D	110	0	94	4	0
4	A	83	0	0	3	0
4	B	53	0	0	3	0
All	All	3676	0	3444	45	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (45) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:260:THR:HG22	1:B:305:VAL:HG12	1.61	0.83
1:B:274:LYS:HZ3	1:B:276:ASN:HB2	1.44	0.82
1:B:290:LYS:HG2	1:B:303:VAL:HG22	1.76	0.68
1:B:274:LYS:HD2	1:B:275:PHE:N	2.12	0.65
1:B:290:LYS:HE2	1:B:303:VAL:HG13	1.78	0.65
1:B:274:LYS:NZ	1:B:276:ASN:HB2	2.13	0.64
1:A:356:ASP:OD2	4:A:501:HOH:O	2.16	0.63
1:A:295:GLN:HE21	2:C:1:NAG:H62	1.63	0.61
1:A:297:ASN:HD22	2:C:1:NAG:H83	1.69	0.56
1:B:265:ASP:OD1	3:D:1:NAG:N2	2.38	0.56
1:B:278:TYR:HB2	1:B:320:LYS:HB3	1.86	0.56
1:A:360:LYS:O	1:A:414:LYS:HD3	2.05	0.55
1:B:397:VAL:HG22	1:B:405:PHE:CE1	2.42	0.55
1:A:361:ASN:OD1	1:A:361:ASN:N	2.33	0.54
1:B:312:ASP:HB3	1:B:319:TYR:OH	2.09	0.53
1:A:269:GLU:OE1	1:A:269:GLU:N	2.35	0.53
1:B:253:ILE:HG22	4:B:523:HOH:O	2.09	0.52
1:B:322:LYS:HG3	1:B:333:GLU:HG2	1.93	0.51
1:B:402:GLY:O	4:B:501:HOH:O	2.19	0.51
1:B:297:ASN:OD1	1:B:299:THR:HG22	2.12	0.50
1:A:353:PRO:HG3	1:A:363:VAL:CG1	2.41	0.50
1:B:274:LYS:NZ	4:B:504:HOH:O	2.39	0.49
1:B:259:VAL:HG23	1:B:308:VAL:HG11	1.94	0.49
1:A:244:PRO:HB3	1:A:336:ILE:HD11	1.95	0.49
1:A:347:GLN:NE2	1:A:370:LYS:HE3	2.28	0.48
1:B:358:LEU:O	1:B:414:LYS:NZ	2.25	0.48

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:342:GLN:OE1	4:A:502:HOH:O	2.19	0.48
1:B:351:LEU:HB2	1:B:366:THR:HB	1.96	0.47
1:A:421:ASN:OD1	4:A:503:HOH:O	2.20	0.47
1:B:301:ARG:NH1	3:D:2:NAG:H81	2.31	0.46
1:B:246:LYS:HE3	3:D:6:MAN:H61	1.98	0.45
1:B:274:LYS:HD2	1:B:275:PHE:H	1.82	0.45
1:A:355:ARG:H	1:A:355:ARG:HG2	1.48	0.44
1:B:260:THR:HG21	3:D:6:MAN:H2	1.99	0.44
1:B:353:PRO:HD3	1:B:365:LEU:HD12	2.00	0.43
1:B:266:VAL:N	1:B:299:THR:OG1	2.52	0.43
1:A:266:VAL:HB	1:A:300:TYR:HB2	2.00	0.43
1:B:248:LYS:NZ	1:B:380:GLU:OE2	2.40	0.43
1:B:268:HIS:HE1	1:B:298:SER:C	2.27	0.43
1:B:286:ASN:OD1	1:B:286:ASN:N	2.52	0.42
1:B:274:LYS:HB3	1:B:324:SER:HB2	2.01	0.42
1:B:295:GLN:HB2	1:B:297:ASN:OD1	2.20	0.42
1:A:367:CYS:HB2	1:A:381:TRP:CZ2	2.55	0.42
1:B:357:GLU:OE1	1:B:364:SER:HB2	2.20	0.41
1:B:238:PRO:HB2	1:B:328:LEU:HD11	2.03	0.41

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	205/231 (89%)	202 (98%)	3 (2%)	0	100	100
1	B	205/231 (89%)	191 (93%)	14 (7%)	0	100	100
All	All	410/462 (89%)	393 (96%)	17 (4%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	194/211 (92%)	192 (99%)	2 (1%)	73	84
1	B	194/211 (92%)	183 (94%)	11 (6%)	17	21
All	All	388/422 (92%)	375 (97%)	13 (3%)	32	42

All (13) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	361	ASN
1	A	443	LEU
1	B	262	VAL
1	B	263	VAL
1	B	282	VAL
1	B	286	ASN
1	B	292	ARG
1	B	293	GLU
1	B	302	VAL
1	B	303	VAL
1	B	363	VAL
1	B	397	VAL
1	B	424	SER

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	295	GLN
1	A	311	GLN
1	A	347	GLN
1	B	347	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

18 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	NAG	C	1	2,1	14,14,15	0.59	0	17,19,21	0.50	0
2	NAG	C	2	2	14,14,15	0.24	0	17,19,21	0.61	0
2	BMA	C	3	2	11,11,12	1.51	2 (18%)	15,15,17	1.03	1 (6%)
2	MAN	C	4	2	11,11,12	1.08	1 (9%)	15,15,17	1.46	2 (13%)
2	NAG	C	5	2	14,14,15	0.33	0	17,19,21	0.47	0
2	GAL	C	6	2	11,11,12	1.59	1 (9%)	15,15,17	1.06	1 (6%)
2	MAN	C	7	2	11,11,12	1.21	2 (18%)	15,15,17	1.48	1 (6%)
2	NAG	C	8	2	14,14,15	0.35	0	17,19,21	0.63	0
2	FUC	C	9	2	10,10,11	0.96	0	14,14,16	1.10	1 (7%)
3	NAG	D	1	3,1	14,14,15	0.82	1 (7%)	17,19,21	0.82	1 (5%)
3	NAG	D	2	3	14,14,15	0.51	0	17,19,21	0.46	0
3	BMA	D	3	3	11,11,12	0.87	1 (9%)	15,15,17	0.93	1 (6%)
3	MAN	D	4	3	11,11,12	0.84	0	15,15,17	1.21	2 (13%)
3	NAG	D	5	3	14,14,15	0.36	0	17,19,21	0.68	1 (5%)
3	MAN	D	6	3	11,11,12	1.64	1 (9%)	15,15,17	1.07	1 (6%)
3	MAN	D	7	3	11,11,12	0.98	1 (9%)	15,15,17	1.19	2 (13%)
3	NAG	D	8	3	14,14,15	0.27	0	17,19,21	0.59	0
3	FUC	D	9	3	10,10,11	0.86	0	14,14,16	0.92	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAG	C	1	2,1	-	2/6/23/26	0/1/1/1
2	NAG	C	2	2	-	0/6/23/26	0/1/1/1
2	BMA	C	3	2	-	0/2/19/22	0/1/1/1
2	MAN	C	4	2	-	0/2/19/22	0/1/1/1
2	NAG	C	5	2	-	0/6/23/26	0/1/1/1
2	GAL	C	6	2	-	0/2/19/22	0/1/1/1
2	MAN	C	7	2	-	1/2/19/22	0/1/1/1
2	NAG	C	8	2	-	4/6/23/26	0/1/1/1
2	FUC	C	9	2	-	-	0/1/1/1
3	NAG	D	1	3,1	-	2/6/23/26	0/1/1/1
3	NAG	D	2	3	-	2/6/23/26	0/1/1/1
3	BMA	D	3	3	-	0/2/19/22	0/1/1/1
3	MAN	D	4	3	-	2/2/19/22	0/1/1/1
3	NAG	D	5	3	-	0/6/23/26	0/1/1/1
3	MAN	D	6	3	-	2/2/19/22	1/1/1/1
3	MAN	D	7	3	-	2/2/19/22	0/1/1/1
3	NAG	D	8	3	-	4/6/23/26	0/1/1/1
3	FUC	D	9	3	-	-	0/1/1/1

All (10) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	C	6	GAL	C1-C2	4.17	1.62	1.52
2	C	3	BMA	C2-C3	3.69	1.58	1.52
3	D	6	MAN	O5-C5	3.28	1.49	1.43
2	C	4	MAN	O5-C5	2.60	1.48	1.43
3	D	7	MAN	O5-C5	2.42	1.48	1.43
2	C	3	BMA	O2-C2	-2.41	1.38	1.43
3	D	1	NAG	O5-C1	2.37	1.47	1.43
2	C	7	MAN	C2-C3	-2.21	1.49	1.52
3	D	3	BMA	C2-C3	2.15	1.55	1.52
2	C	7	MAN	O2-C2	-2.02	1.39	1.43

All (14) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	C	7	MAN	O2-C2-C3	-4.80	100.20	110.15
2	C	4	MAN	C1-O5-C5	4.28	117.92	112.19
2	C	4	MAN	O2-C2-C3	-2.98	103.98	110.15
3	D	7	MAN	O2-C2-C3	-2.96	104.02	110.15
3	D	4	MAN	C1-O5-C5	2.94	116.13	112.19

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	D	7	MAN	C1-O5-C5	2.82	115.97	112.19
2	C	6	GAL	O2-C2-C1	2.76	115.53	109.22
3	D	1	NAG	C1-O5-C5	2.67	115.76	112.19
3	D	6	MAN	C1-O5-C5	2.39	115.38	112.19
3	D	4	MAN	O2-C2-C3	-2.32	105.35	110.15
2	C	3	BMA	C1-O5-C5	2.11	115.02	112.19
2	C	9	FUC	C2-C3-C4	-2.04	107.27	110.86
3	D	3	BMA	C1-O5-C5	2.03	114.91	112.19
3	D	5	NAG	C1-O5-C5	2.01	114.87	112.19

There are no chirality outliers.

All (21) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	C	8	NAG	O5-C5-C6-O6
3	D	2	NAG	O5-C5-C6-O6
2	C	8	NAG	C4-C5-C6-O6
3	D	2	NAG	C4-C5-C6-O6
2	C	1	NAG	C8-C7-N2-C2
2	C	1	NAG	O7-C7-N2-C2
3	D	1	NAG	O5-C5-C6-O6
3	D	4	MAN	O5-C5-C6-O6
3	D	1	NAG	C4-C5-C6-O6
3	D	7	MAN	C4-C5-C6-O6
3	D	8	NAG	C4-C5-C6-O6
3	D	8	NAG	O5-C5-C6-O6
3	D	6	MAN	C4-C5-C6-O6
2	C	8	NAG	C1-C2-N2-C7
3	D	8	NAG	C1-C2-N2-C7
3	D	7	MAN	O5-C5-C6-O6
3	D	4	MAN	C4-C5-C6-O6
2	C	8	NAG	C3-C2-N2-C7
3	D	8	NAG	C3-C2-N2-C7
3	D	6	MAN	O5-C5-C6-O6
2	C	7	MAN	C4-C5-C6-O6


All (1) ring outliers are listed below:


Mol	Chain	Res	Type	Atoms
3	D	6	MAN	C1-C2-C3-C4-C5-O5

4 monomers are involved in 6 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	C	1	NAG	2	0
3	D	1	NAG	1	0
3	D	6	MAN	2	0
3	D	2	NAG	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.





5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	207/231 (89%)	0.27	3 (1%) 73 70	33, 46, 67, 76	0
1	B	207/231 (89%)	1.16	53 (25%) 2 1	30, 56, 100, 108	0
All	All	414/462 (89%)	0.72	56 (13%) 8 6	30, 49, 96, 108	0

All (56) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	273	VAL	5.6
1	B	275	PHE	5.1
1	B	323	VAL	5.0
1	B	238	PRO	4.7
1	B	300	TYR	4.5
1	B	284	VAL	4.2
1	B	263	VAL	4.2
1	B	266	VAL	4.2
1	B	277	TRP	4.0
1	B	321	CYS	3.9
1	B	291	PRO	3.8
1	B	271	PRO	3.8
1	B	262	VAL	3.7
1	B	292	ARG	3.6
1	B	303	VAL	3.6
1	B	287	ALA	3.5
1	B	296	TYR	3.4
1	B	302	VAL	3.4
1	B	328	LEU	3.3
1	B	331	PRO	3.2
1	B	327	ALA	3.2
1	B	272	GLU	3.2
1	B	285	HIS	3.2
1	B	306	LEU	3.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	319	TYR	3.0
1	B	274	LYS	3.0
1	B	261	CYS	2.9
1	B	359	THR	2.9
1	B	240	VAL	2.9
1	B	299	THR	2.9
1	B	332	ILE	2.7
1	B	324	SER	2.7
1	B	282	VAL	2.6
1	B	304	SER	2.6
1	B	264	VAL	2.5
1	B	239	SER	2.5
1	B	286	ASN	2.5
1	B	268	HIS	2.5
1	B	305	VAL	2.5
1	B	267	SER	2.4
1	B	288	LYS	2.3
1	B	289	THR	2.3
1	A	443	LEU	2.3
1	B	329	PRO	2.2
1	B	322	LYS	2.2
1	B	355	ARG	2.2
1	A	438	GLN	2.2
1	B	290	LYS	2.2
1	B	363	VAL	2.2
1	B	389	ASN	2.1
1	B	301	ARG	2.1
1	B	260	THR	2.1
1	B	294	GLU	2.0
1	B	444	SER	2.0
1	A	238	PRO	2.0
1	B	330	ALA	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum,




median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	NAG	D	8	14/15	0.47	0.16	89,96,105,109	0
3	FUC	D	9	10/11	0.50	0.16	107,111,113,113	0
3	NAG	D	2	14/15	0.56	0.17	96,101,105,105	0
2	NAG	C	8	14/15	0.57	0.15	75,84,86,89	0
3	MAN	D	4	11/12	0.66	0.13	81,83,90,92	0
3	NAG	D	1	14/15	0.67	0.14	103,106,110,111	0
3	MAN	D	6	11/12	0.69	0.17	61,69,78,79	0
3	BMA	D	3	11/12	0.75	0.12	81,85,90,93	0
3	NAG	D	5	14/15	0.80	0.12	73,74,80,81	0
3	MAN	D	7	11/12	0.80	0.10	72,83,86,88	0
2	GAL	C	6	11/12	0.83	0.20	49,56,60,64	0
2	MAN	C	7	11/12	0.86	0.09	62,65,74,76	0
2	NAG	C	1	14/15	0.88	0.09	48,52,56,60	0
2	MAN	C	4	11/12	0.89	0.09	47,49,50,52	0
2	FUC	C	9	10/11	0.90	0.09	49,53,56,61	0
2	NAG	C	2	14/15	0.91	0.09	45,48,53,54	0
2	BMA	C	3	11/12	0.91	0.08	46,51,56,58	0
2	NAG	C	5	14/15	0.92	0.08	43,52,54,54	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around Chain C:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.