



Full wwPDB X-ray Structure Validation Report ⓘ

Jun 30, 2026 – 04:08 AM EDT

PDB ID : 9ELC / pdb_00009elc
Title : Structure of glucocerebrosidase in complex with a covalent inhibitor
Authors : Pluvinage, B.; Boraston, A.B.
Deposited on : 2024-12-04
Resolution : 2.20 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtrriage (Phenix) : 2.0
EDS : 3.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.015 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.50


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	6164 (2.20-2.20)
Clashscore	190562	6851 (2.20-2.20)
Ramachandran outliers	187476	6768 (2.20-2.20)
Sidechain outliers	187428	6769 (2.20-2.20)
RSRZ outliers	180081	6166 (2.20-2.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	497	 3% 92% 8%
1	B	497	 2% 92% 7%
2	E	3	 100%
3	C	2	 50% 50%
3	F	2	 100%

2 Entry composition i


There are 8 unique types of molecules in this entry. The entry contains 8786 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Glucosylceramidase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	497	Total 3926	C 2525	N 675	O 710	S 16	0	2	0
1	B	495	Total 3884	C 2494	N 667	O 707	S 16	0	2	0

- Molecule 2 is an oligosaccharide called 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-[alpha-L-fucopyranose-(1-6)]2-acetamido-2-deoxy-beta-D-glucopyranose.




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
2	E	3	Total 38	C 22	N 2	O 14	0	0	0

- Molecule 3 is an oligosaccharide called 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
3	F	2	Total 28	C 16	N 2	O 10	0	0	0
3	C	2	Total 28	C 16	N 2	O 10	0	0	0

- Molecule 4 is (1S,2R,3S,4S,5S)-5-(hexylamino)-4-(hexyloxy)cyclohexane-1,2,3-triol (CCD ID: A1BLE) (formula: C₁₈H₃₇NO₄) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	N	O		
4	A	1	17	12	1	4	0	0
4	B	1	14	9	1	4	0	0

- Molecule 5 is 2-acetamido-2-deoxy-beta-D-glucopyranose (CCD ID: NAG) (formula: $C_8H_{15}NO_6$).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	N	O		
5	A	1	14	8	1	5	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	N	O		
5	B	1	14	8	1	5	0	0

- Molecule 6 is SULFATE ION (CCD ID: SO4) (formula: O₄S).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	O	S		
6	A	1	5	4	1	0	0
6	A	1	5	4	1	0	0
6	A	1	5	4	1	0	0
6	A	1	5	4	1	0	0
6	A	1	5	4	1	0	0
6	A	1	5	4	1	0	0
6	B	1	5	4	1	0	0
6	B	1	5	4	1	0	0
6	B	1	5	4	1	0	0
6	B	1	5	4	1	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
6	B	1	Total	O	S	0	0
			5	4	1		
6	B	1	Total	O	S	0	0
			5	4	1		

- Molecule 7 is 1,2-ETHANEDIOL (CCD ID: EDO) (formula: C₂H₆O₂).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
7	A	1	Total	C	O	0	0
			4	2	2		
7	A	1	Total	C	O	0	0
			4	2	2		
7	A	1	Total	C	O	0	0
			4	2	2		
7	A	1	Total	C	O	0	0
			4	2	2		
7	A	1	Total	C	O	0	0
			4	2	2		
7	A	1	Total	C	O	0	0
			4	2	2		
7	A	1	Total	C	O	0	0
			4	2	2		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
7	A	1	Total C O 4 2 2	0	0
7	A	1	Total C O 4 2 2	0	0
7	A	1	Total C O 4 2 2	0	0
7	A	1	Total C O 4 2 2	0	0
7	B	1	Total C O 4 2 2	0	0
7	B	1	Total C O 4 2 2	0	0
7	B	1	Total C O 4 2 2	0	0
7	B	1	Total C O 4 2 2	0	0
7	B	1	Total C O 4 2 2	0	0
7	B	1	Total C O 4 2 2	0	0
7	B	1	Total C O 4 2 2	0	0
7	B	1	Total C O 4 2 2	0	0
7	B	1	Total C O 4 2 2	0	0
7	B	1	Total C O 4 2 2	0	0
7	B	1	Total C O 4 2 2	0	0


- Molecule 8 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
8	A	335	Total O 335 335	0	0
8	B	332	Total O 332 332	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Glucosylceramidase




- Molecule 1: Glucosylceramidase



- Molecule 2: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-[alpha-L-fucopyranose-(1-6)]2-acetamido-2-deoxy-beta-D-glucopyranose



- Molecule 3: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose



- Molecule 3: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain C: 


MAG1
MAG2

4 Data and refinement statistics

Property	Value	Source
Space group	C 2 2 21	Depositor
Cell constants a, b, c, α , β , γ	110.36Å 284.81Å 91.59Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	29.92 – 2.20 29.92 – 2.20	Depositor EDS
% Data completeness (in resolution range)	96.8 (29.92-2.20) 96.7 (29.92-2.20)	Depositor EDS
R_{merge}	0.09	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.57 (at 2.20Å)	Xtrriage
Refinement program	PHENIX 1.18_3845	Depositor
R, R_{free}	0.173 , 0.210 0.174 , 0.210	Depositor DCC
R_{free} test set	3632 reflections (4.94%)	wwPDB-VP
Wilson B-factor (Å ²)	32.7	Xtrriage
Anisotropy	0.087	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 36.2	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.34$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	8786	wwPDB-VP
Average B, all atoms (Å ²)	34.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.66% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: A1BLE, NAG, FUC, SO4, EDO

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.21	1/4046 (0.0%)	0.40	0/5520
1	B	0.21	1/4001 (0.0%)	0.40	2/5459 (0.0%)
All	All	0.21	2/8047 (0.0%)	0.40	2/10979 (0.0%)

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	340	GLU	CD-OE2	6.67	1.38	1.25
1	A	340	GLU	CD-OE2	6.57	1.37	1.25

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	340	GLU	CG-CD-OE2	5.66	131.41	118.40
1	B	340	GLU	OE1-CD-OE2	-5.29	110.21	122.90

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3926	0	3821	22	0
1	B	3884	0	3751	25	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	E	38	0	34	0	0
3	C	28	0	25	0	0
3	F	28	0	25	0	0
4	A	17	0	0	3	0
4	B	14	0	0	0	0
5	A	14	0	13	0	0
5	B	14	0	13	0	0
6	A	30	0	0	0	0
6	B	30	0	0	0	0
7	A	52	0	78	1	0
7	B	44	0	66	8	0
8	A	335	0	0	0	0
8	B	332	0	0	2	0
All	All	8786	0	7826	47	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (47) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:451[B]:HIS:HE2	7:B:909:EDO:H22	1.45	0.80
1:B:44:ARG:HH12	7:B:913:EDO:H21	1.50	0.76
1:B:418:TYR:HE2	7:B:914:EDO:H21	1.62	0.65
1:B:255:HIS:HB2	7:B:916:EDO:H22	1.78	0.64
1:B:262:ARG:HH22	7:B:919:EDO:H22	1.62	0.63
1:B:176:ALA:HB3	1:B:230:VAL:HG12	1.83	0.61
1:A:254:GLU:HG2	7:A:810:EDO:H12	1.84	0.59
1:A:162:HIS:O	1:A:166:GLN:HG3	2.03	0.59
1:A:287:LEU:HB3	1:A:291:TRP:CD1	2.39	0.58
1:A:1:ALA:HB2	1:A:27:ASP:OD1	2.07	0.55
1:A:312:TRP:HH2	1:A:367:ILE:HG13	1.74	0.52
1:A:339:SER:O	1:A:379:THR:OG1	2.28	0.52
1:A:284:GLN:NE2	1:A:314:LEU:H	2.08	0.51
1:B:277:ARG:HH12	7:B:917:EDO:H11	1.76	0.50
1:B:363:TYR:HD2	1:B:420:LEU:HD22	1.77	0.49
1:A:426:PHE:HB3	1:A:493:LEU:HD21	1.95	0.49
1:B:287:LEU:HB3	1:B:291:TRP:CD1	2.49	0.48
1:B:127:ASP:HB3	1:B:246:PHE:CG	2.49	0.47
1:B:367:ILE:HD11	1:B:420:LEU:HD13	1.96	0.47
1:B:236:PRO:HB2	1:B:251:PHE:HB2	1.96	0.47

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:67:LEU:HG	1:A:470:LEU:HD11	1.97	0.47
1:A:284:GLN:HG3	1:A:314:LEU:HD12	1.97	0.47
1:B:496:ARG:HD3	8:B:1022:HOH:O	2.16	0.46
1:B:426:PHE:HB3	1:B:493:LEU:HD21	1.97	0.46
1:B:139:PRO:HA	1:B:184:TRP:CD1	2.51	0.46
1:A:82:GLY:HA3	1:A:118:ILE:O	2.15	0.45
1:A:194:LYS:HB2	1:A:242:SER:HA	1.98	0.45
1:A:313:TYR:HB2	4:A:801:A1BLE:C22	2.46	0.45
1:A:457:VAL:HG22	1:A:493:LEU:HD23	1.98	0.45
1:A:311:HIS:HB3	4:A:801:A1BLE:C22	2.47	0.44
1:A:199:GLY:HA3	1:A:203:ASP:OD2	2.17	0.44
1:B:387:PRO:HD3	1:B:404:VAL:O	2.16	0.44
1:A:387:PRO:HD3	1:A:404:VAL:O	2.18	0.44
1:B:40:TYR:HA	1:B:50:GLU:O	2.18	0.43
1:B:426:PHE:O	1:B:451[A]:HIS:NE2	2.50	0.43
1:B:152:GLU:HA	1:B:156:LEU:HD12	2.01	0.43
1:A:127:ASP:OD2	4:A:801:A1BLE:O11	2.36	0.43
1:B:188:ASN:HB3	7:B:918:EDO:H21	2.01	0.43
1:B:234:ASN:ND2	1:B:235:GLU:HG3	2.34	0.42
1:B:176:ALA:HB2	1:B:227:PHE:CE2	2.54	0.42
1:A:176:ALA:HB2	1:A:227:PHE:CE2	2.53	0.42
1:A:207:GLN:NE2	1:A:263:ASP:OD1	2.42	0.42
1:B:363:TYR:CD2	1:B:420:LEU:HD22	2.54	0.42
1:B:187:THR:OG1	7:B:918:EDO:H22	2.21	0.41
1:B:353:ARG:NH2	8:B:1009:HOH:O	2.46	0.41
1:A:176:ALA:HB3	1:A:230:VAL:HG12	2.02	0.41
1:A:321:LYS:HE2	1:A:321:LYS:HB2	1.78	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	497/497 (100%)	472 (95%)	22 (4%)	3 (1%)	21	23
1	B	493/497 (99%)	477 (97%)	13 (3%)	3 (1%)	21	23
All	All	990/994 (100%)	949 (96%)	35 (4%)	6 (1%)	21	23

All (6) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	233	GLU
1	A	281	LEU
1	A	381	TRP
1	B	281	LEU
1	B	381	TRP
1	B	233	GLU

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	420/424 (99%)	415 (99%)	5 (1%)	63	78
1	B	413/424 (97%)	410 (99%)	3 (1%)	76	87
All	All	833/848 (98%)	825 (99%)	8 (1%)	68	81

All (8) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	317	LEU
1	A	343	VAL
1	A	346	LYS
1	A	381	TRP
1	A	493	LEU
1	B	242	SER
1	B	381	TRP
1	B	420	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such

sidechains are listed below:

Mol	Chain	Res	Type
1	A	169	GLN
1	A	396	ASN
1	A	432	GLN
1	B	223	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

7 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	NAG	C	1	1,3	14,14,15	0.37	0	17,19,21	0.56	0
3	NAG	C	2	3	14,14,15	0.47	0	17,19,21	0.69	1 (5%)
2	NAG	E	1	1,2	14,14,15	0.45	0	17,19,21	0.39	0
2	NAG	E	2	2	14,14,15	0.49	0	17,19,21	0.52	0
2	FUC	E	3	2	10,10,11	0.83	0	14,14,16	0.81	0
3	NAG	F	1	1,3	14,14,15	0.30	0	17,19,21	0.43	0
3	NAG	F	2	3	14,14,15	0.41	0	17,19,21	0.62	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	NAG	C	1	1,3	-	2/6/23/26	0/1/1/1
3	NAG	C	2	3	-	2/6/23/26	0/1/1/1
2	NAG	E	1	1,2	-	0/6/23/26	0/1/1/1
2	NAG	E	2	2	-	0/6/23/26	0/1/1/1
2	FUC	E	3	2	-	-	0/1/1/1
3	NAG	F	1	1,3	-	0/6/23/26	0/1/1/1
3	NAG	F	2	3	-	0/6/23/26	0/1/1/1

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	C	2	NAG	C1-O5-C5	2.10	115.00	112.19

There are no chirality outliers.


All (4) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
3	C	1	NAG	C8-C7-N2-C2
3	C	1	NAG	O7-C7-N2-C2
3	C	2	NAG	C8-C7-N2-C2
3	C	2	NAG	O7-C7-N2-C2


There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.







5.6 Ligand geometry [i](#)

40 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
6	SO4	A	805	-	4,4,4	0.24	0	6,6,6	0.15	0
6	SO4	B	904	-	4,4,4	0.24	0	6,6,6	0.26	0
7	EDO	B	911	-	3,3,3	0.48	0	2,2,2	0.30	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
7	EDO	A	819	-	3,3,3	0.43	0	2,2,2	0.48	0
7	EDO	A	813	-	3,3,3	0.40	0	2,2,2	0.53	0
7	EDO	A	812	-	3,3,3	0.44	0	2,2,2	0.38	0
6	SO4	B	903	-	4,4,4	0.20	0	6,6,6	0.17	0
7	EDO	A	814	-	3,3,3	0.46	0	2,2,2	0.28	0
7	EDO	A	811	-	3,3,3	0.41	0	2,2,2	0.49	0
6	SO4	A	803	-	4,4,4	0.21	0	6,6,6	0.20	0
6	SO4	A	807	-	4,4,4	0.23	0	6,6,6	0.14	0
6	SO4	B	907	-	4,4,4	0.23	0	6,6,6	0.11	0
6	SO4	B	905	-	4,4,4	0.24	0	6,6,6	0.16	0
7	EDO	B	912	-	3,3,3	0.42	0	2,2,2	0.49	0
7	EDO	A	815	-	3,3,3	0.46	0	2,2,2	0.32	0
6	SO4	A	808	-	4,4,4	0.25	0	6,6,6	0.18	0
6	SO4	A	804	-	4,4,4	0.27	0	6,6,6	0.24	0
7	EDO	A	818	-	3,3,3	0.46	0	2,2,2	0.33	0
5	NAG	B	902	1	14,14,15	0.29	0	17,19,21	0.77	1 (5%)
6	SO4	A	806	-	4,4,4	0.23	0	6,6,6	0.20	0
4	A1BLE	B	901	1	14,14,23	2.09	3 (21%)	17,19,28	1.36	4 (23%)
7	EDO	B	910	-	3,3,3	0.50	0	2,2,2	0.34	0
6	SO4	B	908	-	4,4,4	0.25	0	6,6,6	0.05	0
7	EDO	B	914	-	3,3,3	0.40	0	2,2,2	0.54	0
7	EDO	A	816	-	3,3,3	0.42	0	2,2,2	0.44	0
7	EDO	B	915	-	3,3,3	0.46	0	2,2,2	0.35	0
7	EDO	B	913	-	3,3,3	0.38	0	2,2,2	0.60	0
7	EDO	A	820	-	3,3,3	0.40	0	2,2,2	0.53	0
7	EDO	A	809	-	3,3,3	0.41	0	2,2,2	0.44	0
7	EDO	A	817	-	3,3,3	0.40	0	2,2,2	0.63	0
7	EDO	B	916	-	3,3,3	0.48	0	2,2,2	0.30	0
7	EDO	A	821	-	3,3,3	0.41	0	2,2,2	0.42	0
7	EDO	B	918	-	3,3,3	0.48	0	2,2,2	0.23	0
7	EDO	B	919	-	3,3,3	0.42	0	2,2,2	0.46	0
7	EDO	B	909	-	3,3,3	0.38	0	2,2,2	0.56	0
7	EDO	B	917	-	3,3,3	0.42	0	2,2,2	0.49	0
7	EDO	A	810	-	3,3,3	0.37	0	2,2,2	0.54	0
4	A1BLE	A	801	1	17,17,23	2.09	4 (23%)	20,22,28	1.46	4 (20%)
5	NAG	A	802	1	14,14,15	0.28	0	17,19,21	0.51	0
6	SO4	B	906	-	4,4,4	0.23	0	6,6,6	0.20	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
7	EDO	B	911	-	-	0/1/1/1	-
7	EDO	A	819	-	-	0/1/1/1	-
7	EDO	A	813	-	-	0/1/1/1	-
7	EDO	A	812	-	-	0/1/1/1	-
7	EDO	A	814	-	-	1/1/1/1	-
7	EDO	A	811	-	-	0/1/1/1	-
7	EDO	B	912	-	-	0/1/1/1	-
7	EDO	A	815	-	-	0/1/1/1	-
7	EDO	A	818	-	-	0/1/1/1	-
5	NAG	B	902	1	-	0/6/23/26	0/1/1/1
4	A1BLE	B	901	1	-	2/5/25/34	0/1/1/1
7	EDO	B	910	-	-	0/1/1/1	-
7	EDO	B	914	-	-	0/1/1/1	-
7	EDO	A	816	-	-	0/1/1/1	-
7	EDO	B	915	-	-	0/1/1/1	-
7	EDO	B	913	-	-	1/1/1/1	-
7	EDO	A	820	-	-	0/1/1/1	-
7	EDO	A	809	-	-	0/1/1/1	-
7	EDO	A	817	-	-	0/1/1/1	-
7	EDO	B	916	-	-	0/1/1/1	-
7	EDO	A	821	-	-	0/1/1/1	-
7	EDO	B	918	-	-	1/1/1/1	-
7	EDO	B	919	-	-	0/1/1/1	-
7	EDO	B	909	-	-	0/1/1/1	-
7	EDO	B	917	-	-	0/1/1/1	-
7	EDO	A	810	-	-	1/1/1/1	-
4	A1BLE	A	801	1	-	6/8/28/34	0/1/1/1
5	NAG	A	802	1	-	0/6/23/26	0/1/1/1

All (7) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	801	A1BLE	C05-N07	4.65	1.57	1.47
4	B	901	A1BLE	C06-C05	4.51	1.58	1.52
4	B	901	A1BLE	C05-N07	4.34	1.56	1.47
4	A	801	A1BLE	C19-N07	4.30	1.56	1.47
4	A	801	A1BLE	C06-C05	3.60	1.57	1.52
4	A	801	A1BLE	C01-C02	2.05	1.57	1.52
4	B	901	A1BLE	C01-C06	2.02	1.57	1.52

All (9) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	801	A1BLE	C04-C05-C06	3.05	115.40	109.75
4	B	901	A1BLE	C02-C01-C06	2.88	116.23	109.68
4	A	801	A1BLE	C02-C01-C06	2.85	116.15	109.68
5	B	902	NAG	C1-O5-C5	2.75	115.87	112.19
4	A	801	A1BLE	O11-C01-C06	-2.66	103.12	109.94
4	B	901	A1BLE	C04-C05-C06	2.55	114.48	109.75
4	A	801	A1BLE	C04-C05-N07	-2.49	106.53	112.04
4	B	901	A1BLE	O10-C02-C01	-2.03	105.60	110.38
4	B	901	A1BLE	C04-C05-N07	-2.02	107.57	112.04

There are no chirality outliers.

All (12) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	A	801	A1BLE	C05-C06-O12-C17
4	B	901	A1BLE	C04-C05-N07-C19
4	B	901	A1BLE	C06-C05-N07-C19
4	A	801	A1BLE	C06-C05-N07-C19
7	A	810	EDO	O1-C1-C2-O2
4	A	801	A1BLE	C20-C21-C22-C23
4	A	801	A1BLE	C20-C19-N07-C05
4	A	801	A1BLE	C04-C05-N07-C19
7	B	918	EDO	O1-C1-C2-O2
7	B	913	EDO	O1-C1-C2-O2
4	A	801	A1BLE	C19-C20-C21-C22
7	A	814	EDO	O1-C1-C2-O2


There are no ring outliers.


9 monomers are involved in 12 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
7	B	914	EDO	1	0
7	B	913	EDO	1	0
7	B	916	EDO	1	0
7	B	918	EDO	2	0
7	B	919	EDO	1	0
7	B	909	EDO	1	0
7	B	917	EDO	1	0
7	A	810	EDO	1	0
4	A	801	A1BLE	3	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths,

bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [\(i\)](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [\(i\)](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	497/497 (100%)	-0.38	13 (2%) 57 54	16, 31, 47, 75	2 (0%)
1	B	495/497 (99%)	-0.37	12 (2%) 59 56	18, 31, 48, 82	2 (0%)
All	All	992/994 (99%)	-0.37	25 (2%) 58 55	16, 31, 47, 82	4 (0%)

All (25) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	344	GLY	5.9
1	A	343	VAL	5.0
1	B	313	TYR	4.7
1	B	316	PHE	4.5
1	B	312	TRP	4.1
1	B	319	PRO	4.0
1	A	316	PHE	3.7
1	A	378	TRP	3.6
1	B	347	PHE	3.5
1	B	345	SER	3.4
1	A	345	SER	2.9
1	A	313	TYR	2.8
1	A	397	PHE	2.8
1	B	314	LEU	2.8
1	A	318	ALA	2.8
1	A	31	PHE	2.8
1	A	406	ILE	2.7
1	B	344	GLY	2.7
1	B	31	PHE	2.5
1	A	347	PHE	2.4
1	A	312	TRP	2.4
1	A	317	LEU	2.2
1	B	315	ASP	2.2
1	B	348	TRP	2.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	1	ALA	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [\(i\)](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [\(i\)](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	NAG	E	2	14/15	0.52	0.17	50,59,70,76	0
3	NAG	C	2	14/15	0.74	0.13	56,63,72,74	0
2	FUC	E	3	10/11	-	-	44,57,65,70	0
3	NAG	F	1	14/15	-	-	29,37,42,45	0
3	NAG	F	2	14/15	-	-	36,49,66,70	0
2	NAG	E	1	14/15	0.80	0.12	37,47,54,55	0
3	NAG	C	1	14/15	0.91	0.10	29,47,58,63	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around Chain E:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around Chain F:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around Chain C:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.4 Ligands

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
7	EDO	A	811	4/4	0.59	0.24	46,53,58,59	0
5	NAG	B	902	14/15	0.66	0.18	43,62,66,71	0
4	A1BLE	A	801	17/23	0.68	0.20	38,49,60,62	0
7	EDO	B	917	4/4	0.74	0.23	46,48,52,58	0
7	EDO	B	916	4/4	0.75	0.20	46,48,50,51	0
7	EDO	A	819	4/4	0.75	0.15	40,42,44,47	0
7	EDO	B	919	4/4	0.77	0.18	55,57,63,63	0
7	EDO	A	812	4/4	0.78	0.15	44,50,52,55	0
7	EDO	A	821	4/4	0.79	0.13	54,60,66,67	0
7	EDO	B	918	4/4	0.81	0.16	39,47,51,54	0
7	EDO	A	818	4/4	0.81	0.15	49,49,53,54	0
7	EDO	A	809	4/4	0.83	0.20	42,45,49,52	0
7	EDO	B	911	4/4	0.83	0.17	38,42,46,48	0
7	EDO	A	817	4/4	0.84	0.30	31,40,48,52	0
7	EDO	A	820	4/4	0.84	0.16	47,50,53,56	0
6	SO4	A	808	5/5	0.85	0.22	57,61,66,67	0
7	EDO	B	909	4/4	0.85	0.14	39,44,47,57	0
7	EDO	A	814	4/4	0.85	0.16	47,50,52,59	0
7	EDO	B	913	4/4	0.85	0.16	39,46,47,53	0
7	EDO	A	810	4/4	0.86	0.16	30,41,44,51	0
7	EDO	B	915	4/4	0.86	0.18	46,49,51,53	0
7	EDO	B	914	4/4	0.87	0.15	34,38,41,48	0
4	A1BLE	B	901	14/23	0.87	0.13	32,42,48,52	0
7	EDO	A	816	4/4	0.87	0.16	44,49,51,53	0
5	NAG	A	802	14/15	0.88	0.09	32,38,42,44	0
6	SO4	B	908	5/5	0.88	0.15	63,65,74,76	0
7	EDO	B	912	4/4	0.89	0.11	35,36,43,44	0
7	EDO	B	910	4/4	0.89	0.13	32,34,35,42	0
6	SO4	B	906	5/5	0.91	0.10	54,59,67,75	0
6	SO4	A	807	5/5	0.91	0.18	59,64,64,89	0
6	SO4	A	805	5/5	0.91	0.10	56,62,69,77	0
6	SO4	B	907	5/5	0.92	0.15	57,62,68,88	0
6	SO4	A	806	5/5	0.93	0.10	44,47,58,68	0
7	EDO	A	815	4/4	0.94	0.08	35,36,41,41	0
6	SO4	B	905	5/5	0.94	0.08	43,48,54,57	0
7	EDO	A	813	4/4	0.95	0.10	28,29,34,48	0
6	SO4	A	804	5/5	0.97	0.06	38,39,42,44	0


Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
6	SO4	B	904	5/5	0.98	0.06	37,39,44,45	0
6	SO4	B	903	5/5	0.99	0.05	31,34,40,50	0
6	SO4	A	803	5/5	0.99	0.04	31,37,37,44	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.





6.5 Other polymers [i](#)

There are no such residues in this entry.