



Full wwPDB X-ray Structure Validation Report ⓘ

Dec 15, 2025 – 06:45 pm GMT

PDB ID : 9G63 / pdb_00009g63
Title : The structure of *Aspergillus fumigatus* UDP-GlcNAc pyrophosphorylase in complex with fragments
Authors : Yan, K.
Deposited on : 2024-07-17
Resolution : 2.03 Å (reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	1.8.4, CSD as541be (2020)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
buster-report	:	1.1.7 (2018)
Percentile statistics	:	20231227.v01 (using entries in the PDB archive December 27th 2023)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.47


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.03 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	164625	12358 (2.04-2.00)
Clashscore	180529	13897 (2.04-2.00)
Ramachandran outliers	177936	13770 (2.04-2.00)
Sidechain outliers	177891	13769 (2.04-2.00)
RSRZ outliers	164620	12358 (2.04-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	B	484	
2	A	484	

2 Entry composition

There are 7 unique types of molecules in this entry. The entry contains 7485 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called UDP-N-acetylglucosamine diphosphorylase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	B	471	Total	C	N	O	S	0	3	0
			3690	2356	629	692	13			

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	26	GLY	-	expression tag	UNP A0A229XUD0


- Molecule 2 is a protein called UDP-N-acetylglucosamine diphosphorylase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	A	460	Total	C	N	O	S	0	2	0
			3563	2275	607	667	14			

There is a discrepancy between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	26	GLY	-	expression tag	UNP A0A229XUD0

- Molecule 3 is 2-acetamido-2-deoxy-1-O-phosphono-alpha-D-glucopyranose (CCD ID: GN1) (formula: C₈H₁₆NO₉P).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	B	1	Total	C	N	O	P	0
			19	8	1	9	1	

- Molecule 4 is 4-bromanyl-2-methyl-pyrazol-3-amine (CCD ID: A1IHU) (formula: $C_4H_6BrN_3$) (labeled as "Ligand of Interest" by depositor).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	B	1	Total	Br	C	N	0	0
			8	1	4	3		
4	B	1	Total	Br	C	N	0	0
			8	1	4	3		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	B	1	Total	Br	C	N	0	0
			8	1	4	3		

- Molecule 5 is furan-2-yl(piperazin-1-yl)methanone (CCD ID: A1IIR) (formula: C₉H₁₂N₂O₂) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
5	B	1	Total	C	N	O	0	0
			13	9	2	2		

- Molecule 6 is CHLORIDE ION (CCD ID: CL) (formula: Cl).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	B	1	Total Cl	0	0
			1 1		

- Molecule 7 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
7	B	104	Total O	0	0
			104 104		
7	A	71	Total O	0	0
			71 71		


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: UDP-N-acetylglucosamine diphosphorylase



- Molecule 2: UDP-N-acetylglucosamine diphosphorylase



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	55.49Å 136.84Å 143.05Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	63.39 – 2.03 63.39 – 2.03	Depositor EDS
% Data completeness (in resolution range)	99.8 (63.39-2.03) 99.9 (63.39-2.03)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.06 (at 2.03Å)	Xtriage
Refinement program	REFMAC 5.8.0352	Depositor
R, R_{free}	0.213 , 0.264 0.221 , 0.269	Depositor DCC
R_{free} test set	1906 reflections (2.66%)	wwPDB-VP
Wilson B-factor (Å ²)	37.9	Xtriage
Anisotropy	0.321	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 37.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	0.006 for -h,l,k	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	7485	wwPDB-VP
Average B, all atoms (Å ²)	47.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.37% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: A1IHU, YCM, GN1, CL, A1IIR

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	B	0.64	0/3767	1.04	4/5086 (0.1%)
2	A	0.63	0/3643	1.06	2/4922 (0.0%)
All	All	0.63	0/7410	1.05	6/10008 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	B	0	3

There are no bond length outliers.

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	446	ASP	CA-CB-CG	6.56	119.16	112.60
1	B	506	GLU	CB-CG-CD	6.49	123.63	112.60
1	B	201	GLU	CB-CG-CD	5.70	122.29	112.60
1	B	249	ASN	CA-CB-CG	5.50	118.10	112.60
2	A	111	LYS	CB-CA-C	-5.28	101.87	110.85
2	A	64	HIS	CB-CA-C	5.12	119.56	110.85

There are no chirality outliers.

All (3) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	B	382	ARG	Sidechain
1	B	73[A]	ARG	Sidechain

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Group
1	B	73[B]	ARG	Sidechain

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	B	3690	0	3671	47	0
2	A	3563	0	3510	43	0
3	B	19	0	0	0	0
4	B	24	0	0	0	0
5	B	13	0	0	1	0
6	B	1	0	0	0	0
7	A	71	0	0	3	0
7	B	104	0	0	5	0
All	All	7485	0	7181	88	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (88) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:504:PHE:CZ	1:B:506:GLU:HG2	2.11	0.85
2:A:28:SER:O	2:A:30:GLU:N	2.10	0.84
1:B:101:ALA:HB2	1:B:419:LEU:HD13	1.68	0.75
2:A:137:GLY:HA3	2:A:196:THR:HG21	1.70	0.72
2:A:237:GLU:OE1	2:A:241:LYS:NZ	2.19	0.68
2:A:281[A]:CYS:SG	2:A:434:SER:HB3	2.34	0.68
1:B:307:ARG:HD3	7:B:800:HOH:O	1.95	0.67
1:B:227:CYS:HB3	1:B:385:ILE:HD13	1.76	0.65
2:A:230:ASN:ND2	2:A:382:ARG:HH12	1.95	0.65
2:A:374:HIS:HE1	7:A:640:HOH:O	1.80	0.64
2:A:141:ARG:HB3	2:A:142:LEU:HD23	1.79	0.63
1:B:266:ARG:NH1	1:B:367:GLU:OE2	2.32	0.62
1:B:309:ARG:HH11	1:B:309:ARG:CG	2.12	0.60
2:A:374:HIS:CE1	7:A:640:HOH:O	2.54	0.60

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:230:ASN:HD21	1:B:382:ARG:HH12	1.50	0.59
1:B:69:PHE:CE1	1:B:226:PRO:HD3	2.38	0.59
2:A:230:ASN:ND2	2:A:382:ARG:NH1	2.50	0.58
2:A:236:MET:O	2:A:387:CYS:HB2	2.03	0.58
1:B:69:PHE:HA	1:B:73[A]:ARG:HH22	1.69	0.58
2:A:334:ASP:N	2:A:334:ASP:OD1	2.39	0.56
2:A:379:HIS:CD2	2:A:409:PHE:CD2	2.95	0.55
1:B:189:TYR:CE1	1:B:218:ILE:HD13	2.41	0.55
2:A:138:GLN:HG3	2:A:140:THR:HG23	1.89	0.54
1:B:230:ASN:ND2	1:B:382:ARG:HH22	2.05	0.54
1:B:95[A]:GLU:OE2	1:B:350:LYS:NZ	2.39	0.54
1:B:154:GLY:O	1:B:451:LYS:HE2	2.07	0.54
2:A:385:ILE:HD12	2:A:403:GLY:HA3	1.89	0.54
1:B:504:PHE:CZ	1:B:506:GLU:CG	2.89	0.53
2:A:141:ARG:HG2	2:A:437:LYS:O	2.09	0.53
2:A:44:GLN:O	2:A:47:VAL:HG22	2.10	0.52
2:A:48:PHE:O	2:A:51:VAL:HG23	2.09	0.52
1:B:307:ARG:NH1	7:B:704:HOH:O	2.36	0.51
1:B:309:ARG:CG	1:B:309:ARG:NH1	2.73	0.51
1:B:101:ALA:HB2	1:B:419:LEU:CD1	2.38	0.51
1:B:273:ILE:N	1:B:273:ILE:HD12	2.26	0.50
1:B:80:LYS:O	1:B:84:PRO:HB3	2.12	0.50
2:A:52:ASP:O	2:A:53:GLU:HB3	2.12	0.50
2:A:125:ALA:HB1	2:A:180:SER:HB3	1.94	0.49
1:B:444:GLU:HA	1:B:449:THR:HG21	1.94	0.49
1:B:106:LEU:HD21	1:B:428:ARG:HG2	1.94	0.48
2:A:137:GLY:CA	2:A:196:THR:HG21	2.42	0.48
1:B:57:GLY:HA2	2:A:331:SER:HB3	1.95	0.48
1:B:506:GLU:O	1:B:507:LYS:C	2.56	0.48
1:B:309:ARG:HH11	1:B:309:ARG:HG3	1.78	0.47
1:B:82:LEU:C	1:B:84:PRO:HD3	2.40	0.46
2:A:142:LEU:HA	2:A:439:ALA:HB2	1.96	0.46
2:A:276:TYR:CE1	2:A:283:VAL:HG11	2.50	0.46
1:B:286:ALA:O	1:B:287:ASP:C	2.58	0.46
1:B:507:LYS:N	1:B:507:LYS:HE2	2.31	0.46
2:A:146:ALA:HB1	2:A:147:PRO:HD2	1.98	0.45
1:B:133:LEU:HD11	1:B:278:VAL:HB	1.98	0.45
1:B:384:LYS:HE3	7:B:799:HOH:O	2.17	0.45
1:B:57:GLY:HA2	2:A:331:SER:CB	2.47	0.45
1:B:308:LYS:HD2	1:B:317:LEU:HD21	1.99	0.45
1:B:105:ILE:HD11	7:B:770:HOH:O	2.17	0.45

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:A:142:LEU:HD21	2:A:148:LYS:HE3	1.99	0.45
2:A:52:ASP:O	2:A:53:GLU:CB	2.65	0.45
2:A:104:SER:O	2:A:108:SER:HB2	2.17	0.45
1:B:302:ALA:HA	1:B:423:ALA:O	2.17	0.44
1:B:70:ASP:HB3	1:B:73[A]:ARG:HG2	1.99	0.44
1:B:69:PHE:HA	1:B:73[A]:ARG:NH2	2.31	0.44
2:A:106:LEU:HD12	2:A:106:LEU:N	2.34	0.43
1:B:218:ILE:HD12	1:B:218:ILE:N	2.34	0.43
1:B:446:ASP:HB2	1:B:447:PRO:CD	2.50	0.42
2:A:39:TYR:CE2	2:A:66:LEU:HD22	2.54	0.42
2:A:197:ARG:HH11	2:A:221:GLU:HG3	1.84	0.42
1:B:179:ILE:HD12	1:B:179:ILE:O	2.19	0.42
2:A:484:ILE:HD11	2:A:507:LYS:HA	2.00	0.42
1:B:134:MET:HE2	1:B:275:THR:HG22	2.02	0.42
2:A:230:ASN:HD21	2:A:382:ARG:NH1	2.16	0.42
2:A:302:ALA:HA	2:A:423:ALA:O	2.20	0.41
2:A:62:LEU:HG	2:A:240:PHE:HA	2.03	0.41
2:A:277[B]:CYS:SG	2:A:279:ASP:OD2	2.78	0.41
1:B:419:LEU:HD22	1:B:419:LEU:HA	1.92	0.41
1:B:492:GLU:C	1:B:494:LEU:H	2.29	0.41
1:B:234:ILE:CG2	1:B:242:VAL:HG13	2.51	0.41
2:A:74:ILE:HD12	2:A:226:PRO:HD2	2.02	0.41
2:A:28:SER:C	2:A:30:GLU:N	2.79	0.41
2:A:169:ILE:CG1	2:A:285:VAL:HG11	2.51	0.41
2:A:476:VAL:O	2:A:476:VAL:HG13	2.21	0.41
2:A:109:ASP:OD1	2:A:111:LYS:HG2	2.21	0.40
2:A:198:LYS:N	2:A:199:PRO:CD	2.85	0.40
1:B:51:VAL:O	1:B:52:ASP:C	2.64	0.40
1:B:138:GLN:HE22	1:B:195:PRO:HG2	1.86	0.40
1:B:388:ILE:HD12	1:B:395:PHE:HA	2.03	0.40
1:B:320:GLN:HG2	7:B:783:HOH:O	2.22	0.40
1:B:407:GLU:HB2	5:B:605:A1IIR:C10	2.52	0.40
2:A:284:LYS:NZ	7:A:609:HOH:O	2.53	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	B	467/484 (96%)	453 (97%)	14 (3%)	0	100	100
2	A	452/484 (93%)	432 (96%)	14 (3%)	6 (1%)	10	5
All	All	919/968 (95%)	885 (96%)	28 (3%)	6 (1%)	19	14

All (6) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	A	53	GLU
2	A	89	ASP
2	A	29	ALA
2	A	55	GLN
2	A	136	GLY
2	A	345	GLN

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	B	393/407 (97%)	378 (96%)	15 (4%)	28	26
2	A	376/408 (92%)	358 (95%)	18 (5%)	21	18
All	All	769/815 (94%)	736 (96%)	33 (4%)	25	22

All (33) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	100	ILE
1	B	119	GLU
1	B	138	GLN
1	B	174	LEU
1	B	197	ARG
1	B	241	LYS
1	B	309	ARG
1	B	348	VAL
1	B	382	ARG
1	B	388	ILE
1	B	399	GLU
1	B	419	LEU
1	B	454	ILE
1	B	506	GLU
1	B	507	LYS
2	A	55	GLN
2	A	56	THR
2	A	59	ARG
2	A	62	LEU
2	A	80	LYS
2	A	142	LEU
2	A	145	SER
2	A	179	ILE
2	A	182	LYS
2	A	205	GLU
2	A	275	THR
2	A	297	LYS
2	A	310	ASN
2	A	329	GLU
2	A	334	ASP
2	A	337	THR
2	A	380	VAL
2	A	388	ILE

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (13) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	33	GLN
1	B	64	HIS
1	B	230	ASN
1	B	320	GLN
1	B	322	ASN
1	B	438	ASN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
2	A	34	GLN
2	A	55	GLN
2	A	61	GLN
2	A	83	ASN
2	A	222	GLN
2	A	230	ASN
2	A	310	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

1 non-standard protein/DNA/RNA residue is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
1	YCM	B	277	1	7,9,10	0.59	0	4,10,12	0.40	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	YCM	B	277	1	-	2/6/8/10	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (2) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
1	B	277	YCM	SG-CD-CE-NZ2
1	B	277	YCM	SG-CD-CE-OZ1

There are no ring outliers.

No monomer is involved in short contacts.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 6 ligands modelled in this entry, 1 is monoatomic - leaving 5 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	GN1	B	601	-	18,19,19	0.75	0	27,28,28	0.93	2 (7%)
4	A1IHU	B	603	-	6,8,8	2.20	2 (33%)	5,11,11	3.13	4 (80%)
4	A1IHU	B	604	-	6,8,8	2.14	2 (33%)	5,11,11	5.80	2 (40%)
5	A1IIR	B	605	-	10,14,14	3.73	4 (40%)	12,18,18	1.84	2 (16%)
4	A1IHU	B	602	-	6,8,8	3.23	3 (50%)	5,11,11	3.00	2 (40%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	GN1	B	601	-	-	0/10/31/31	0/1/1/1
4	A1IHU	B	603	-	-	-	0/1/1/1
4	A1IHU	B	604	-	-	-	0/1/1/1
5	A1IIR	B	605	-	-	0/4/16/16	1/2/2/2
4	A1IHU	B	602	-	-	-	0/1/1/1

All (11) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	B	605	A1IIR	C08-C07	-9.61	1.35	1.49
4	B	602	A1IHU	CAA-NAB	6.17	1.59	1.47
5	B	605	A1IIR	C04-N03	4.68	1.55	1.47
4	B	603	A1IHU	CAA-NAB	4.36	1.55	1.47
4	B	604	A1IHU	BRAG-CAD	4.19	1.99	1.89
4	B	602	A1IHU	CAD-CAC	-3.82	1.30	1.38
5	B	605	A1IIR	C02-N03	3.72	1.53	1.47
4	B	602	A1IHU	CAE-CAD	-2.96	1.31	1.37
4	B	603	A1IHU	CAD-CAC	-2.52	1.33	1.38
4	B	604	A1IHU	CAE-CAD	-2.33	1.32	1.37
5	B	605	A1IIR	C04-C05	2.06	1.56	1.51

All (12) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	B	604	A1IHU	CAA-NAB-CAC	9.27	139.99	128.91
4	B	604	A1IHU	CAE-NAF-NAB	8.92	113.51	104.23
4	B	602	A1IHU	CAE-NAF-NAB	6.01	110.49	104.23
4	B	603	A1IHU	CAE-NAF-NAB	5.65	110.11	104.23
5	B	605	A1IIR	C04-N03-C02	4.76	121.79	112.62
5	B	605	A1IIR	C02-N03-C07	-2.65	114.40	122.78
4	B	603	A1IHU	BRAG-CAD-CAE	2.63	128.66	124.51
4	B	603	A1IHU	CAE-CAD-CAC	2.42	110.21	107.16
3	B	601	GN1	O1-P-OP3	-2.34	100.36	109.39
4	B	603	A1IHU	CAA-NAB-CAC	-2.07	126.43	128.91
3	B	601	GN1	C4-C3-C2	-2.03	107.37	110.34
4	B	602	A1IHU	CAE-CAD-CAC	2.01	109.70	107.16

There are no chirality outliers.

There are no torsion outliers.


All (1) ring outliers are listed below:

Mol	Chain	Res	Type	Atoms
5	B	605	A1IIR	C01-C02-C04-C05-N03-N06


1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	B	605	A1IIR	1	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.




Ligand A1IHU B 603




Ligand A1IHU B 604



Ligand A1IIR B 605





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ > 2			OWAB(Å ²)	Q < 0.9
1	B	470/484 (97%)	0.05	14 (2%)	52	51	21, 40, 77, 92	3 (0%)
2	A	460/484 (95%)	0.33	22 (4%)	36	36	23, 48, 81, 115	2 (0%)
All	All	930/968 (96%)	0.19	36 (3%)	44	43	21, 44, 79, 115	5 (0%)

All (36) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	A	388	ILE	5.5
1	B	27	PRO	4.5
2	A	338	ALA	4.0
2	A	337	THR	3.9
2	A	333	ILE	3.9
2	A	476	VAL	3.9
1	B	476	VAL	3.3
1	B	471	THR	3.2
2	A	136	GLY	3.1
1	B	470	ILE	3.0
1	B	141	ARG	2.8
2	A	142	LEU	2.8
1	B	395	PHE	2.6
1	B	140	THR	2.6
1	B	52	ASP	2.5
2	A	347	ASP	2.5
2	A	385	ILE	2.5
2	A	140	THR	2.5
2	A	330	TYR	2.5
2	A	380	VAL	2.4
1	B	240	PHE	2.4
1	B	73[A]	ARG	2.4
2	A	310	ASN	2.4
2	A	335	LYS	2.4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	83	ASN	2.4
2	A	179	ILE	2.3
2	A	194	GLY	2.3
2	A	33	GLN	2.2
1	B	420	GLU	2.2
1	B	507	LYS	2.2
2	A	92	ALA	2.1
1	B	388	ILE	2.1
2	A	29	ALA	2.1
2	A	88	ASP	2.1
2	A	419	LEU	2.1
2	A	386	PRO	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
1	YCM	B	277	10/11	0.97	0.09	29,32,39,41	0

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
5	A1IIR	B	605	13/13	0.83	0.14	49,53,62,66	0
4	A1IHU	B	604	8/8	0.85	0.15	41,59,73,117	0
4	A1IHU	B	602	8/8	0.89	0.14	54,61,67,88	0
3	GN1	B	601	19/19	0.98	0.04	27,30,40,42	0
4	A1IHU	B	603	8/8	0.99	0.06	35,37,39,41	0

Continued on next page...




Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
6	CL	B	606	1/1	0.99	0.08	48,48,48,48	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around A1IIR B 605:

2mF_o-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around A1IHU B 604:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around A1IHU B 602:**


$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around GN1 B 601:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.5 Other polymers [i](#)

There are no such residues in this entry.