



# Full wwPDB X-ray Structure Validation Report i

Feb 4, 2026 – 12:10 pm GMT

PDB ID : 9GED / pdb\_00009ged  
Title : Entamoeba histolytica Gal/GalNAc lectin heavy chain residues 808-992  
Authors : Gerard, S.F.; Higgins, M.K.  
Deposited on : 2024-08-07  
Resolution : 2.48 Å (reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>  
with specific help available everywhere you see the i symbol.

The types of validation reports are described at  
<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references](#) i) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0  
Mogul : 1.8.4, CSD as541be (2020)  
Xtriage (Phenix) : 2.0  
EDS : 3.0  
Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)  
CCP4 : 9.0.010 (Gargrove)  
Density-Fitness : 1.0.12  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.47


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## X-RAY DIFFRACTION


The reported resolution of this entry is 2.48 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	164625	7106 (2.50-2.46)
Clashscore	180529	7991 (2.50-2.46)
Ramachandran outliers	177936	7888 (2.50-2.46)
Sidechain outliers	177891	7890 (2.50-2.46)
RSRZ outliers	164620	7106 (2.50-2.46)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.



## 2 Entry composition (i)

There are 3 unique types of molecules in this entry. The entry contains 4505 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Galactose/N-acetyl-D-galactosamine lectin heavy subunit 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	185	Total	C	N	O	S	0	0	0
			1491	933	241	302	15			
1	B	180	Total	C	N	O	S	0	0	0
			1457	914	235	293	15			
1	C	179	Total	C	N	O	S	0	0	0
			1450	909	234	292	15			

There are 9 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	811	SER	CYS	conflict	UNP P32022
A	972	SER	CYS	conflict	UNP P32022
A	993	GLY	THR	conflict	UNP P32022
B	811	SER	CYS	conflict	UNP P32022
B	972	SER	CYS	conflict	UNP P32022
B	993	GLY	THR	conflict	UNP P32022
C	811	SER	CYS	conflict	UNP P32022
C	972	SER	CYS	conflict	UNP P32022
C	993	GLY	THR	conflict	UNP P32022

- Molecule 2 is 2-acetamido-2-deoxy-beta-D-glucopyranose (CCD ID: NAG) (formula: C<sub>8</sub>H<sub>15</sub>NO<sub>6</sub>).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	1	Total C N O 14 8 1 5	0	0
2	B	1	Total C N O 14 8 1 5	0	0
2	C	1	Total C N O 14 8 1 5	0	0


- Molecule 3 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	18	Total O 18 18	0	0
3	B	23	Total O 23 23	0	0
3	C	24	Total O 24 24	0	0


### 3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Galactose/N-acetyl-D-galactosamine lectin heavy subunit 1



- Molecule 1: Galactose/N-acetyl-D-galactosamine lectin heavy subunit 1



- Molecule 1: Galactose/N-acetyl-D-galactosamine lectin heavy subunit 1



## 4 Data and refinement statistics i

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	83.14Å 145.08Å 85.94Å 90.00° 114.28° 90.00°	Depositor
Resolution (Å)	34.47 – 2.48 34.47 – 2.48	Depositor EDS
% Data completeness (in resolution range)	62.4 (34.47-2.48) 62.4 (34.47-2.48)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle^1$	1.53 (at 2.48Å)	Xtriage
Refinement program	BUSTER 2.10.4 (26-JUL-2023)	Depositor
$R$ , $R_{free}$	0.236 , 0.259 0.233 , 0.256	Depositor DCC
$R_{free}$ test set	995 reflections (3.04%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	44.7	Xtriage
Anisotropy	0.093	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.30 , 27.7	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.90	EDS
Total number of atoms	4505	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	53.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.58% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality i

### 5.1 Standard geometry i

Bond lengths and bond angles in the following residue types are not validated in this section: NAG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.68	0/1517	1.00	0/2045
1	B	0.64	0/1482	1.02	0/1998
1	C	0.62	0/1475	0.96	0/1988
All	All	0.65	0/4474	0.99	0/6031

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts i

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1491	0	1437	15	0
1	B	1457	0	1404	10	0
1	C	1450	0	1395	7	0
2	A	14	0	13	0	0
2	B	14	0	13	0	0
2	C	14	0	13	0	0
3	A	18	0	0	0	0
3	B	23	0	0	0	0
3	C	24	0	0	0	0
All	All	4505	0	4275	31	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (31) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:929:ILE:HD12	1:B:934:LYS:HB2	1.69	0.74
1:A:981:GLU:HG2	1:A:986:LYS:HG2	1.70	0.72
1:B:981:GLU:HG2	1:B:986:LYS:HG2	1.71	0.72
1:C:981:GLU:HG2	1:C:986:LYS:HG2	1.73	0.69
1:A:811:SER:HB2	1:A:818:GLU:H	1.64	0.60
1:A:835:LYS:NZ	1:A:933:GLU:HG2	2.17	0.59
1:A:858:CYS:HB2	1:A:881:ALA:HB1	1.85	0.58
1:B:929:ILE:HB	1:B:931:THR:HG22	1.90	0.54
1:C:858:CYS:HB2	1:C:881:ALA:HB1	1.90	0.52
1:B:929:ILE:HG22	1:B:930:GLU:N	2.25	0.51
1:A:872:TYR:HB3	1:A:879:VAL:HG11	1.93	0.51
1:A:929:ILE:HG12	1:A:936:GLN:HG2	1.92	0.50
1:A:858:CYS:HB3	1:A:916:TYR:HE1	1.77	0.50
1:A:858:CYS:HB3	1:A:916:TYR:CE1	2.46	0.50
1:C:835:LYS:NZ	1:C:933:GLU:HG2	2.28	0.48
1:C:872:TYR:HB3	1:C:879:VAL:HG11	1.94	0.48
1:A:833:ASP:HA	1:A:928:LEU:HD11	1.97	0.47
1:A:982:LYS:HG2	1:B:880:THR:HG23	1.96	0.47
1:B:872:TYR:HB3	1:B:879:VAL:HG11	1.97	0.47
1:B:929:ILE:HG22	1:B:930:GLU:H	1.81	0.46
1:A:835:LYS:HZ2	1:A:933:GLU:HG2	1.80	0.45
1:A:974:ILE:HD12	1:A:974:ILE:HA	1.85	0.44
1:C:928:LEU:HD13	1:C:932:ARG:HD2	2.00	0.43
1:B:858:CYS:HB2	1:B:881:ALA:HB1	2.00	0.42
1:C:875:CYS:HB3	1:C:926:CYS:HB2	1.98	0.42
1:A:884:HIS:O	1:A:912:ALA:HA	2.21	0.41
1:C:884:HIS:O	1:C:912:ALA:HA	2.21	0.41
1:B:978:VAL:HG23	1:B:989:THR:HG23	2.03	0.41
1:B:884:HIS:O	1:B:912:ALA:HA	2.20	0.40
1:A:862:ASP:OD1	1:A:864:ASN:HB3	2.21	0.40
1:A:875:CYS:HB3	1:A:926:CYS:HB2	1.97	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles (i)

### 5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	183/189 (97%)	178 (97%)	5 (3%)	0	100 100
1	B	176/189 (93%)	173 (98%)	3 (2%)	0	100 100
1	C	175/189 (93%)	172 (98%)	3 (2%)	0	100 100
All	All	534/567 (94%)	523 (98%)	11 (2%)	0	100 100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	174/176 (99%)	167 (96%)	7 (4%)	27 48
1	B	170/176 (97%)	165 (97%)	5 (3%)	37 61
1	C	169/176 (96%)	167 (99%)	2 (1%)	67 84
All	All	513/528 (97%)	499 (97%)	14 (3%)	40 64

All (14) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	812	GLU
1	A	929	ILE
1	A	931	THR
1	A	950	VAL
1	A	960	ILE

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	A	974	ILE
1	A	989	THR
1	B	826	GLU
1	B	950	VAL
1	B	955	LYS
1	B	960	ILE
1	B	989	THR
1	C	960	ILE
1	C	989	THR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	897	HIS
1	A	964	ASN
1	B	964	ASN
1	C	876	GLN
1	C	964	ASN

### 5.3.3 RNA [\(i\)](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [\(i\)](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [\(i\)](#)

There are no oligosaccharides in this entry.

### 5.6 Ligand geometry [\(i\)](#)

3 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond

length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
2	NAG	C	1001	1	14,14,15	0.24	0	17,19,21	0.82	1 (5%)
2	NAG	B	1001	1	14,14,15	0.21	0	17,19,21	0.77	1 (5%)
2	NAG	A	1001	1	14,14,15	0.24	0	17,19,21	0.63	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAG	C	1001	1	-	0/6/23/26	0/1/1/1
2	NAG	B	1001	1	-	2/6/23/26	0/1/1/1
2	NAG	A	1001	1	-	0/6/23/26	0/1/1/1

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed( $^{\circ}$ )	Ideal( $^{\circ}$ )
2	C	1001	NAG	C1-O5-C5	2.46	115.53	112.19
2	B	1001	NAG	C1-O5-C5	2.36	115.39	112.19

There are no chirality outliers.

All (2) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	B	1001	NAG	C8-C7-N2-C2
2	B	1001	NAG	O7-C7-N2-C2

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers [\(i\)](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [\(i\)](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [\(i\)](#)

### 6.1 Protein, DNA and RNA chains [\(i\)](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	185/189 (97%)	0.67	20 (10%) 12 12	30, 54, 85, 96	0
1	B	180/189 (95%)	0.58	15 (8%) 19 18	33, 48, 75, 91	0
1	C	179/189 (94%)	0.54	11 (6%) 28 26	29, 56, 83, 96	0
All	All	544/567 (95%)	0.60	46 (8%) 18 17	29, 53, 80, 96	0

All (46) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	972	SER	5.6
1	B	816	PHE	4.3
1	B	972	SER	4.2
1	A	974	ILE	4.2
1	B	810	VAL	4.1
1	B	935	VAL	3.7
1	A	812	GLU	3.7
1	C	968	GLN	3.6
1	C	816	PHE	3.3
1	A	933	GLU	3.2
1	B	991	LYS	3.2
1	A	968	GLN	3.1
1	A	971	GLU	3.1
1	A	989	THR	2.9
1	B	968	GLN	2.9
1	A	931	THR	2.9
1	C	859	LEU	2.9
1	A	969	PRO	2.8
1	B	971	GLU	2.8
1	A	965	PHE	2.8
1	A	986	LYS	2.8
1	C	916	TYR	2.8
1	B	990	CYS	2.8

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	A	991	LYS	2.7
1	B	970	ILE	2.7
1	B	973	LYS	2.6
1	A	992	ASN	2.6
1	A	813	SER	2.6
1	A	970	ILE	2.6
1	C	992	ASN	2.6
1	A	987	THR	2.5
1	B	974	ILE	2.5
1	C	809	SER	2.4
1	C	983	ASP	2.3
1	B	934	LYS	2.3
1	A	815	LYS	2.2
1	C	971	GLU	2.2
1	A	823	GLN	2.2
1	A	973	LYS	2.2
1	B	926	CYS	2.1
1	B	989	THR	2.1
1	C	972	SER	2.1
1	B	876	GLN	2.1
1	C	961	ILE	2.1
1	A	816	PHE	2.1
1	C	960	ILE	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [\(i\)](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [\(i\)](#)

There are no oligosaccharides in this entry.

## 6.4 Ligands [\(i\)](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

*Continued on next page...*

*Continued from previous page...*

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
2	NAG	B	1001	14/15	0.80	0.13	50,51,52,53	0
2	NAG	A	1001	14/15	0.90	0.10	44,44,45,45	0
2	NAG	C	1001	14/15	0.91	0.10	40,41,43,43	0

## 6.5 Other polymers [\(i\)](#)

There are no such residues in this entry.