



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 30, 2026 – 04:45 pm BST

PDB ID : 9GTZ / pdb_00009gtz
Title : Xenopus tropicalis Interleukin Enhancer-Binding Factor 3 (ILF3) and Interleukin Enhancer-Binding Factor 2 (ILF2) heterodimer.
Authors : Talbot, A.J.; Mancini, E.J.
Deposited on : 2024-09-18
Resolution : 2.53 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Xtrriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.48.1


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.53 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	164625	1004 (2.54-2.54)
Clashscore	180529	1055 (2.54-2.54)
Ramachandran outliers	177936	1048 (2.54-2.54)
Sidechain outliers	177891	1048 (2.54-2.54)
RSRZ outliers	164620	1004 (2.54-2.54)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	332	 19% 82% 17%
2	B	335	 13% 75% 11% 1% 13%

2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 4786 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Interleukin enhancer-binding factor 2 homolog.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	332	2460	1557	423	469	11	0	6	0

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	41	SER	THR	conflict	UNP Q6P8G1
A	163	GLY	GLU	conflict	UNP Q6P8G1

- Molecule 2 is a protein called Interleukin enhancer-binding factor 3.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	B	291	2183	1374	389	407	13	0	6	0

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	20	SER	VAL	conflict	UNP Q6GL57


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	74	Total	O	0	0
			74	74		
3	B	69	Total	O	0	0
			69	69		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Interleukin enhancer-binding factor 2 homolog



- Molecule 2: Interleukin enhancer-binding factor 3



4 Data and refinement statistics i

Property	Value	Source
Space group	P 31 2 1	Depositor
Cell constants a, b, c, α , β , γ	80.97Å 80.97Å 230.36Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	76.82 – 2.53 76.79 – 2.53	Depositor EDS
% Data completeness (in resolution range)	100.0 (76.82-2.53) 99.8 (76.79-2.53)	Depositor EDS
R_{merge}	0.14	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.24 (at 2.55Å)	Xtrriage
Refinement program	PHENIX 1.21.2_5419:	Depositor
R, R_{free}	0.220 , (Not available) 0.288 , 0.352	Depositor DCC
R_{free} test set	1475 reflections (4.90%)	wwPDB-VP
Wilson B-factor (Å ²)	62.5	Xtrriage
Anisotropy	0.014	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 55.2	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	0.018 for -h,-k,l	Xtrriage
F_o, F_c correlation	0.92	EDS
Total number of atoms	4786	wwPDB-VP
Average B, all atoms (Å ²)	65.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.18% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.16	0/2508	0.38	0/3422
2	B	0.16	0/2218	0.39	0/3016
All	All	0.16	0/4726	0.38	0/6438

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2460	0	2387	38	0
2	B	2183	0	2136	31	0
3	A	74	0	0	0	0
3	B	69	0	0	0	0
All	All	4786	0	4523	64	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

All (64) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:17:VAL:HG22	2:B:324:TYR:HD1	1.10	1.17

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:298:MET:HE3	2:B:302:GLU:HB3	1.39	1.04
1:A:17:VAL:HG22	2:B:324:TYR:CD1	1.92	1.03
1:A:17:VAL:CG2	2:B:324:TYR:HD1	1.90	0.83
2:B:298:MET:HE1	2:B:302:GLU:OE2	1.85	0.76
2:B:298:MET:CE	2:B:302:GLU:HB3	2.18	0.73
2:B:204:THR:HG21	2:B:239:LEU:HG	1.75	0.68
2:B:298:MET:HE1	2:B:302:GLU:CD	2.20	0.67
1:A:251:GLY:HA2	1:A:254:LEU:HD13	1.76	0.65
1:A:58:ILE:HD11	1:A:75:VAL:HG21	1.78	0.65
2:B:260:ILE:O	2:B:264:VAL:HG23	1.98	0.63
1:A:254:LEU:HD11	1:A:284:CYS:HB2	1.82	0.61
1:A:57:VAL:O	1:A:61:LEU:HD12	2.01	0.60
1:A:175:LEU:HA	1:A:178:ILE:HD12	1.84	0.59
1:A:151:THR:HA	1:A:171:LEU:HD22	1.83	0.58
1:A:275:MET:HB3	1:A:279:GLN:HB2	1.83	0.58
1:A:227:MET:HE3	1:A:227:MET:HA	1.86	0.57
2:B:31:VAL:O	2:B:35:VAL:HG23	2.06	0.56
2:B:144:LEU:HG	2:B:199:LEU:HD21	1.89	0.55
2:B:204:THR:HG23	2:B:235[A]:CYS:HB3	1.89	0.55
1:A:212:PRO:HG3	1:A:279:GLN:HB3	1.89	0.55
1:A:190:HIS:HB3	1:A:193:VAL:HG13	1.89	0.55
1:A:322:VAL:HG13	2:B:355:GLN:HG2	1.90	0.53
1:A:17:VAL:HG11	2:B:332:SER:C	2.33	0.53
1:A:191:SER:O	1:A:195:VAL:HG23	2.08	0.53
1:A:285:TYR:HB2	1:A:327:SER:HB2	1.89	0.53
1:A:103[A]:PRO:HD3	1:A:152:THR:HG21	1.91	0.53
2:B:209:ARG:O	2:B:213:GLU:HG3	2.09	0.53
1:A:153:VAL:HB	1:A:155:PRO:HD2	1.91	0.52
2:B:7:PHE:HD2	2:B:302:GLU:HG2	1.74	0.52
2:B:21:VAL:HG21	2:B:267:ASP:HB2	1.93	0.51
2:B:7:PHE:CD2	2:B:302:GLU:HG2	2.47	0.50
2:B:169:GLU:HA	2:B:169:GLU:OE1	2.13	0.49
1:A:50:LEU:HD23	1:A:146:VAL:HG23	1.94	0.49
2:B:202:ARG:HH11	2:B:202:ARG:HG2	1.77	0.48
1:A:58:ILE:HA	1:A:61:LEU:HD12	1.94	0.48
1:A:156:ASN:HA	1:A:159:LYS:HD2	1.95	0.48
1:A:34:ASN:CG	1:A:195:VAL:HG13	2.39	0.47
2:B:298:MET:CE	2:B:302:GLU:CD	2.86	0.47
1:A:193:VAL:HG12	1:A:226:VAL:HG11	1.97	0.47
1:A:282:MET:HA	1:A:285:TYR:CE2	2.51	0.46
2:B:194:GLU:HB3	2:B:195:PRO:HD3	1.97	0.45

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:213:LEU:HB3	1:A:218:LEU:HD21	1.98	0.45
2:B:202:ARG:HG2	2:B:202:ARG:NH1	2.32	0.45
1:A:3:ILE:HD12	1:A:4:PRO:HD2	1.99	0.45
2:B:298:MET:HE3	2:B:302:GLU:CB	2.28	0.45
1:A:54:ILE:O	1:A:58:ILE:HG23	2.17	0.45
1:A:282:MET:HA	1:A:285:TYR:CZ	2.52	0.45
1:A:58:ILE:HA	1:A:61:LEU:CD1	2.48	0.44
1:A:135:THR:HA	1:A:170:VAL:HG21	2.00	0.44
2:B:42:LEU:HD13	2:B:124:VAL:HG21	1.99	0.44
1:A:193:VAL:O	1:A:197:ILE:HG13	2.17	0.43
2:B:298:MET:CE	2:B:302:GLU:OE2	2.62	0.43
1:A:275:MET:HE3	1:A:275:MET:HB2	1.94	0.43
1:A:196:LEU:O	1:A:200:LEU:HG	2.19	0.42
1:A:110:ALA:O	1:A:113:ILE:HG13	2.20	0.42
1:A:292:ARG:O	1:A:295:SER:HB2	2.20	0.42
2:B:355:GLN:HB2	2:B:358:GLU:HB3	2.01	0.42
2:B:234:LYS:H	2:B:234:LYS:HG2	1.70	0.42
1:A:286:THR:O	1:A:290:LEU:HG	2.19	0.41
2:B:187:ILE:HG23	2:B:201:ILE:HB	2.01	0.41
2:B:296:ARG:CZ	2:B:298:MET:SD	3.09	0.41
2:B:291:ILE:HD11	2:B:303:ALA:HB1	2.03	0.41
1:A:190:HIS:O	1:A:194[A]:LYS:HG3	2.21	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	330/332 (99%)	324 (98%)	6 (2%)	0	100	100
2	B	283/335 (84%)	279 (99%)	4 (1%)	0	100	100
All	All	613/667 (92%)	603 (98%)	10 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	257/288 (89%)	245 (95%)	12 (5%)	22	32
2	B	226/292 (77%)	215 (95%)	11 (5%)	21	30
All	All	483/580 (83%)	460 (95%)	23 (5%)	21	31

All (23) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	35	GLN
1	A	70	VAL
1	A	111	LEU
1	A	128	LEU
1	A	152	THR
1	A	172	GLN
1	A	193	VAL
1	A	196	LEU
1	A	231	THR
1	A	277	LEU
1	A	291[A]	VAL
1	A	323	ILE
2	B	8	LEU
2	B	122	ARG
2	B	130	VAL
2	B	154	ILE
2	B	157	LEU
2	B	179	ILE
2	B	187	ILE
2	B	253	ASN
2	B	260	ILE
2	B	315	ILE
2	B	355	GLN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such

sidechains are listed below:

Mol	Chain	Res	Type
1	A	223	HIS
1	A	280	GLN
2	B	355	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

The following chains have linkage breaks:

Mol	Chain	Number of breaks
2	B	1

All chain breaks are listed below:

Model	Chain	Residue-1	Atom-1	Residue-2	Atom-2	Distance (Å)
1	B	340:ARG	C	355:GLN	N	4.77

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	332/332 (100%)	1.26	63 (18%) 4 5	39, 64, 101, 120	4 (1%)
2	B	291/335 (86%)	1.14	42 (14%) 7 9	40, 61, 91, 118	5 (1%)
All	All	623/667 (93%)	1.21	105 (16%) 5 6	39, 62, 99, 120	9 (1%)

All (105) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	152	THR	6.0
2	B	298	MET	5.8
1	A	20	ALA	5.1
2	B	36	SER	4.9
1	A	60	ASN	4.8
1	A	57	VAL	4.7
2	B	187	ILE	4.6
2	B	51	GLN	4.6
1	A	62	ILE	4.5
1	A	72	ILE	4.3
1	A	17	VAL	4.3
1	A	323	ILE	4.2
1	A	304	GLY	3.9
1	A	70	VAL	3.8
2	B	194	GLU	3.8
2	B	210	GLU	3.8
2	B	286	LEU	3.7
1	A	150	ILE	3.7
1	A	101	ILE	3.6
1	A	61	LEU	3.6
1	A	71	GLN	3.5
1	A	105	LEU	3.5
1	A	183	TRP	3.4
1	A	99	LEU	3.4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	166	LEU	3.4
1	A	134	GLU	3.4
2	B	355	GLN	3.2
1	A	111	LEU	3.2
2	B	240	ALA	3.2
1	A	154	PRO	3.1
1	A	96	VAL	3.1
1	A	264	CYS	3.0
2	B	256	LYS	3.0
2	B	50	ASP	3.0
2	B	206	PRO	2.9
2	B	212	MET	2.9
2	B	149	ARG	2.9
1	A	63	VAL	2.9
2	B	6	ILE	2.9
1	A	333	LYS	2.9
2	B	164	LEU	2.8
2	B	179	ILE	2.8
1	A	97	VAL	2.8
2	B	259	VAL	2.8
1	A	107	ALA	2.8
1	A	142	ALA	2.8
2	B	140	LEU	2.8
1	A	275	MET	2.8
1	A	127	VAL	2.8
2	B	192	THR	2.7
1	A	184	PHE	2.7
1	A	13	ALA	2.7
2	B	294	ALA	2.7
1	A	55	ASN	2.7
2	B	214	LYS	2.7
2	B	291	ILE	2.7
1	A	272	HIS	2.6
1	A	58	ILE	2.6
1	A	98	ILE	2.6
1	A	7	PHE	2.5
1	A	160	LEU	2.5
2	B	386	PRO	2.5
1	A	168	ILE	2.5
2	B	120	THR	2.5
2	B	331	ALA	2.4
1	A	139	ILE	2.4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	151	THR	2.4
2	B	195	PRO	2.4
2	B	119	ARG	2.4
2	B	49	ILE	2.3
1	A	64	ALA	2.3
2	B	168	PHE	2.3
1	A	258	VAL	2.3
2	B	340	ARG	2.3
2	B	176	TYR	2.3
1	A	93	ALA	2.3
1	A	153	VAL	2.2
1	A	277	LEU	2.2
1	A	266	SER	2.2
2	B	141	ASP	2.2
1	A	77	GLN	2.2
1	A	216	TRP	2.2
2	B	28	LEU	2.2
1	A	234	PRO	2.2
1	A	74	GLU	2.2
1	A	78	VAL	2.2
2	B	207	LEU	2.1
1	A	173	SER	2.1
1	A	205	SER	2.1
1	A	155	PRO	2.1
1	A	51	VAL	2.1
2	B	208	VAL	2.1
2	B	136	LEU	2.1
1	A	19	PRO	2.1
2	B	258	CYS	2.1
2	B	260	ILE	2.1
1	A	157	LEU	2.1
2	B	385	LEU	2.1
2	B	7	PHE	2.1
2	B	186	SER	2.1
1	A	8	HIS	2.1
1	A	124	PRO	2.0
1	A	145	THR	2.0
1	A	115	VAL	2.0
1	A	170	VAL	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.