



# Full wwPDB X-ray Structure Validation Report i

Nov 24, 2025 – 12:16 pm GMT

PDB ID : 9HJM / pdb\_00009hhm  
Title : Crystal structure of phosphatidyl inositol 4-kinase II beta in complex with HH5129  
Authors : Klima, M.; Boura, E.  
Deposited on : 2024-11-22  
Resolution : 2.25 Å (reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>  
with specific help available everywhere you see the i symbol.

The types of validation reports are described at  
<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references](#) i) were used in the production of this report:

MolProbity : 4.5-2 with Phenix2.0  
Mogul : 1.8.4, CSD as541be (2020)  
Xtriage (Phenix) : 2.0  
EDS : 3.0  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)  
CCP4 : 9.0.010 (Gargrove)  
Density-Fitness : 1.0.12  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.46


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## X-RAY DIFFRACTION


The reported resolution of this entry is 2.25 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	164625	1763 (2.26-2.26)
Clashscore	180529	1919 (2.26-2.26)
Ramachandran outliers	177936	1884 (2.26-2.26)
Sidechain outliers	177891	1885 (2.26-2.26)
RSRZ outliers	164620	1763 (2.26-2.26)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.



## 2 Entry composition (i)

There are 3 unique types of molecules in this entry. The entry contains 7458 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Phosphatidylinositol 4-kinase type 2-beta,Endolysin.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	446	Total	C	N	O	S	0	1	0
			3612	2336	608	657	11			
1	B	451	Total	C	N	O	S	0	1	0
			3609	2326	608	664	11			

There are 16 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	87	GLY	-	expression tag	UNP Q8TCG2
A	88	ALA	-	expression tag	UNP Q8TCG2
A	89	MET	-	expression tag	UNP Q8TCG2
A	1011	GLY	ARG	conflict	UNP P00720
A	1053	THR	CYS	conflict	UNP P00720
A	1096	ALA	CYS	conflict	UNP P00720
A	1136	ARG	ILE	conflict	UNP P00720
A	331	VAL	ILE	conflict	UNP Q8TCG2
B	87	GLY	-	expression tag	UNP Q8TCG2
B	88	ALA	-	expression tag	UNP Q8TCG2
B	89	MET	-	expression tag	UNP Q8TCG2
B	1011	GLY	ARG	conflict	UNP P00720
B	1053	THR	CYS	conflict	UNP P00720
B	1096	ALA	CYS	conflict	UNP P00720
B	1136	ARG	ILE	conflict	UNP P00720
B	331	VAL	ILE	conflict	UNP Q8TCG2



- Molecule 2 is (1 {S},2 {S},4 {S},5 {R})-6-[[[(2 {R},3 {S},4 {R},5 {R})-5-(6-aminopurin-9-yl)-3,4-bis(oxidanyl)oxolan-2-yl]methoxy-bis(oxidanylidene)-\$1^{\wedge}\{6\}-phosphanyl]oxy-bis(oxidanylidene)-\$1^{\wedge}\{6\}-phosphanyl]methyl-bis(oxidanylidene)-\$1^{\wedge}\{6\}-phosphanyl]oxycyclohexane-1,2,3,4,5-pentol (CCD ID: A1IVA) (formula: C<sub>17</sub>H<sub>25</sub>N<sub>5</sub>O<sub>17</sub>P<sub>3</sub>) (labeled as "Ligand of Interest" by depositor).




### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Phosphatidylinositol 4-kinase type 2-beta, Endolysin



- Molecule 1: Phosphatidylinositol 4-kinase type 2-beta, Endolysin



## 4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 41 21 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	105.24Å 105.24Å 211.64Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	39.27 – 2.25 39.27 – 2.25	Depositor EDS
% Data completeness (in resolution range)	99.8 (39.27-2.25) 99.8 (39.27-2.25)	Depositor EDS
$R_{merge}$	0.28	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle^1$	1.38 (at 2.24Å)	Xtriage
Refinement program	PHENIX 1.20.1_4487	Depositor
$R$ , $R_{free}$	0.216 , 0.240 0.216 , 0.240	Depositor DCC
$R_{free}$ test set	2856 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	52.5	Xtriage
Anisotropy	0.092	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.33 , 27.1	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	7458	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	59.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.13% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [\(i\)](#)

### 5.1 Standard geometry [\(i\)](#)

Bond lengths and bond angles in the following residue types are not validated in this section: A1IVA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.10	0/3699	0.27	0/5009
1	B	0.10	0/3692	0.28	0/5003
All	All	0.10	0/7391	0.27	0/10012

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [\(i\)](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3612	0	3561	9	0
1	B	3609	0	3539	18	0
2	A	42	0	0	0	0
2	B	31	0	0	0	0
3	A	86	0	0	0	0
3	B	78	0	0	0	0
All	All	7458	0	7100	27	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (27) close contacts within the same asymmetric unit are listed below, sorted by their clash

magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:386:ILE:HB	1:B:425:ILE:HG23	1.92	0.51
1:A:100:PHE:HZ	1:A:210:VAL:HG12	1.75	0.50
1:B:187:LEU:HD11	1:B:406:ASP:HB2	1.94	0.49
1:B:1010:GLU:OE2	1:B:1144:ARG:NH1	2.43	0.49
1:A:263:TYR:HB3	1:A:311:LEU:HB3	1.94	0.49
1:B:1025:THR:HG22	1:B:1031:LEU:HA	1.95	0.49
1:A:306:GLY:HA2	1:A:363:HIS:HB2	1.97	0.46
1:B:218:ASN:O	1:B:252:LYS:NZ	2.48	0.46
1:A:1019:ASP:OD1	1:A:1023:TYR:N	2.37	0.45
1:B:149:PRO:HG2	1:B:152:GLU:HB2	1.99	0.45
1:A:187:LEU:HD11	1:A:406:ASP:HB2	2.00	0.44
1:B:306:GLY:HA2	1:B:363:HIS:HB2	1.99	0.44
1:B:1026:ILE:HG13	1:B:1027:GLY:N	2.32	0.44
1:B:1049:ILE:HD12	1:B:1061:GLU:HB3	1.99	0.43
1:B:404:LYS:HA	1:B:409:PHE:CG	2.53	0.43
1:B:1032:LEU:HD23	1:B:1032:LEU:HA	1.89	0.43
1:A:1077:ILE:HG12	1:A:1102:VAL:HG21	2.01	0.42
1:A:1010:GLU:OE2	1:A:1144:ARG:NH1	2.44	0.42
1:B:1051:ARG:NH1	1:B:1053:THR:HG21	2.34	0.42
1:B:125:ARG:HG2	1:B:127:SER:O	2.19	0.42
1:B:1033:THR:HG22	1:B:1041:ALA:HB2	2.00	0.42
1:B:1041:ALA:O	1:B:1045:LEU:N	2.48	0.41
1:B:386:ILE:HD11	1:B:432:LEU:HD12	2.02	0.41
1:B:1020:THR:HG21	1:B:1141:THR:HG22	2.03	0.41
1:B:378:ILE:HD12	1:B:378:ILE:HA	1.95	0.41
1:A:100:PHE:CZ	1:A:210:VAL:HG12	2.54	0.40
1:A:1086:VAL:HG21	1:A:1117:LEU:HB3	2.03	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles

### 5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	439/517 (85%)	435 (99%)	4 (1%)	0	100	100
1	B	444/517 (86%)	429 (97%)	15 (3%)	0	100	100
All	All	883/1034 (85%)	864 (98%)	19 (2%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [\(i\)](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	380/449 (85%)	380 (100%)	0	100	100
1	B	378/449 (84%)	378 (100%)	0	100	100
All	All	758/898 (84%)	758 (100%)	0	100	100

There are no protein residues with a non-rotameric sidechain to report.

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	1131	ASN
1	B	1139	ASN
1	B	443	GLN

### 5.3.3 RNA [\(i\)](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [\(i\)](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [\(i\)](#)

There are no oligosaccharides in this entry.

## 5.6 Ligand geometry [\(i\)](#)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	A1IVA	B	1201	-	25,33,45	1.11	1 (4%)	25,52,71	0.75	1 (4%)
2	A1IVA	A	1201	-	39,45,45	1.02	2 (5%)	43,71,71	0.83	2 (4%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	A1IVA	B	1201	-	-	6/14/38/67	0/3/3/4
2	A1IVA	A	1201	-	-	2/17/67/67	0/4/4/4

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	1201	A1IVA	PA-O5'	4.90	1.68	1.59
2	B	1201	A1IVA	PA-O5'	4.58	1.68	1.59
2	A	1201	A1IVA	OI4-Cl4	-2.85	1.39	1.43

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	1201	A1IVA	PG-OI4-Cl4	-4.08	111.94	120.28
2	A	1201	A1IVA	C5-C6-N6	2.34	123.91	120.35
2	B	1201	A1IVA	C5-C6-N6	2.34	123.90	120.35

There are no chirality outliers.



All (8) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	B	1201	A1IVA	PG-C3B-PB-O1B
2	B	1201	A1IVA	PG-C3B-PB-O2B
2	B	1201	A1IVA	PG-C3B-PB-O3A
2	B	1201	A1IVA	C5'-O5'-PA-O1A
2	A	1201	A1IVA	CI4-OI4-PG-C3B
2	B	1201	A1IVA	C5'-O5'-PA-O2A
2	A	1201	A1IVA	PB-O3A-PA-O2A
2	B	1201	A1IVA	PB-O3A-PA-O2A

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



## 5.7 Other polymers (i)

There are no such residues in this entry.

## 5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.

## 6 Fit of model and data i

### 6.1 Protein, DNA and RNA chains i

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	446/517 (86%)	0.11	13 (2%) 54 54	23, 53, 89, 130	1 (0%)
1	B	451/517 (87%)	0.23	30 (6%) 25 24	22, 55, 121, 142	1 (0%)
All	All	897/1034 (86%)	0.17	43 (4%) 36 36	22, 54, 107, 142	2 (0%)

All (43) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	156	GLY	4.7
1	A	359	ALA	4.2
1	A	250	PRO	3.9
1	B	1024	TYR	3.7
1	B	1016	ILE	3.7
1	B	1055	GLY	3.6
1	B	357	TRP	3.4
1	B	155	TYR	3.3
1	B	1038	LEU	3.2
1	B	1032	LEU	3.1
1	B	1034	LYS	3.0
1	B	1020	THR	2.9
1	B	1056	VAL	2.9
1	B	337	LEU	2.8
1	A	213	VAL	2.8
1	B	1031	LEU	2.8
1	A	357	TRP	2.7
1	B	98	ASN	2.7
1	B	1042	LYS	2.6
1	B	128	GLN	2.6
1	B	1054	ASN	2.6
1	B	1057	ILE	2.6
1	A	1162	ASN	2.6
1	A	1041	ALA	2.5

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	B	130	SER	2.5
1	B	1014	LEU	2.5
1	B	1045	LEU	2.4
1	B	1059	LYS	2.3
1	B	1046	ASP	2.3
1	B	1162	ASN	2.3
1	A	132	GLY	2.3
1	B	1036	PRO	2.2
1	A	336	PHE	2.2
1	B	1039	ASN	2.2
1	A	219	TYR	2.2
1	B	132	GLY	2.2
1	B	1103	PHE	2.2
1	A	122	PHE	2.1
1	A	1039	ASN	2.1
1	B	1041	ALA	2.1
1	B	119	VAL	2.1
1	A	131	SER	2.1
1	A	1026	ILE	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [\(i\)](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [\(i\)](#)

There are no oligosaccharides in this entry.


## 6.4 Ligands [\(i\)](#)


In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
2	A1IVA	B	1201	31/42	0.77	0.15	51,103,126,137	0
2	A1IVA	A	1201	42/42	0.89	0.10	55,76,140,144	0

The following is a graphical depiction of the model fit to experimental electron density of all

instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.





## 6.5 Other polymers [\(i\)](#)

There are no such residues in this entry.