



Full wwPDB X-ray Structure Validation Report ⓘ

Dec 1, 2025 – 12:51 pm GMT

PDB ID : 9HIM / pdb_00009him
Title : ManDH5 E303Q in complex with mannatriose after co-crystallization with mannohexaose at 1.36 angstroms resolution a beta-D-Mannanase of GH5 family from Dictyoglomus thermophilum
Authors : Sivron, Y.; Romano, A.; Shoham, Y.; Shoham, G.
Deposited on : 2024-11-26
Resolution : 1.36 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	1.8.4, CSD as541be (2020)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20231227.v01 (using entries in the PDB archive December 27th 2023)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.46


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.36 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	164625	1089 (1.36-1.36)
Clashscore	180529	1157 (1.36-1.36)
Ramachandran outliers	177936	1146 (1.36-1.36)
Sidechain outliers	177891	1146 (1.36-1.36)
RSRZ outliers	164620	1088 (1.36-1.36)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	568	5% 93% 6%
2	C	3	33% 67%

2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 9884 atoms, of which 4607 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called DUF5060 domain-containing protein.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	563	Total	C	H	N	O	S	0	4	0
			9321	3116	4575	767	854	9			

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	303	GLN	GLU	engineered mutation	UNP A0A7C3MIF0

- Molecule 2 is an oligosaccharide called beta-D-mannopyranose-(1-4)-beta-D-mannopyranose-(1-4)-beta-D-mannopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
2	C	3	Total	C	H	O	0	0	0
			66	18	32	16			


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	496	Total	O	0	1
			497	497		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: DUF5060 domain-containing protein



- Molecule 2: beta-D-mannopyranose-(1-4)-beta-D-mannopyranose-(1-4)-beta-D-mannopyranose



4 Data and refinement statistics

Property	Value	Source
Space group	I 2 2 2	Depositor
Cell constants a, b, c, α , β , γ	95.48Å 99.35Å 153.62Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	45.10 – 1.36 45.10 – 1.36	Depositor EDS
% Data completeness (in resolution range)	100.0 (45.10-1.36) 100.0 (45.10-1.36)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.61 (at 1.36Å)	Xtriage
Refinement program	PHENIX 1.17.1_3660	Depositor
R, R_{free}	0.147 , 0.168 0.148 , 0.169	Depositor DCC
R_{free} test set	7764 reflections (4.84%)	wwPDB-VP
Wilson B-factor (Å ²)	14.2	Xtriage
Anisotropy	0.215	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.39 , 35.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	0.010 for k,h,-l	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	9884	wwPDB-VP
Average B, all atoms (Å ²)	22.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.86% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: BMA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.39	0/4910	0.63	0/6662

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4746	4575	4581	24	0
2	C	34	32	28	3	0
3	A	497	0	0	1	0
All	All	5277	4607	4609	24	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (24) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:392:GLU:OE1	2:C:1:BMA:H1	1.60	0.99

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:92:ILE:CD1	1:A:106:ILE:HD12	2.11	0.80
1:A:92:ILE:HD12	1:A:106:ILE:HD12	1.68	0.76
1:A:10:LEU:O	1:A:114:ARG:NH1	2.25	0.69
1:A:488:ILE:HD11	1:A:521:ILE:HG21	1.78	0.65
1:A:537:LYS:NZ	1:A:540:GLU:OE1	2.24	0.63
1:A:418:THR:HG21	1:A:427:ALA:HB2	1.88	0.54
1:A:273:GLU:HG2	1:A:277[A]:TYR:CE2	2.42	0.54
1:A:303:GLN:NE2	2:C:1:BMA:O1	2.43	0.51
1:A:521:ILE:HD12	1:A:525:ILE:HG21	1.94	0.48
1:A:419:ILE:HG21	1:A:448:LEU:HD11	1.98	0.45
1:A:92:ILE:HD12	1:A:106:ILE:CD1	2.44	0.45
1:A:34:LEU:C	1:A:34:LEU:HD12	2.42	0.44
1:A:160:CYS:HA	1:A:187:VAL:HA	1.98	0.44
1:A:399:GLY:HA2	1:A:431:TRP:CZ2	2.52	0.44
1:A:204:LYS:HD3	1:A:277[A]:TYR:OH	2.17	0.44
1:A:354:ILE:HG22	1:A:356:GLY:H	1.81	0.44
1:A:462:PHE:HB2	3:A:635:HOH:O	2.17	0.43
1:A:525:ILE:HG22	1:A:563:LEU:HD23	1.99	0.42
1:A:488:ILE:HD11	1:A:521:ILE:CG2	2.47	0.42
1:A:193:TRP:CZ3	2:C:3:BMA:H5	2.55	0.41
1:A:488:ILE:HB	1:A:561:ILE:HB	2.03	0.41
1:A:526:LYS:NZ	1:A:562:GLU:OE1	2.50	0.41

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	565/568 (100%)	552 (98%)	13 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	501/508 (99%)	501 (100%)	0	100	100

There are no protein residues with a non-rotameric sidechain to report.

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	272	ASN
1	A	303	GLN
1	A	440	ASN
1	A	467	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

3 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	BMA	C	1	2	12,12,12	1.62	3 (25%)	17,17,17	2.40	9 (52%)
2	BMA	C	2	2	11,11,12	2.21	4 (36%)	15,15,17	2.72	7 (46%)
2	BMA	C	3	2	11,11,12	2.01	4 (36%)	15,15,17	1.59	4 (26%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	BMA	C	1	2	-	0/2/22/22	0/1/1/1
2	BMA	C	2	2	-	0/2/19/22	0/1/1/1
2	BMA	C	3	2	-	0/2/19/22	0/1/1/1

All (11) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	C	2	BMA	O2-C2	-4.52	1.33	1.43
2	C	3	BMA	C2-C3	3.89	1.58	1.52
2	C	2	BMA	C2-C3	3.73	1.58	1.52
2	C	1	BMA	O5-C1	-2.66	1.36	1.42
2	C	2	BMA	C1-C2	2.62	1.58	1.52
2	C	2	BMA	O5-C5	-2.51	1.38	1.43
2	C	3	BMA	O5-C1	-2.50	1.39	1.43
2	C	3	BMA	C1-C2	2.46	1.57	1.52
2	C	3	BMA	O2-C2	-2.43	1.38	1.43
2	C	1	BMA	O6-C6	-2.28	1.32	1.42
2	C	1	BMA	C3-C2	2.06	1.57	1.52

All (20) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	C	2	BMA	C1-C2-C3	-6.88	101.21	109.67
2	C	2	BMA	C1-O5-C5	4.34	118.08	112.19
2	C	1	BMA	C1-O5-C5	4.30	121.77	113.66
2	C	1	BMA	C6-C5-C4	4.29	123.05	113.00
2	C	1	BMA	O2-C2-C3	3.72	118.95	110.35
2	C	3	BMA	O5-C5-C6	3.44	112.60	107.20
2	C	1	BMA	O5-C1-C2	3.15	115.91	110.28
2	C	1	BMA	O4-C4-C5	-2.85	102.21	109.30
2	C	2	BMA	O5-C5-C4	-2.80	104.01	110.83

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	C	2	BMA	O2-C2-C3	2.79	115.72	110.14
2	C	1	BMA	O6-C6-C5	-2.73	101.93	111.29
2	C	1	BMA	C4-C3-C2	2.51	115.20	110.82
2	C	1	BMA	O3-C3-C4	-2.44	104.71	110.35
2	C	2	BMA	C2-C3-C4	2.41	115.06	110.89
2	C	2	BMA	O4-C4-C3	-2.23	105.19	110.35
2	C	3	BMA	O5-C1-C2	2.17	114.12	110.77
2	C	1	BMA	C1-C2-C3	2.10	114.68	110.31
2	C	2	BMA	C3-C4-C5	2.09	113.96	110.24
2	C	3	BMA	C3-C4-C5	-2.08	106.52	110.24
2	C	3	BMA	O2-C2-C1	-2.02	105.02	109.15

There are no chirality outliers.


There are no torsion outliers.

There are no ring outliers.

2 monomers are involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	C	1	BMA	2	0
2	C	3	BMA	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.



5.6 Ligand geometry

There are no ligands in this entry.

5.7 Other polymers

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ > 2		OWAB(Å ²)	Q < 0.9
1	A	563/568 (99%)	-0.04	31 (5%)	32 40	10, 19, 42, 59	4 (0%)

All (31) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	403	TRP	6.0
1	A	510	SER	5.4
1	A	509	ASP	4.9
1	A	520	LEU	4.7
1	A	508	MET	4.3
1	A	521	ILE	4.2
1	A	18	SER	4.0
1	A	454[A]	ASP	3.8
1	A	354	ILE	3.7
1	A	511	SER	3.7
1	A	544	ASP	3.1
1	A	466	THR	3.0
1	A	507	ASN	3.0
1	A	465	LEU	2.9
1	A	124	TYR	2.9
1	A	563	LEU	2.7
1	A	541	LEU	2.7
1	A	564	LEU	2.6
1	A	496	LYS	2.6
1	A	119	ASP	2.5
1	A	547	VAL	2.5
1	A	419	ILE	2.4
1	A	19	LYS	2.4
1	A	522	LYS	2.3
1	A	519	LEU	2.3
1	A	545	LYS	2.3
1	A	548	ILE	2.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	505	ILE	2.2
1	A	538	ILE	2.2
1	A	526	LYS	2.1
1	A	455	ILE	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)


There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	BMA	C	1	12/12	-	-	19,35,43,45	23
2	BMA	C	2	11/12	-	-	25,33,39,41	21
2	BMA	C	3	11/12	-	-	39,43,51,52	22

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.



6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.