



Full wwPDB X-ray Structure Validation Report i

Jan 5, 2026 – 03:49 pm GMT

PDB ID : 9HRG / pdb_00009hrg
Title : Crystal structure of the Class V (G61A) GTP aptamer variant in complex with GTP
Authors : Stafflinger, H.; Woehnert, J.; Schindelin, H.
Deposited on : 2024-12-18
Resolution : 2.05 Å (reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>
with specific help available everywhere you see the i symbol.

The types of validation reports are described at
<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references](#) i) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	?? (??), CSD ??CSD?? (????)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
buster-report	:	1.1.7 (2018)
Percentile statistics	:	20231227.v01 (using entries in the PDB archive December 27th 2023)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.47


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION


The reported resolution of this entry is 2.05 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	164625	2096 (2.04-2.04)
Clashscore	180529	2229 (2.04-2.04)
RSRZ outliers	164620	2096 (2.04-2.04)
RNA backbone	3690	1036 (2.40-1.68)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$ The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.



2 Entry composition (i)

There are 5 unique types of molecules in this entry. The entry contains 6356 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a RNA chain called Class V (G61A) GTP aptamer.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	68	Total	C	N	O	P	0	1	0
			1482	658	264	491	69			
1	B	68	Total	C	N	O	P	0	1	0
			1482	658	264	491	69			
1	C	68	Total	C	N	O	P	0	1	0
			1482	658	264	491	69			
1	D	68	Total	C	N	O	P	0	1	0
			1482	658	264	491	69			

- Molecule 2 is POTASSIUM ION (CCD ID: K) (formula: K).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	2	Total	K	0	0
			2	2		
2	B	2	Total	K	0	0
			2	2		
2	C	2	Total	K	0	0
			2	2		
2	D	2	Total	K	0	0
			2	2		

- Molecule 3 is GUANOSINE-5'-TRIPHOSPHATE (CCD ID: GTP) (formula: C₁₀H₁₆N₅O₁₄P₃) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total	C	N	O	P	0	0
			32	10	5	14	3		
3	B	1	Total	C	N	O	P	0	0
			32	10	5	14	3		
3	C	1	Total	C	N	O	P	0	0
			32	10	5	14	3		
3	D	1	Total	C	N	O	P	0	0
			32	10	5	14	3		

- Molecule 4 is MAGNESIUM ION (CCD ID: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	3	Total 3	Mg 3	0	0
4	B	3	Total 3	Mg 3	0	0
4	C	3	Total 3	Mg 3	0	0
4	D	3	Total 3	Mg 3	0	0

- Molecule 5 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	60	Total O 60 60	0	0

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	B	82	Total O 82 82	0	0
5	C	52	Total O 52 52	0	0
5	D	86	Total O 86 86	0	0


3 Residue-property plots [\(i\)](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.



- Molecule 1: Class V (G61A) GTP aptamer





- Molecule 1: Class V (G61A) GTP aptamer



- Molecule 1: Class V (G61A) GTP aptamer



- Molecule 1: Class V (G61A) GTP aptamer



4 Data and refinement statistics i

Property	Value	Source
Space group	P 1 21 1	Deposited
Cell constants a, b, c, α , β , γ	52.22Å 126.25Å 68.60Å 90.00° 90.50° 90.00°	Deposited
Resolution (Å)	34.60 – 2.05 34.60 – 2.05	Deposited EDS
% Data completeness (in resolution range)	97.8 (34.60-2.05) 97.9 (34.60-2.05)	Deposited EDS
R_{merge}	0.06	Deposited
R_{sym}	(Not available)	Deposited
$\langle I/\sigma(I) \rangle^1$	1.58 (at 2.05Å)	Xtriage
Refinement program	REFMAC 5.8.0430 (refmacat 0.4.88), PHENIX 1.20.1_4487	Deposited
R , R_{free}	0.207 , 0.233 0.219 , 0.237	Deposited DCC
R_{free} test set	2719 reflections (4.89%)	wwPDB-1
Wilson B-factor (Å ²)	58.6	Xtriage
Anisotropy	0.155	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 60.2	EDS
L-test for twinning ²	$\langle L \rangle = 0.51$, $\langle L^2 \rangle = 0.35$	Xtriage
Estimated twinning fraction	0.075 for h,-k,-l	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	6356	wwPDB-1
Average B, all atoms (Å ²)	69.0	wwPDB-1

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 24.89 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 3.5176e-03. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality i

5.1 Standard geometry i

Bond lengths and bond angles in the following residue types are not validated in this section: GTP, MG, K

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.29	0/1656	0.51	0/2583
1	B	0.34	0/1656	0.54	0/2583
1	C	0.30	0/1656	0.51	0/2583
1	D	0.33	0/1656	0.53	0/2583
All	All	0.32	0/6624	0.52	0/10332

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts i

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1482	0	743	9	0
1	B	1482	0	743	13	0
1	C	1482	0	743	8	0
1	D	1482	0	743	7	0
2	A	2	0	0	0	0
2	B	2	0	0	0	0
2	C	2	0	0	0	0
2	D	2	0	0	0	0
3	A	32	0	12	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	B	32	0	12	0	0
3	C	32	0	12	0	0
3	D	32	0	12	0	0
4	A	3	0	0	0	0
4	B	3	0	0	0	0
4	C	3	0	0	0	0
4	D	3	0	0	0	0
5	A	60	0	0	0	0
5	B	82	0	0	1	0
5	C	52	0	0	1	0
5	D	86	0	0	0	0
All	All	6356	0	3020	35	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (35) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:43:G:H4'	1:D:44:U:H5'	1.51	0.93
1:B:43:G:H4'	1:B:44:U:H5'	1.56	0.85
1:C:44:U:H4'	1:C:45[B]:A:H5"	1.67	0.77
1:C:1:G:OP2	5:C:201:HOH:O	2.05	0.73
1:A:30:U:O2'	1:A:32:G:O6	2.09	0.67
1:C:43:G:H4'	1:C:44:U:H5"	1.78	0.66
1:A:43:G:H4'	1:A:44:U:H5"	1.79	0.65
1:A:30:U:O2'	1:A:31:C:OP1	2.16	0.64
1:B:40:G:N3	1:B:40:G:H3'	2.14	0.63
1:C:32:G:O2'	1:C:33:G:O5'	2.18	0.61
1:D:32:G:O2'	1:D:33:G:O5'	2.18	0.59
1:B:60:A:N7	5:B:201:HOH:O	2.32	0.58
1:D:40:G:H3'	1:D:40:G:N3	2.21	0.56
1:C:32:G:O2'	1:C:33:G:O4'	2.21	0.55
1:C:40:G:N3	1:C:40:G:H3'	2.22	0.54
1:A:40:G:H3'	1:A:40:G:N3	2.24	0.53
1:A:32:G:O2'	1:A:33:G:O5'	2.26	0.52
1:C:22:U:H2'	1:C:23:C:C6	2.45	0.51
1:A:9:U:O2'	1:B:41:G:H1'	2.11	0.50
1:A:22:U:H2'	1:A:23:C:C6	2.47	0.49
1:B:43:G:H4'	1:B:44:U:C5'	2.35	0.48
1:A:45[A]:A:O4'	1:B:45[A]:A:O4'	2.31	0.48

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:32:G:O2'	1:B:33:G:O4'	2.24	0.47
1:B:31:C:H5'	1:B:32:G:C5	2.50	0.46
1:A:32:G:O2'	1:A:33:G:O4'	2.32	0.46
1:D:15:A:H2'	1:D:16:G:O4'	2.17	0.45
1:B:30:U:O2'	1:B:32:G:O6	2.28	0.45
1:B:15:A:H2'	1:B:16:G:O4'	2.17	0.45
1:C:30:U:O2'	1:C:32:G:O6	2.26	0.45
1:D:49:A:C2	1:D:52:G:C6	3.05	0.44
1:D:22:U:H2'	1:D:23:C:C6	2.53	0.44
1:B:30:U:H3'	1:B:31:C:O2	2.17	0.44
1:B:31:C:H5'	1:B:32:G:C6	2.55	0.42
1:D:31:C:H5'	1:D:32:G:C5	2.54	0.42
1:B:45[A]:A:OP1	1:B:45[A]:A:H3'	2.20	0.41

There are no symmetry-related clashes.

5.3 Torsion angles [\(i\)](#)

5.3.1 Protein backbone [\(i\)](#)

There are no protein molecules in this entry.

5.3.2 Protein sidechains [\(i\)](#)

There are no protein molecules in this entry.

5.3.3 RNA [\(i\)](#)

Mol	Chain	Analysed	Backbone Outliers	Pucker Outliers
1	A	66/68 (97%)	7 (10%)	0
1	B	66/68 (97%)	9 (13%)	0
1	C	66/68 (97%)	8 (12%)	0
1	D	66/68 (97%)	8 (12%)	0
All	All	264/272 (97%)	32 (12%)	0

All (32) RNA backbone outliers are listed below:

Mol	Chain	Res	Type
1	A	31	C

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	32	G
1	A	33	G
1	A	40	G
1	A	44	U
1	A	51	U
1	A	52	G
1	B	31	C
1	B	32	G
1	B	33	G
1	B	40	G
1	B	44	U
1	B	46	G
1	B	51	U
1	B	52	G
1	B	61	A
1	C	31	C
1	C	32	G
1	C	33	G
1	C	40	G
1	C	44	U
1	C	46	G
1	C	51	U
1	C	52	G
1	D	31	C
1	D	32	G
1	D	33	G
1	D	40	G
1	D	44	U
1	D	46	G
1	D	51	U
1	D	52	G

There are no RNA pucker outliers to report.

5.4 Non-standard residues in protein, DNA, RNA chains [\(i\)](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [\(i\)](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry (i)

Of 24 ligands modelled in this entry, 20 are monoatomic - leaving 4 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.


There are no chirality outliers.


There are no torsion outliers.


There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.







5.7 Other polymers [\(i\)](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [\(i\)](#)

There are no chain breaks in this entry.

6 Fit of model and data i

6.1 Protein, DNA and RNA chains i

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	68/68 (100%)	0.00	1 (1%) 71 74	39, 68, 97, 134	1 (1%)
1	B	68/68 (100%)	-0.13	1 (1%) 71 74	39, 60, 105, 125	1 (1%)
1	C	68/68 (100%)	-0.05	1 (1%) 71 74	39, 67, 109, 125	1 (1%)
1	D	68/68 (100%)	-0.13	1 (1%) 71 74	39, 61, 101, 130	1 (1%)
All	All	272/272 (100%)	-0.08	4 (1%) 71 74	39, 65, 110, 134	4 (1%)

All (4) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	45[A]	A	3.9
1	B	45[A]	A	3.5
1	A	45[A]	A	2.6
1	C	45[A]	A	2.3

6.2 Non-standard residues in protein, DNA, RNA chains i

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates i


There are no oligosaccharides in this entry.


6.4 Ligands i

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
4	MG	C	106	1/1	0.81	0.34	91,91,91,91	0
4	MG	B	106	1/1	0.82	0.15	82,82,82,82	0
4	MG	D	106	1/1	0.88	0.10	79,79,79,79	0
4	MG	A	104	1/1	0.90	0.07	61,61,61,61	0
4	MG	A	106	1/1	0.90	0.19	80,80,80,80	0
3	GTP	A	103	32/32	0.90	0.09	45,56,103,109	0
3	GTP	C	103	32/32	0.90	0.10	45,54,114,117	0
3	GTP	D	103	32/32	0.90	0.09	40,55,97,100	0
3	GTP	B	103	32/32	0.92	0.08	43,53,97,100	0
2	K	C	102	1/1	0.95	0.09	49,49,49,49	0
2	K	A	102	1/1	0.95	0.08	50,50,50,50	0
4	MG	B	105	1/1	0.96	0.04	55,55,55,55	0
2	K	A	101	1/1	0.97	0.07	48,48,48,48	0
2	K	C	101	1/1	0.98	0.06	48,48,48,48	0
2	K	B	101	1/1	0.98	0.04	47,47,47,47	0
2	K	D	102	1/1	0.98	0.04	44,44,44,44	0
4	MG	B	104	1/1	0.98	0.03	56,56,56,56	0
2	K	B	102	1/1	0.99	0.06	45,45,45,45	0
4	MG	C	104	1/1	0.99	0.04	58,58,58,58	0
4	MG	C	105	1/1	0.99	0.03	59,59,59,59	0
2	K	D	101	1/1	0.99	0.10	45,45,45,45	0
4	MG	D	104	1/1	0.99	0.03	63,63,63,63	0
4	MG	D	105	1/1	0.99	0.03	53,53,53,53	0
4	MG	A	105	1/1	0.99	0.04	53,53,53,53	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.





6.5 Other polymers [\(i\)](#)

There are no such residues in this entry.