



Full wwPDB X-ray Structure Validation Report i

Jan 19, 2026 – 12:23 pm GMT

PDB ID : 9I25 / pdb_00009i25
Title : WxLIP from Enterococcus faecium locus A bound to long WxL
Authors : Williamson, M.P.; Hassan, M.U.
Deposited on : 2025-01-17
Resolution : 1.91 Å (reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>
with specific help available everywhere you see the i symbol.

The types of validation reports are described at
<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references](#) ①) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.47


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION


The reported resolution of this entry is 1.91 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	164625	1028 (1.92-1.92)
Clashscore	180529	1100 (1.92-1.92)
Ramachandran outliers	177936	1087 (1.92-1.92)
Sidechain outliers	177891	1087 (1.92-1.92)
RSRZ outliers	164620	1028 (1.92-1.92)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.



2 Entry composition (i)

There are 3 unique types of molecules in this entry. The entry contains 4238 atoms, of which 2064 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called EfmWxLIP1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	C	251	Total	C	H	N	O	43	0	0
			3950	1255	1975	331	389			

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
C	135	LYS	THR	conflict	UNP A0A6N3AEW7

- Molecule 2 is a protein called WxL domain surface protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace	
2	A	13	Total	C	H	N	O	S	4	0	0
			182	59	89	14	19	1			


- Molecule 3 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	C	100	Total O 100 100	0	0
3	A	6	Total O 6 6	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: EfmWxLIP1



- Molecule 2: WxL domain surface protein



4 Data and refinement statistics i

Property	Value	Source
Space group	P 61	Depositor
Cell constants a, b, c, α , β , γ	105.15Å 105.15Å 67.17Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	54.00 – 1.91 54.00 – 1.91	Depositor EDS
% Data completeness (in resolution range)	99.9 (54.00-1.91) 100.0 (54.00-1.91)	Depositor EDS
R_{merge}	0.25	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle^1$	1.00 (at 1.91Å)	Xtriage
Refinement program	REFMAC 5.8.0405	Depositor
R , R_{free}	0.217, 0.261 0.215, 0.258	Depositor DCC
R_{free} test set	1690 reflections (5.13%)	wwPDB-VP
Wilson B-factor (Å ²)	34.5	Xtriage
Anisotropy	0.484	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.40, 32.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	0.033 for h,-h-k,-l	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	4238	wwPDB-VP
Average B, all atoms (Å ²)	49.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.99% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality i

5.1 Standard geometry i

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	C	1.01	2/2011 (0.1%)	1.53	15/2730 (0.5%)
2	A	1.30	1/95 (1.1%)	1.72	2/130 (1.5%)
All	All	1.02	3/2106 (0.1%)	1.54	17/2860 (0.6%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	C	0	4
2	A	0	1
All	All	0	5

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	C	30	GLU	CD-OE1	-5.79	1.14	1.25
1	C	256	THR	C-O	-5.61	1.17	1.24
2	A	219	GLU	CD-OE2	5.44	1.35	1.25

All (17) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	63	THR	CA-CB-OG1	-10.40	94.00	109.60
1	C	4	ASP	CA-CB-CG	6.88	119.48	112.60
1	C	70	ASN	CA-CB-CG	-6.83	105.77	112.60
1	C	5	PHE	CB-CA-C	6.30	123.94	110.07
1	C	50	GLU	CA-CB-CG	-6.24	101.62	114.10
1	C	97	HIS	CB-CA-C	-6.23	103.05	111.89
2	A	221	THR	CA-CB-OG1	-5.83	100.85	109.60
1	C	122	HIS	CA-CB-CG	5.80	119.60	113.80
1	C	233	LYS	CB-CA-C	-5.66	98.05	110.67

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	219	GLU	CB-CG-CD	5.52	121.99	112.60
1	C	207	TYR	N-CA-CB	-5.50	102.66	110.81
1	C	56	ALA	CA-C-O	-5.50	114.32	120.54
1	C	226	ASN	CA-CB-CG	-5.42	107.18	112.60
1	C	83	ASP	CA-CB-CG	5.35	117.95	112.60
1	C	72	THR	CA-CB-OG1	-5.20	101.80	109.60
1	C	207	TYR	CB-CA-C	5.11	118.25	109.15
1	C	37	VAL	N-CA-CB	-5.06	103.14	111.44

There are no chirality outliers.

All (5) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
2	A	214	GLY	Peptide
1	C	116	ARG	Sidechain
1	C	143	ARG	Sidechain
1	C	240	ARG	Sidechain
1	C	248	GLY	Peptide

5.2 Too-close contacts [\(i\)](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	C	1975	1975	1970	31	0
2	A	93	89	88	1	0
3	A	6	0	0	1	0
3	C	100	0	0	6	0
All	All	2174	2064	2058	31	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

All (31) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:248:GLY:HA3	3:C:353:HOH:O	1.46	1.15

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:39:ILE:HD12	1:C:92:ILE:HD11	1.49	0.95
1:C:4:ASP:HB2	3:C:308:HOH:O	1.72	0.89
1:C:2:ALA:O	1:C:146:TYR:OH	2.01	0.79
1:C:127:GLU:OE1	1:C:143:ARG:NH1	2.19	0.74
1:C:27:VAL:O	3:C:301:HOH:O	2.09	0.70
1:C:191:ILE:HA	3:C:304:HOH:O	1.93	0.67
1:C:191:ILE:HB	1:C:217:VAL:HG13	1.77	0.65
1:C:63:THR:HG23	3:A:303:HOH:O	2.02	0.59
1:C:260:THR:HG21	1:C:262:LYS:HE3	1.86	0.57
1:C:53:VAL:HG23	1:C:87:ILE:HD12	1.86	0.57
1:C:122:HIS:ND1	1:C:147:ASN:ND2	2.51	0.57
1:C:116:ARG:NH1	1:C:151:VAL:HG11	2.23	0.54
1:C:39:ILE:HD12	1:C:92:ILE:CD1	2.32	0.53
1:C:139:GLN:O	1:C:139:GLN:NE2	2.42	0.49
1:C:53:VAL:HG11	1:C:85:LEU:HB2	1.95	0.49
1:C:5:PHE:CZ	1:C:146:TYR:CD1	3.01	0.48
1:C:7:ILE:HG21	1:C:123:VAL:HG21	1.96	0.47
1:C:252:GLU:OE2	1:C:254:ARG:NH2	2.48	0.46
1:C:5:PHE:CE1	1:C:146:TYR:CD1	3.05	0.45
1:C:2:ALA:N	3:C:308:HOH:O	2.51	0.44
1:C:146:TYR:HA	2:A:220:LEU:O	2.18	0.43
1:C:60:ASP:OD2	1:C:224:HIS:HD2	2.01	0.43
1:C:5:PHE:CE2	1:C:146:TYR:HE1	2.37	0.42
1:C:50:GLU:OE2	1:C:91:GLU:OE1	2.38	0.42
1:C:5:PHE:CZ	1:C:146:TYR:CE1	3.09	0.41
1:C:250:SER:CB	3:C:353:HOH:O	2.69	0.41
1:C:81:VAL:HG11	1:C:119:ALA:HB2	2.02	0.41
1:C:94:VAL:HG22	1:C:100:THR:OG1	2.20	0.41
1:C:5:PHE:CZ	1:C:125:PRO:HD3	2.57	0.40
1:C:181:LEU:O	1:C:224:HIS:HA	2.21	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	C	247/277 (89%)	240 (97%)	5 (2%)	2 (1%)	16 6
2	A	11/226 (5%)	11 (100%)	0	0	100 100
All	All	258/503 (51%)	251 (97%)	5 (2%)	2 (1%)	16 6

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	C	174	ASN
1	C	173	VAL

5.3.2 Protein sidechains [\(i\)](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	C	221/241 (92%)	211 (96%)	10 (4%)	23 9
2	A	10/196 (5%)	10 (100%)	0	100 100
All	All	231/437 (53%)	221 (96%)	10 (4%)	25 10

All (10) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	37	VAL
1	C	43	SER
1	C	86	LEU
1	C	176	LYS
1	C	200	ILE
1	C	205	GLN
1	C	208	ILE
1	C	217	VAL
1	C	250	SER
1	C	263	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (8) such sidechains are listed below:

Mol	Chain	Res	Type
1	C	70	ASN
1	C	147	ASN
1	C	153	GLN
1	C	156	GLN
1	C	210	ASN
1	C	213	ASN
1	C	224	HIS
1	C	257	GLN

5.3.3 RNA [\(i\)](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [\(i\)](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [\(i\)](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [\(i\)](#)

There are no ligands in this entry.

5.7 Other polymers [\(i\)](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [\(i\)](#)

There are no chain breaks in this entry.

6 Fit of model and data i

6.1 Protein, DNA and RNA chains i

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	C	251/277 (90%)	-0.08	10 (3%) 43 48	28, 44, 81, 122	0
2	A	13/226 (5%)	0.25	1 (7%) 21 26	33, 39, 73, 74	0
All	All	264/503 (52%)	-0.06	11 (4%) 41 47	28, 44, 81, 122	0

All (11) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	2	ALA	5.7
1	C	173	VAL	4.8
1	C	175	ALA	3.8
1	C	3	GLY	3.8
1	C	140	ILE	2.8
2	A	214	GLY	2.6
1	C	127	GLU	2.4
1	C	233	LYS	2.3
1	C	263	LYS	2.3
1	C	139	GLN	2.1
1	C	207	TYR	2.0

6.2 Non-standard residues in protein, DNA, RNA chains i

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates i

There are no oligosaccharides in this entry.

6.4 Ligands [\(i\)](#)

There are no ligands in this entry.

6.5 Other polymers [\(i\)](#)

There are no such residues in this entry.