



Full wwPDB X-ray Structure Validation Report ⓘ

Nov 17, 2025 – 03:06 pm GMT

PDB ID : 9IA4 / pdb_00009ia4
Title : Bc8.108 Fab
Authors : Mechaly, A.E.; Haouz, A.; Beretta, M.; Caillet-Saguy, C.; Mouquet, H.
Deposited on : 2025-02-07
Resolution : 2.36 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Xtrriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.46


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.36 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	164625	1460 (2.36-2.36)
Clashscore	180529	1571 (2.36-2.36)
Ramachandran outliers	177936	1559 (2.36-2.36)
Sidechain outliers	177891	1559 (2.36-2.36)
RSRZ outliers	164620	1460 (2.36-2.36)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	215	
2	B	228	

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 3367 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Heavy chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	215	1649	1034	282	328	5	0	0	0

- Molecule 2 is a protein called Light chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	B	219	1617	1024	267	320	6	0	1	0


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	63	Total	O	0	0
			63	63		
3	B	38	Total	O	0	0
			38	38		

3 Residue-property plots [i](#)


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Heavy chain

Chain A: 



- Molecule 2: Light chain

Chain B: 



4 Data and refinement statistics

Property	Value	Source
Space group	C 2 2 21	Depositor
Cell constants a, b, c, α , β , γ	94.38Å 146.16Å 84.72Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	43.29 – 2.36 43.29 – 2.36	Depositor EDS
% Data completeness (in resolution range)	100.0 (43.29-2.36) 99.9 (43.29-2.36)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.19	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.57 (at 2.37Å)	Xtrriage
Refinement program	PHENIX 1.21.2_5419	Depositor
R, R_{free}	0.218 , 0.280 0.218 , 0.280	Depositor DCC
R_{free} test set	1225 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å ²)	50.3	Xtrriage
Anisotropy	0.283	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 45.7	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	3367	wwPDB-VP
Average B, all atoms (Å ²)	65.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.30% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality i

5.1 Standard geometry i

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.34	0/1686	0.53	0/2292
2	B	0.34	0/1656	0.55	0/2256
All	All	0.34	0/3342	0.54	0/4548

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts i

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1649	0	1611	11	0
2	B	1617	0	1581	13	0
3	A	63	0	0	1	0
3	B	38	0	0	0	0
All	All	3367	0	3192	24	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (24) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:130:PRO:HB3	2:B:156:TYR:HB3	1.61	0.82

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:50:ASP:HB2	1:A:53:LYS:HD2	1.73	0.69
1:A:143:ARG:NH1	3:A:301:HOH:O	2.34	0.60
2:B:162:THR:HG23	2:B:210[A]:ASN:HB3	1.86	0.56
1:A:125:GLN:HG2	1:A:130:THR:O	2.07	0.54
2:B:87:ARG:O	2:B:122:VAL:HG21	2.08	0.54
2:B:14:PRO:HD3	2:B:123:SER:C	2.36	0.49
2:B:14:PRO:HD3	2:B:123:SER:O	2.12	0.49
2:B:162:THR:CG2	2:B:210[A]:ASN:HB3	2.43	0.49
1:A:202:LEU:HD13	1:A:206:VAL:HG23	1.95	0.48
1:A:12:SER:HA	1:A:106:GLU:HG3	1.96	0.48
1:A:198:THR:HG22	1:A:205:PRO:HG3	1.98	0.46
1:A:62:PHE:CD1	1:A:75:ILE:HG12	2.53	0.44
2:B:209:VAL:HB	2:B:218:VAL:HG23	2.00	0.43
1:A:33:LEU:HD22	1:A:71:PHE:CG	2.53	0.43
1:A:179:THR:HG22	1:A:181:THR:HG23	2.01	0.43
2:B:88:SER:HA	2:B:122:VAL:HG23	2.00	0.42
1:A:81:GLU:H	1:A:81:GLU:CD	2.28	0.42
2:B:13:LYS:N	2:B:13:LYS:HE2	2.34	0.42
2:B:68:LEU:HD11	2:B:81:MET:HG2	2.01	0.42
2:B:99:VAL:HA	2:B:111:PHE:O	2.19	0.42
2:B:211:HIS:HB3	2:B:216:THR:HB	2.01	0.42
1:A:208:LYS:HD3	1:A:208:LYS:HA	1.91	0.41

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	213/215 (99%)	205 (96%)	8 (4%)	0	100	100
2	B	212/228 (93%)	195 (92%)	15 (7%)	2 (1%)	14	14
All	All	425/443 (96%)	400 (94%)	23 (5%)	2 (0%)	25	28

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	B	157	PHE
2	B	25	SER

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	185/185 (100%)	182 (98%)	3 (2%)	58 71
2	B	180/192 (94%)	176 (98%)	4 (2%)	47 59
All	All	365/377 (97%)	358 (98%)	7 (2%)	58 65

All (7) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	69	THR
1	A	155	LEU
1	A	203	SER
2	B	13	LYS
2	B	52	ILE
2	B	210[A]	ASN
2	B	210[B]	ASN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	215/215 (100%)	0.14	4 (1%) 66 71	36, 50, 84, 140	0
2	B	219/228 (96%)	1.27	49 (22%) 3 2	39, 71, 136, 166	1 (0%)
All	All	434/443 (97%)	0.71	53 (12%) 10 12	36, 59, 119, 166	1 (0%)

All (53) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	B	185	GLY	8.0
2	B	157	PHE	6.8
2	B	156	TYR	6.4
2	B	161	VAL	6.3
2	B	108	TRP	5.9
2	B	104	ASP	5.5
2	B	158	PRO	5.2
2	B	184	SER	4.9
2	B	123	SER	4.8
2	B	181	LEU	4.7
2	B	124	SER	4.3
2	B	145	GLY	3.9
2	B	160	PRO	3.8
2	B	188	SER	3.7
2	B	129	GLY	3.7
2	B	128	LYS	3.7
2	B	130	PRO	3.6
2	B	218	VAL	3.5
2	B	212	LYS	3.5
2	B	182	GLN	3.4
2	B	159	GLU	3.4
2	B	127	THR	3.3
2	B	209	VAL	3.3
2	B	216	THR	3.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
2	B	26	GLY	3.2
2	B	183	SER	3.1
2	B	210[A]	ASN	3.1
2	B	1	GLN	3.1
2	B	162	THR	2.9
2	B	226	SER	2.8
2	B	180	VAL	2.8
2	B	201	GLY	2.8
2	B	169	ALA	2.7
2	B	211	HIS	2.6
2	B	213	PRO	2.6
2	B	125	ALA	2.5
2	B	122	VAL	2.5
2	B	205	TYR	2.4
2	B	170	LEU	2.4
1	A	181	THR	2.4
2	B	190	SER	2.3
2	B	215	ASN	2.3
2	B	63	LYS	2.2
2	B	126	SER	2.1
2	B	206	ILE	2.1
1	A	213	GLY	2.1
2	B	131	SER	2.1
2	B	197	SER	2.1
2	B	27	GLY	2.1
1	A	215	CYS	2.1
2	B	167	SER	2.0
2	B	155	ASP	2.0
1	A	182	LEU	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.