



Full wwPDB X-ray Structure Validation Report i

Jan 14, 2026 – 08:45 pm GMT

PDB ID : 9IDU / pdb_00009idu
Title : Human Deoxyhypusine Synthase Fragment Screening Campaign - ligand VT00215
Authors : Wilk, P.; Wator-Wilk, E.; Krojer, T.; Grudnik, P.
Deposited on : 2025-02-15
Resolution : 1.34 Å (reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>
with specific help available everywhere you see the i symbol.

The types of validation reports are described at
<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references](#) i) were used in the production of this report:

MolProbity : 4.5-2 with Phenix2.0
Mogul : 1.8.4, CSD as541be (2020)
Xtriage (Phenix) : 2.0
EDS : 3.0
buster-report : 1.1.7 (2018)
Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.47


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION


The reported resolution of this entry is 1.34 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	164625	1904 (1.36-1.32)
Clashscore	180529	2038 (1.36-1.32)
Ramachandran outliers	177936	2016 (1.36-1.32)
Sidechain outliers	177891	2016 (1.36-1.32)
RSRZ outliers	164620	1903 (1.36-1.32)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.



The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	A1I2T	A	403	-	-	X	-
3	A1I2T	B	403	-	-	-	X

2 Entry composition [i](#)


There are 5 unique types of molecules in this entry. The entry contains 6183 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Deoxyhypusine synthase.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	350	Total	C	N	O	S	0	9	0
			2779	1772	470	520	17			
1	B	352	Total	C	N	O	S	0	6	0
			2781	1770	471	523	17			

- Molecule 2 is NICOTINAMIDE-ADENINE-DINUCLEOTIDE (CCD ID: NAD) (formula: $C_{21}H_{27}N_7O_{14}P_2$).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2		A		Total	C	N	O	P	
		1		44	21	7	14	2	0
2		B		Total	C	N	O	P	
		1		44	21	7	14	2	0

- Molecule 3 is methyl 3,5-dimethyl-1 {H}-pyrazole-4-carboxylate (CCD ID: A1I2T) (formula: $C_7H_{10}N_2O_2$) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total C N O 11 7 2 2	0	0
3	A	1	Total C N O 11 7 2 2	0	0
3	B	1	Total C N O 11 7 2 2	0	0

- Molecule 4 is (4R)-2-METHYLPENTANE-2,4-DIOL (CCD ID: MRD) (formula: C₆H₁₄O₂).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	B	1	Total C O 8 6 2	0	0


- Molecule 5 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	249	Total O 249 249	0	0
5	B	245	Total O 245 245	0	0


3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Deoxyhypusine synthase



- Molecule 1: Deoxyhypusine synthase



ASP

4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 32 2 1	Depositor
Cell constants a, b, c, α , β , γ	105.00 Å 105.00 Å 160.34 Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	46.08 – 1.34 46.08 – 1.34	Depositor EDS
% Data completeness (in resolution range)	97.7 (46.08-1.34) 97.7 (46.08-1.34)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle^1$	0.99 (at 1.34 Å)	Xtriage
Refinement program	REFMAC 5.8.0419	Depositor
R , R_{free}	0.199, 0.213 0.208, 0.221	Depositor DCC
R_{free} test set	11108 reflections (4.86%)	wwPDB-VP
Wilson B-factor (Å ²)	24.8	Xtriage
Anisotropy	0.127	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37, 41.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.51$, $\langle L^2 \rangle = 0.34$	Xtriage
Estimated twinning fraction	0.012 for -h,-k,l	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	6183	wwPDB-VP
Average B, all atoms (Å ²)	31.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.08% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality i

5.1 Standard geometry i

Bond lengths and bond angles in the following residue types are not validated in this section: NAD, A1I2T, CSS, MRD

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z > 5	RMSZ	# Z > 5
1	A	1.21	5/2846 (0.2%)	1.39	15/3859 (0.4%)
1	B	1.15	5/2845 (0.2%)	1.31	13/3857 (0.3%)
All	All	1.18	10/5691 (0.2%)	1.35	28/7716 (0.4%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	4

All (10) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	275[A]	CYS	N-CA	11.46	1.60	1.46
1	A	275[B]	CYS	N-CA	11.46	1.60	1.46
1	B	65	GLN	CD-OE1	7.04	1.36	1.23
1	B	271	ILE	C-O	-5.58	1.17	1.24
1	A	60	GLY	C-O	5.54	1.30	1.23
1	B	25	PRO	CA-C	5.22	1.54	1.51
1	B	286[A]	VAL	C-O	5.03	1.30	1.24
1	B	286[B]	VAL	C-O	5.03	1.30	1.24
1	A	261[A]	GLU	C-O	5.02	1.29	1.24
1	A	261[B]	GLU	C-O	5.02	1.29	1.24

All (28) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	241[A]	LEU	CA-C-O	10.18	131.34	120.55

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	241[B]	LEU	CA-C-O	10.18	131.34	120.55
1	A	261[A]	GLU	CA-C-O	9.76	130.90	120.55
1	A	261[B]	GLU	CA-C-O	9.76	130.90	120.55
1	B	65	GLN	CG-CD-NE2	-9.44	102.23	116.40
1	A	261[A]	GLU	N-CA-C	8.72	120.79	111.28
1	A	261[B]	GLU	N-CA-C	8.72	120.79	111.28
1	B	268[A]	THR	CA-C-O	8.24	129.41	120.10
1	B	268[B]	THR	CA-C-O	8.24	129.41	120.10
1	B	241[A]	LEU	N-CA-C	7.33	119.27	111.28
1	B	241[B]	LEU	N-CA-C	7.33	119.27	111.28
1	A	229	ILE	O-C-N	7.15	125.87	121.37
1	B	286[A]	VAL	CA-C-O	6.90	128.13	120.95
1	B	286[B]	VAL	CA-C-O	6.90	128.13	120.95
1	A	53	GLY	CA-C-O	-6.45	117.42	122.52
1	B	65	GLN	OE1-CD-NE2	6.35	128.95	122.60
1	A	274	LYS	CA-C-N	5.97	132.44	121.94
1	A	274	LYS	C-N-CA	5.97	132.44	121.94
1	A	24	LEU	CA-C-N	5.32	125.86	120.38
1	A	24	LEU	C-N-CA	5.32	125.86	120.38
1	B	305	TYR	CA-CB-CG	5.27	123.38	113.90
1	A	187	LEU	CA-C-O	-5.26	114.16	120.10
1	B	286[A]	VAL	N-CA-C	5.20	115.92	110.62
1	B	286[B]	VAL	N-CA-C	5.20	115.92	110.62
1	A	275[A]	CYS	CA-C-O	-5.09	114.28	120.54
1	A	275[B]	CYS	CA-C-O	-5.09	114.28	120.54
1	A	333	ASP	CA-CB-CG	5.08	117.68	112.60
1	A	305	TYR	CA-CB-CG	5.05	122.98	113.90

There are no chirality outliers.

All (4) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	275[A]	CYS	Mainchain
1	A	275[B]	CYS	Mainchain
1	A	85	ALA	Mainchain
1	A	86[A]	ASP	Mainchain

5.2 Too-close contacts [\(i\)](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen

atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2779	0	2769	26	0
1	B	2781	0	2769	22	0
2	A	44	0	26	0	0
2	B	44	0	26	1	0
3	A	22	0	0	12	0
3	B	11	0	0	1	0
4	B	8	0	14	0	0
5	A	249	0	0	4	1
5	B	245	0	0	7	1
All	All	6183	0	5604	48	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (48) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:329:LYS:HE3	5:B:696:HOH:O	1.22	1.33
1:A:24:LEU:HD13	3:A:403:A1I2T:C01	1.74	1.17
3:A:402:A1I2T:C01	5:A:565:HOH:O	2.27	0.82
1:A:24:LEU:CD2	3:A:403:A1I2T:C11	2.33	0.82
1:B:214:ILE:O	3:B:403:A1I2T:N08	2.13	0.81
1:B:27:GLU:O	5:B:502:HOH:O	2.06	0.72
1:A:137[B]:GLU:OE1	5:A:501:HOH:O	2.09	0.71
1:A:24:LEU:N	3:A:403:A1I2T:N08	2.17	0.69
1:A:24:LEU:HD21	3:A:403:A1I2T:C11	2.21	0.69
1:A:24:LEU:CD1	3:A:403:A1I2T:C01	2.64	0.68
1:B:14:LEU:C	1:B:14:LEU:HD13	2.20	0.67
1:A:47:GLU:OE1	5:A:502:HOH:O	2.12	0.66
1:B:75:GLU:OE1	5:B:503:HOH:O	2.14	0.65
1:A:285[B]:VAL:HG12	2:B:402:NAD:O3D	1.98	0.64
1:B:95:THR:HG23	5:B:601:HOH:O	2.00	0.62
1:A:359:MET:O	1:A:359:MET:HE3	2.03	0.57
1:A:24:LEU:C	3:A:403:A1I2T:C07	2.77	0.57
1:A:24:LEU:CD1	3:A:403:A1I2T:O02	2.53	0.56
1:B:21:SER:CB	1:B:29:THR:HG21	2.35	0.56
1:A:275[A]:CYS:SG	1:A:302:TYR:HE2	2.29	0.55
1:B:14:LEU:HD11	1:B:18:LEU:HD11	1.87	0.54
1:A:103:TYR:N	1:A:286[A]:VAL:HG11	2.22	0.54

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:264:ARG:O	1:B:268[A]:THR:HG23	2.08	0.53
1:B:88:THR:O	5:B:504:HOH:O	2.18	0.53
1:A:24:LEU:O	3:A:403:A1I2T:C07	2.57	0.52
1:B:21:SER:HB2	1:B:29:THR:HG21	1.92	0.52
1:A:83:GLN:HG3	1:A:84:HIS:N	2.25	0.51
1:A:295:LEU:HD12	1:A:295:LEU:O	2.11	0.50
1:A:28:SER:OG	3:A:403:A1I2T:C01	2.49	0.49
1:A:222:TYR:OH	1:A:226:LYS:NZ	2.45	0.48
1:A:147:TYR:CZ	3:A:402:A1I2T:C10	2.95	0.48
1:B:19:LYS:O	1:B:22:SER:OG	2.31	0.47
1:A:359:MET:HE3	1:A:359:MET:HA	1.98	0.45
1:A:226:LYS:NZ	5:A:507:HOH:O	2.44	0.45
1:B:95:THR:CG2	5:B:601:HOH:O	2.62	0.45
1:A:97:CYS:HA	1:A:275[A]:CYS:HB3	2.00	0.43
1:B:27:GLU:H	1:B:27:GLU:CD	2.25	0.43
1:A:164:ASN:HB3	1:B:327:TRP:CE2	2.54	0.43
1:A:359:MET:HE2	1:A:363:MET:SD	2.60	0.42
1:B:14:LEU:C	1:B:14:LEU:CD1	2.91	0.42
1:B:123:ASN:OD1	1:B:227:ASN:HB3	2.21	0.41
1:B:269:GLN:NE2	5:B:516:HOH:O	2.53	0.41
1:B:26:PRO:HD2	1:B:27:GLU:OE2	2.20	0.41
1:B:96:SER:HA	1:B:274:LYS:HD3	2.02	0.41
1:B:135:VAL:HA	1:B:221:TYR:CE2	2.56	0.41
1:A:24:LEU:HD13	3:A:403:A1I2T:O02	2.00	0.40
1:B:103:TYR:N	1:B:286[A]:VAL:HG11	2.37	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
5:A:714:HOH:O	5:B:717:HOH:O[4_555]	2.14	0.06

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	354/369 (96%)	349 (99%)	5 (1%)	0	100 100
1	B	355/369 (96%)	349 (98%)	6 (2%)	0	100 100
All	All	709/738 (96%)	698 (98%)	11 (2%)	0	100 100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [\(i\)](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	300/309 (97%)	294 (98%)	6 (2%)	50 17
1	B	301/309 (97%)	297 (99%)	4 (1%)	65 32
All	All	601/618 (97%)	591 (98%)	10 (2%)	60 23

All (10) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	23	THR
1	A	86[A]	ASP
1	A	86[B]	ASP
1	A	109	SER
1	A	115	THR
1	A	274	LYS
1	B	27	GLU
1	B	88	THR
1	B	95	THR
1	B	115	THR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	228	HIS

5.3.3 RNA [\(i\)](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [\(i\)](#)

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
1	CSS	B	177	1	4,6,7	0.79	0	1,6,8	0.62	0
1	CSS	A	177	1	4,6,7	0.84	0	1,6,8	0.57	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	CSS	B	177	1	-	0/1/5/7	-
1	CSS	A	177	1	-	0/1/5/7	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.5 Carbohydrates [\(i\)](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [\(i\)](#)

6 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	NAD	A	401	-	42,48,48	0.67	1 (2%)	50,73,73	0.99	3 (6%)
3	A1I2T	A	403	-	9,11,11	2.05	4 (44%)	8,15,15	2.01	4 (50%)
3	A1I2T	B	403	-	9,11,11	1.77	3 (33%)	8,15,15	1.83	3 (37%)
2	NAD	B	402	-	42,48,48	1.15	4 (9%)	50,73,73	1.11	2 (4%)
4	MRD	B	401	-	7,7,7	0.25	0	9,10,10	0.74	0
3	A1I2T	A	402	-	9,11,11	2.68	5 (55%)	8,15,15	2.12	2 (25%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAD	A	401	-	-	4/26/62/62	0/5/5/5
3	A1I2T	A	403	-	-	2/6/6/6	0/1/1/1
3	A1I2T	B	403	-	-	0/6/6/6	0/1/1/1
2	NAD	B	402	-	-	5/26/62/62	0/5/5/5
4	MRD	B	401	-	-	0/5/5/5	-
3	A1I2T	A	402	-	-	0/6/6/6	0/1/1/1

All (17) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	402	A1I2T	C06-N08	-5.97	1.26	1.34
2	B	402	NAD	C2N-N1N	4.26	1.40	1.35

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	403	A1I2T	C06-N08	-3.74	1.29	1.34
3	B	403	A1I2T	O02-C03	3.20	1.40	1.33
3	A	402	A1I2T	O04-C03	-2.83	1.15	1.22
3	A	403	A1I2T	O02-C01	-2.83	1.38	1.45
3	A	402	A1I2T	O02-C01	-2.57	1.39	1.45
3	A	402	A1I2T	C10-N09	-2.57	1.30	1.34
3	A	403	A1I2T	O04-C03	-2.36	1.16	1.22
2	B	402	NAD	C4N-C3N	2.36	1.43	1.39
2	B	402	NAD	C8A-N7A	-2.31	1.30	1.34
3	B	403	A1I2T	C06-N08	-2.28	1.31	1.34
2	B	402	NAD	PA-O2A	-2.23	1.44	1.55
3	A	403	A1I2T	O02-C03	2.19	1.38	1.33
3	A	402	A1I2T	N09-N08	-2.12	1.33	1.37
2	A	401	NAD	C8A-N7A	-2.08	1.31	1.34
3	B	403	A1I2T	O02-C01	-2.04	1.40	1.45

All (14) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	402	NAD	O2A-PA-O1A	4.26	133.30	112.24
2	A	401	NAD	O4D-C1D-C2D	-4.20	100.79	106.93
3	A	403	A1I2T	O02-C03-C05	3.84	121.47	112.27
3	A	402	A1I2T	C11-C10-N09	3.57	127.50	119.65
2	B	402	NAD	C6N-N1N-C2N	-3.13	119.12	121.97
3	B	403	A1I2T	O02-C03-C05	2.98	119.43	112.27
3	A	402	A1I2T	O02-C03-C05	2.90	119.23	112.27
3	B	403	A1I2T	C07-C06-N08	2.84	125.89	119.65
3	A	403	A1I2T	O02-C03-O04	-2.81	117.95	123.45
3	B	403	A1I2T	C11-C10-N09	2.65	125.47	119.65
2	A	401	NAD	C6N-N1N-C2N	-2.54	119.65	121.97
3	A	403	A1I2T	C11-C10-N09	2.21	124.51	119.65
2	A	401	NAD	C5A-C6A-N6A	2.20	123.69	120.35
3	A	403	A1I2T	C07-C06-N08	2.17	124.41	119.65

There are no chirality outliers.

All (11) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	401	NAD	C5D-O5D-PN-O1N
2	B	402	NAD	C5D-O5D-PN-O1N
3	A	403	A1I2T	O04-C03-O02-C01
3	A	403	A1I2T	C05-C03-O02-C01

Continued on next page...

Continued from previous page...


Mol	Chain	Res	Type	Atoms
2	A	401	NAD	PA-O3-PN-O5D
2	B	402	NAD	PA-O3-PN-O5D
2	A	401	NAD	C5D-O5D-PN-O3
2	B	402	NAD	C5D-O5D-PN-O3
2	A	401	NAD	C5D-O5D-PN-O2N
2	B	402	NAD	C5D-O5D-PN-O2N
2	B	402	NAD	PA-O3-PN-O2N


There are no ring outliers.


4 monomers are involved in 14 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	403	A1I2T	10	0
3	B	403	A1I2T	1	0
2	B	402	NAD	1	0
3	A	402	A1I2T	2	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.











5.7 Other polymers [\(i\)](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [\(i\)](#)

There are no chain breaks in this entry.

6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	349/369 (94%)	1.10	48 (13%) 8 10	11, 27, 70, 115	12 (3%)
1	B	351/369 (95%)	1.13	51 (14%) 7 10	13, 27, 46, 78	7 (1%)
All	All	700/738 (94%)	1.11	99 (14%) 7 10	11, 27, 52, 115	19 (2%)

All (99) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	12	GLY	12.4
1	A	18	LEU	11.0
1	A	14	LEU	10.2
1	B	24	LEU	9.8
1	B	13	ALA	9.8
1	B	17	VAL	9.2
1	B	23	THR	9.1
1	B	15	ALA	9.1
1	B	19	LYS	8.7
1	B	18	LEU	8.6
1	B	25	PRO	8.5
1	A	10	PRO	8.0
1	B	14	LEU	7.9
1	B	22	SER	7.6
1	B	20	HIS	7.3
1	A	17	VAL	7.2
1	B	16	ALA	7.1
1	A	11	ALA	7.0
1	A	24	LEU	6.9
1	B	21	SER	6.8
1	A	9	ALA	6.7
1	B	27	GLU	6.6
1	A	16	ALA	6.5
1	A	77	LEU	6.5

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	87	LEU	6.4
1	B	89	GLN	6.1
1	B	28	SER	6.0
1	A	19	LYS	6.0
1	A	26	PRO	5.7
1	A	91	ARG	5.7
1	A	13	ALA	5.4
1	A	15	ALA	5.4
1	A	20	HIS	5.4
1	B	26	PRO	5.3
1	A	147	TYR	5.1
1	A	25	PRO	4.9
1	A	363	MET	4.8
1	A	22	SER	4.8
1	A	12	GLY	4.7
1	B	85	ALA	4.7
1	B	250	TYR	4.6
1	A	274	LYS	4.3
1	A	28	SER	4.3
1	A	21	SER	4.3
1	A	362	PHE	4.2
1	A	361	ALA	4.2
1	B	94	LEU	4.1
1	A	84	HIS	3.9
1	A	85	ALA	3.6
1	A	305	TYR	3.6
1	B	29	THR	3.5
1	B	192	MET	3.5
1	A	275[A]	CYS	3.5
1	B	241[A]	LEU	3.3
1	A	23	THR	3.3
1	A	83	GLN	3.3
1	A	74	LEU	3.1
1	B	251	LYS	3.1
1	B	88	THR	3.0
1	B	363	MET	3.0
1	B	90	SER	3.0
1	B	87	LEU	2.9
1	B	174	GLU	2.9
1	B	305	TYR	2.8
1	A	195	ASN	2.8
1	A	360	ASP	2.8

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	196	THR	2.8
1	A	332	VAL	2.8
1	A	27	GLU	2.7
1	B	285	VAL	2.6
1	B	91	ARG	2.5
1	B	191	VAL	2.5
1	A	196	THR	2.5
1	B	77	LEU	2.5
1	B	153	LEU	2.5
1	A	261[A]	GLU	2.4
1	B	361	ALA	2.4
1	A	271	ILE	2.3
1	B	286[A]	VAL	2.3
1	B	43	ARG	2.2
1	B	345	LEU	2.2
1	A	76	PRO	2.2
1	B	93	PRO	2.2
1	A	66	VAL	2.2
1	A	93	PRO	2.2
1	A	186	ILE	2.2
1	A	200	LYS	2.2
1	A	272	PHE	2.2
1	B	75	GLU	2.1
1	B	298	ASN	2.1
1	B	74	LEU	2.1
1	A	29	THR	2.1
1	A	157	GLU	2.1
1	B	200	LYS	2.1
1	B	171	VAL	2.1
1	B	256	VAL	2.1
1	B	186	ILE	2.0
1	B	173	ASN	2.0
1	B	84	HIS	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [\(i\)](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
1	CSS	B	177	7/8	0.90	0.15	34,35,49,89	0
1	CSS	A	177	7/8	0.92	0.11	25,28,39,55	0

6.3 Carbohydrates [\(i\)](#)


There are no oligosaccharides in this entry.


6.4 Ligands [\(i\)](#)


In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	A1I2T	B	403	11/11	0.28	0.40	20,20,20,20	11
3	A1I2T	A	403	11/11	0.29	0.25	20,20,20,20	11
3	A1I2T	A	402	11/11	0.63	0.24	20,20,20,20	11
4	MRD	B	401	8/8	0.90	0.19	42,43,49,52	8
2	NAD	A	401	44/44	0.97	0.07	18,21,30,36	0
2	NAD	B	402	44/44	0.98	0.07	18,22,32,34	44


The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.











6.5 Other polymers [\(i\)](#)

There are no such residues in this entry.