



Full wwPDB X-ray Structure Validation Report ⓘ

Jan 14, 2026 – 08:29 pm GMT

PDB ID : 9IDY / pdb_00009idy
Title : Human Deoxyhypusine Synthase Fragment Screening Campaign - ligand VT00219
Authors : Wilk, P.; Water-Wilk, E.; Krojer, T.; Grudnik, P.
Deposited on : 2025-02-15
Resolution : 1.37 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	1.8.4, CSD as541be (2020)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
buster-report	:	1.1.7 (2018)
Percentile statistics	:	20231227.v01 (using entries in the PDB archive December 27th 2023)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.47


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.37 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	164625	3869 (1.40-1.36)
Clashscore	180529	4183 (1.40-1.36)
Ramachandran outliers	177936	4116 (1.40-1.36)
Sidechain outliers	177891	4115 (1.40-1.36)
RSRZ outliers	164620	3867 (1.40-1.36)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	369	
1	B	369	

2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 6065 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Deoxyhypusine synthase.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	353	Total	C	N	O	S	0	8	0
			2800	1782	477	525	16			
1	B	336	Total	C	N	O	S	0	7	0
			2675	1704	452	502	17			

- Molecule 2 is NICOTINAMIDE-ADENINE-DINUCLEOTIDE (CCD ID: NAD) (formula: $C_{21}H_{27}N_7O_{14}P_2$).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	N	O	P	0	0
			44	21	7	14	2		
2	B	1	Total	C	N	O	P	0	0
			44	21	7	14	2		

- Molecule 3 is 1-methyl-1-(phenylmethyl)urea (CCD ID: A1IGN) (formula: $C_9H_{12}N_2O$) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C	N	O	0	0
			12	9	2	1		

- Molecule 4 is (4R)-2-METHYLPENTANE-2,4-DIOL (CCD ID: MRD) (formula: $C_6H_{14}O_2$).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	B	1	Total	C	O	0	0
			8	6	2		

- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	241	Total 241	O 241	0	0
5	B	241	Total 241	O 241	0	0

- Molecule 1: Deoxyhypusine synthase



4 Data and refinement statistics

Property	Value	Source
Space group	P 32 2 1	Depositor
Cell constants a, b, c, α , β , γ	104.97Å 104.97Å 160.42Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	46.09 – 1.37 46.09 – 1.37	Depositor EDS
% Data completeness (in resolution range)	99.5 (46.09-1.37) 99.6 (46.09-1.37)	Depositor EDS
R_{merge}	0.06	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	0.98 (at 1.37Å)	Xtriage
Refinement program	REFMAC 5.8.0419	Depositor
R, R_{free}	0.191 , 0.206 0.201 , 0.215	Depositor DCC
R_{free} test set	10597 reflections (4.95%)	wwPDB-VP
Wilson B-factor (Å ²)	26.0	Xtriage
Anisotropy	0.109	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 41.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.51$, $\langle L^2 \rangle = 0.34$	Xtriage
Estimated twinning fraction	0.014 for -h,-k,l	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	6065	wwPDB-VP
Average B, all atoms (Å ²)	32.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.18% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: A1IGN, CSS, MRD, NAD

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.23	6/2867 (0.2%)	1.39	13/3886 (0.3%)
1	B	1.21	5/2739 (0.2%)	1.32	9/3711 (0.2%)
All	All	1.22	11/5606 (0.2%)	1.36	22/7597 (0.3%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	B	0	2

All (11) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	46[A]	LEU	N-CA	8.27	1.57	1.46
1	B	46[B]	LEU	N-CA	8.27	1.57	1.46
1	B	65	GLN	CD-OE1	7.62	1.38	1.23
1	B	271	ILE	C-O	-6.63	1.16	1.24
1	A	122	HIS	CE1-NE2	6.10	1.38	1.32
1	A	32[A]	ARG	C-O	6.02	1.31	1.23
1	A	32[B]	ARG	C-O	6.02	1.31	1.23
1	B	229	ILE	N-CA	5.83	1.50	1.46
1	A	64[A]	GLN	C-O	5.80	1.30	1.24
1	A	64[B]	GLN	C-O	5.80	1.30	1.24
1	A	152	SER	CA-CB	-5.61	1.43	1.53

All (22) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	241[A]	LEU	CA-C-O	10.66	131.85	120.55
1	B	241[B]	LEU	CA-C-O	10.66	131.85	120.55
1	B	65	GLN	CG-CD-NE2	-9.89	101.57	116.40
1	A	229	ILE	O-C-N	9.31	127.24	121.37
1	A	64[A]	GLN	CA-C-O	7.77	128.88	120.10
1	A	64[B]	GLN	CA-C-O	7.77	128.88	120.10
1	B	241[A]	LEU	N-CA-C	7.58	119.54	111.28
1	B	241[B]	LEU	N-CA-C	7.58	119.54	111.28
1	B	65	GLN	OE1-CD-NE2	6.86	129.46	122.60
1	A	24	LEU	CA-C-N	6.77	127.35	120.38
1	A	24	LEU	C-N-CA	6.77	127.35	120.38
1	A	32[A]	ARG	CA-C-O	6.34	128.80	120.11
1	A	32[B]	ARG	CA-C-O	6.34	128.80	120.11
1	A	229	ILE	CA-C-O	-6.17	114.56	119.25
1	B	29	THR	CA-CB-OG1	-6.16	100.36	109.60
1	B	47	GLU	CB-CG-CD	5.78	122.43	112.60
1	B	305	TYR	CA-CB-CG	5.54	123.87	113.90
1	A	133	GLY	O-C-N	5.52	127.49	122.19
1	A	64[A]	GLN	N-CA-C	5.24	117.73	111.71
1	A	64[B]	GLN	N-CA-C	5.24	117.73	111.71
1	A	305	TYR	CA-CB-CG	5.21	123.28	113.90
1	A	221	TYR	O-C-N	5.02	129.05	122.33

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	B	46[A]	LEU	Mainchain
1	B	46[B]	LEU	Mainchain

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2800	0	2789	22	0
1	B	2675	0	2663	5	0
2	A	44	0	26	0	0
2	B	44	0	26	1	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	A	12	0	0	3	0
4	B	8	0	14	0	0
5	A	241	0	0	3	1
5	B	241	0	0	2	1
All	All	6065	0	5518	27	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (27) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:64[A]:GLN:OE1	3:A:402:A1IGN:C6	1.74	1.34
1:A:64[A]:GLN:OE1	3:A:402:A1IGN:C7	1.89	1.19
1:A:64[A]:GLN:CD	3:A:402:A1IGN:C6	2.31	1.04
1:A:77:LEU:CD1	1:A:274:LYS:HD2	2.22	0.69
1:A:77:LEU:HD11	1:A:274:LYS:HD2	1.74	0.68
1:A:137[B]:GLU:OE1	5:A:501:HOH:O	2.14	0.65
1:A:83:GLN:HG3	1:A:84:HIS:N	2.17	0.60
1:A:285[B]:VAL:HG12	2:B:402:NAD:O3D	2.01	0.59
1:A:359:MET:O	1:A:359:MET:HE3	2.02	0.59
1:A:103:TYR:N	1:A:286:VAL:HG11	2.20	0.56
1:A:226:LYS:NZ	5:A:505:HOH:O	2.34	0.56
1:B:88:THR:O	5:B:503:HOH:O	2.18	0.55
1:A:77:LEU:CD1	1:A:274:LYS:CD	2.87	0.52
1:A:47:GLU:OE1	5:A:502:HOH:O	2.18	0.51
1:A:275:CYS:SG	1:A:302:TYR:HE2	2.36	0.48
1:A:359:MET:HE3	1:A:359:MET:HA	1.95	0.48
1:A:222:TYR:OH	1:A:226:LYS:NZ	2.46	0.47
1:B:269:GLN:NE2	5:B:511:HOH:O	2.47	0.46
1:A:295:LEU:HD12	1:A:295:LEU:O	2.15	0.45
1:A:10:PRO:O	1:A:13:ALA:HB3	2.19	0.42
1:A:137[B]:GLU:OE2	1:A:176:TYR:OH	2.27	0.42
1:B:122:HIS:CE1	1:B:359[B]:MET:HE1	2.54	0.42
1:A:83:GLN:CG	1:A:84:HIS:N	2.84	0.41
1:B:96:SER:HA	1:B:274:LYS:HD3	2.02	0.41
1:B:135:VAL:HA	1:B:221:TYR:CE2	2.56	0.41
1:A:32[A]:ARG:HB2	1:A:32[A]:ARG:CZ	2.51	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
5:A:706:HOH:O	5:B:710:HOH:O[4_555]	2.13	0.07

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	356/369 (96%)	352 (99%)	4 (1%)	0	100	100
1	B	340/369 (92%)	334 (98%)	6 (2%)	0	100	100
All	All	696/738 (94%)	686 (99%)	10 (1%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	302/309 (98%)	299 (99%)	3 (1%)	73	47
1	B	290/309 (94%)	288 (99%)	2 (1%)	81	61
All	All	592/618 (96%)	587 (99%)	5 (1%)	81	57

All (5) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	86[A]	ASP
1	A	86[B]	ASP
1	A	115	THR

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	95	THR
1	B	115	THR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	228	HIS
1	B	89	GLN
1	B	310	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
1	CSS	B	177	1	4,6,7	0.75	0	1,6,8	0.67	0
1	CSS	A	177	1	4,6,7	0.84	0	1,6,8	0.43	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	CSS	B	177	1	-	0/1/5/7	-
1	CSS	A	177	1	-	0/1/5/7	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	NAD	B	402	-	42,48,48	1.22	3 (7%)	50,73,73	1.21	4 (8%)
4	MRD	B	401	-	7,7,7	0.24	0	9,10,10	0.88	0
2	NAD	A	401	-	42,48,48	0.64	1 (2%)	50,73,73	1.00	3 (6%)
3	A1IGN	A	402	-	12,12,12	2.63	3 (25%)	14,15,15	0.87	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAD	B	402	-	-	4/26/62/62	0/5/5/5
4	MRD	B	401	-	-	0/5/5/5	-
2	NAD	A	401	-	-	4/26/62/62	0/5/5/5
3	A1IGN	A	402	-	-	0/8/8/8	0/1/1/1

All (7) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	402	A1IGN	C9-N2	7.01	1.44	1.34
2	B	402	NAD	C2N-N1N	5.07	1.41	1.35
3	A	402	A1IGN	C9-N1	4.83	1.42	1.35
3	A	402	A1IGN	O1-C9	-2.90	1.18	1.24
2	B	402	NAD	C8A-N7A	-2.36	1.30	1.34
2	B	402	NAD	PA-O2A	-2.18	1.45	1.55
2	A	401	NAD	C8A-N7A	-2.01	1.31	1.34

All (7) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	401	NAD	O4D-C1D-C2D	-4.43	100.45	106.93
2	B	402	NAD	O2A-PA-O1A	4.23	133.14	112.24
2	B	402	NAD	C6N-N1N-C2N	-3.19	119.07	121.97
2	B	402	NAD	C5A-C6A-N6A	2.85	124.69	120.35
2	B	402	NAD	O4D-C1D-C2D	-2.67	103.03	106.93
2	A	401	NAD	C6N-N1N-C2N	-2.51	119.68	121.97
2	A	401	NAD	C5A-C6A-N6A	2.14	123.60	120.35

There are no chirality outliers.

All (8) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	401	NAD	C5D-O5D-PN-O1N
2	B	402	NAD	C5D-O5D-PN-O1N
2	A	401	NAD	PA-O3-PN-O5D
2	B	402	NAD	PA-O3-PN-O5D
2	A	401	NAD	C5D-O5D-PN-O3
2	B	402	NAD	C5D-O5D-PN-O3
2	A	401	NAD	C5D-O5D-PN-O2N
2	B	402	NAD	C5D-O5D-PN-O2N


There are no ring outliers.


2 monomers are involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	402	NAD	1	0
3	A	402	A1IGN	3	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will

also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2			OWAB(Å ²)	Q<0.9
1	A	352/369 (95%)	1.07	44 (12%)	9	10	11, 27, 73, 124	11 (3%)
1	B	335/369 (90%)	0.72	26 (7%)	20	22	13, 27, 47, 83	8 (2%)
All	All	687/738 (93%)	0.90	70 (10%)	13	14	11, 27, 58, 124	19 (2%)

All (70) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	18	LEU	11.0
1	A	14	LEU	10.4
1	A	10	PRO	7.6
1	A	16	ALA	7.5
1	A	17	VAL	6.9
1	A	9	ALA	6.7
1	A	87	LEU	6.4
1	A	77	LEU	6.2
1	B	89	GLN	6.2
1	A	79	GLN	6.2
1	A	24	LEU	6.2
1	A	26	PRO	5.9
1	A	11	ALA	5.9
1	A	275	CYS	5.6
1	A	78	SER	5.6
1	A	15	ALA	5.3
1	B	94	LEU	5.1
1	A	13	ALA	5.0
1	A	286	VAL	4.8
1	A	91	ARG	4.8
1	B	85	ALA	4.8
1	A	19	LYS	4.7
1	A	362	PHE	4.6
1	A	22	SER	4.5

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	250	TYR	4.4
1	A	23	THR	4.4
1	A	25	PRO	4.2
1	A	21	SER	4.2
1	A	12	GLY	4.1
1	A	361	ALA	4.1
1	B	90	SER	4.0
1	B	28	SER	4.0
1	A	82	ASP	3.9
1	B	241[A]	LEU	3.9
1	A	20	HIS	3.9
1	B	251	LYS	3.7
1	A	32[A]	ARG	3.6
1	A	84	HIS	3.6
1	A	363	MET	3.6
1	A	274	LYS	3.6
1	A	305	TYR	3.6
1	B	88	THR	3.5
1	A	83	GLN	3.4
1	B	192	MET	3.2
1	A	85	ALA	3.1
1	B	363	MET	2.9
1	B	87	LEU	2.8
1	B	93	PRO	2.7
1	B	196	THR	2.6
1	A	332	VAL	2.6
1	A	64[A]	GLN	2.6
1	A	86[A]	ASP	2.5
1	B	77	LEU	2.5
1	B	91	ARG	2.5
1	B	84	HIS	2.5
1	A	272	PHE	2.5
1	A	76	PRO	2.5
1	A	195	ASN	2.4
1	B	305	TYR	2.4
1	B	74	LEU	2.3
1	A	360	ASP	2.3
1	B	200	LYS	2.2
1	A	254	GLY	2.1
1	B	173	ASN	2.1
1	B	191	VAL	2.1
1	B	75	GLU	2.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	361	ALA	2.1
1	A	49[A]	PHE	2.1
1	B	174	GLU	2.1
1	B	298	ASN	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
1	CSS	B	177	7/8	0.90	0.14	35,37,49,89	0
1	CSS	A	177	7/8	0.93	0.10	26,28,40,51	0

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	A1IGN	A	402	12/12	0.32	0.24	20,20,20,20	0
4	MRD	B	401	8/8	0.90	0.19	46,49,51,52	8
2	NAD	A	401	44/44	0.97	0.07	18,21,30,36	0
2	NAD	B	402	44/44	0.97	0.07	18,22,31,34	44




The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.


Electron density around A1IGN A 402:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around NAD A 401:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.5 Other polymers ⓘ

There are no such residues in this entry.