



Full wwPDB X-ray Structure Validation Report ⓘ

Jun 22, 2026 – 12:13 PM JST

PDB ID : 9L4M / pdb_00009l4m
Title : Structure of bifunctional Ylehd at pH 8 from Yarrowia lipolytica
Authors : Yadav, L.R.; Vanarase, K.N.; Inamdar, S.S.; Mande, S.C.; Kumar, A.R.
Deposited on : 2024-12-20
Resolution : 2.20 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION




The reported resolution of this entry is 2.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	6164 (2.20-2.20)
Clashscore	190562	6851 (2.20-2.20)
Ramachandran outliers	187476	6768 (2.20-2.20)
Sidechain outliers	187428	6769 (2.20-2.20)
RSRZ outliers	180081	6166 (2.20-2.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	352	 7% 89% 7% ..
1	B	352	 12% 83% 10% 7%
1	C	352	 23% 80% 15% 5%

2 Entry composition [i](#)


There are 3 unique types of molecules in this entry. The entry contains 8153 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Alpha/Beta hydrolase protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	340	Total 2764	C 1797	N 452	O 504	S 11	2	0	0
1	B	327	Total 2674	C 1742	N 443	O 479	S 10	1	0	0
1	C	333	Total 2654	C 1731	N 445	O 466	S 12	0	0	0

- Molecule 2 is GLYCEROL (CCD ID: GOL) (formula: C₃H₈O₃).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	C	O		
2	B	1	Total 6	C 3	O 3	0	0


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	27	Total 27	O 27	0	0
3	B	20	Total 20	O 20	0	0
3	C	8	Total 8	O 8	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Alpha/Beta hydrolase protein




- Molecule 1: Alpha/Beta hydrolase protein



- Molecule 1: Alpha/Beta hydrolase protein





4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	92.33Å 58.30Å 217.49Å 90.00° 94.64° 90.00°	Depositor
Resolution (Å)	72.26 – 2.20 72.26 – 2.20	Depositor EDS
% Data completeness (in resolution range)	99.6 (72.26-2.20) 99.9 (72.26-2.20)	Depositor EDS
R_{merge}	0.05	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.55 (at 2.20Å)	Xtrriage
Refinement program	PHENIX v1.21.2.5419	Depositor
R, R_{free}	0.269 , 0.306 0.268 , 0.305	Depositor DCC
R_{free} test set	2838 reflections (4.81%)	wwPDB-VP
Wilson B-factor (Å ²)	42.6	Xtrriage
Anisotropy	0.836	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 37.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	8153	wwPDB-VP
Average B, all atoms (Å ²)	61.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.67% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GOL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.14	0/2863	0.29	0/3904
1	B	0.13	0/2770	0.29	0/3772
1	C	0.14	0/2752	0.27	0/3755
All	All	0.14	0/8385	0.28	0/11431

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2764	0	2564	20	0
1	B	2674	0	2491	23	0
1	C	2654	0	2428	52	0
2	B	6	0	8	0	0
3	A	27	0	0	3	0
3	B	20	0	0	0	0
3	C	8	0	0	1	0
All	All	8153	0	7491	94	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (94) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:291:ILE:HG21	1:C:335:ILE:HD11	1.50	0.93
1:A:299:LEU:HD22	3:A:421:HOH:O	1.78	0.83
1:B:85:MET:HE1	1:B:198:VAL:HG11	1.59	0.83
1:C:305:MET:HA	1:C:305:MET:HE3	1.61	0.82
1:C:184:TRP:CE3	1:C:303:MET:HE3	2.22	0.74
1:C:86:GLU:O	1:C:86:GLU:CD	2.34	0.69
1:C:136:PHE:HE2	1:C:149:VAL:HG11	1.56	0.69
1:B:293:SER:O	1:B:295:ASN:N	2.25	0.68
1:C:290:TYR:HD1	1:C:291:ILE:N	1.92	0.68
1:C:136:PHE:CE2	1:C:149:VAL:HG11	2.30	0.66
1:C:290:TYR:HE2	1:C:304:SER:CB	2.09	0.66
1:B:192:GLN:NE2	1:B:192:GLN:HA	2.12	0.64
1:C:290:TYR:HE2	1:C:304:SER:HG	1.47	0.62
1:C:86:GLU:OE1	1:C:86:GLU:C	2.42	0.62
1:A:296:ASP:OD1	1:A:297:ILE:N	2.33	0.62
1:A:85:MET:HE1	1:A:198:VAL:HG11	1.81	0.62
1:C:134:ALA:HB3	1:C:136:PHE:HE1	1.63	0.62
1:A:228:ALA:HB3	1:A:229:PRO:HD3	1.83	0.61
1:B:192:GLN:HA	1:B:192:GLN:HE21	1.66	0.60
1:A:5:THR:OG1	1:C:205:MET:HE1	2.02	0.60
1:C:86:GLU:O	1:C:86:GLU:OE1	2.22	0.57
1:C:324:SER:OG	1:C:325:HIS:N	2.34	0.57
1:C:319:GLU:HB3	1:C:338:ILE:HD13	1.85	0.57
1:B:118:PHE:CE1	1:B:122:MET:HE3	2.39	0.56
1:C:136:PHE:HB3	1:C:146:VAL:HG13	1.87	0.56
1:C:290:TYR:HE2	1:C:304:SER:OG	1.87	0.56
1:A:116:LYS:NZ	1:A:120:ASP:OD2	2.38	0.56
1:C:177:LEU:HD13	1:C:261:ARG:CD	2.36	0.56
1:C:41:LEU:HD11	1:C:66:ILE:HG23	1.86	0.56
1:B:110:LEU:HG	1:B:263:LEU:HD13	1.87	0.56
1:C:38:VAL:HG23	1:C:134:ALA:CB	2.37	0.54
1:C:316:LEU:HD23	1:C:317:THR:N	2.22	0.54
1:C:85:MET:HE1	1:C:198:VAL:HG11	1.89	0.54
1:A:321:VAL:HG23	1:A:323:THR:OG1	2.10	0.52
1:C:27:ASP:OD2	1:C:65:ARG:NH2	2.43	0.52
1:B:1:MET:O	1:B:2:ALA:HB3	2.10	0.51
1:B:52:TRP:O	1:B:56:ILE:HD12	2.11	0.50
1:B:128:LYS:N	1:B:128:LYS:HD3	2.25	0.50
1:A:197:ALA:O	1:A:201:ILE:HD12	2.12	0.49
1:A:201:ILE:HD12	1:A:201:ILE:N	2.27	0.49

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:296:ASP:OD1	1:C:297:ILE:N	2.46	0.49
1:C:184:TRP:CZ3	1:C:303:MET:HA	2.48	0.48
1:C:55:GLN:NE2	1:C:328:LEU:HD12	2.28	0.48
1:A:197:ALA:O	1:A:200:ASN:N	2.47	0.48
1:C:86:GLU:CD	1:C:86:GLU:C	2.81	0.48
1:A:157:VAL:HG11	1:A:160:ILE:HD11	1.95	0.48
1:B:85:MET:HE2	1:B:259:ARG:CZ	2.44	0.47
1:C:136:PHE:CB	1:C:146:VAL:HG13	2.43	0.47
1:B:46:PRO:O	1:B:214:TYR:OH	2.30	0.47
1:C:85:MET:HE1	1:C:198:VAL:CG1	2.45	0.47
1:A:93:ASP:OD1	1:A:93:ASP:N	2.47	0.47
1:C:167:TYR:HE1	1:C:169:VAL:HG22	1.79	0.47
1:B:44:GLY:HA3	1:B:140:ASP:HB3	1.97	0.46
1:C:332:PRO:HA	1:C:335:ILE:HG22	1.97	0.46
1:B:45:PHE:HE1	1:B:193:PHE:CE2	2.33	0.46
1:C:135:THR:HG23	1:C:159:ALA:HB3	1.97	0.46
1:C:44:GLY:HA3	1:C:140:ASP:HB3	1.98	0.45
1:C:134:ALA:HB3	1:C:136:PHE:CE1	2.48	0.45
1:C:136:PHE:CD1	1:C:136:PHE:N	2.85	0.44
1:C:166:TYR:CE2	1:C:303:MET:HB2	2.52	0.44
1:A:165:TRP:CD1	1:A:166:TYR:H	2.35	0.44
1:C:136:PHE:N	1:C:136:PHE:HD1	2.16	0.44
1:B:146:VAL:N	1:B:147:PRO:CD	2.81	0.44
1:C:55:GLN:HE21	1:C:328:LEU:HD12	1.83	0.43
1:C:126:LEU:HD22	1:C:131:ILE:HG21	1.99	0.43
1:B:110:LEU:HD12	1:B:110:LEU:O	2.18	0.43
1:A:85:MET:HE2	1:A:259:ARG:CZ	2.47	0.43
1:C:125:LEU:O	1:C:129:LEU:HD22	2.19	0.43
1:C:38:VAL:HG12	1:C:65:ARG:HB3	2.01	0.43
1:C:41:LEU:HD12	1:C:41:LEU:N	2.34	0.42
1:B:37:ARG:NH1	1:B:346:LEU:HD21	2.34	0.42
1:C:45:PHE:CD1	1:C:46:PRO:HA	2.54	0.42
1:A:215:THR:HG21	1:A:225:ASP:HB2	2.01	0.42
1:C:174:TRP:HB2	1:C:265:TRP:CZ2	2.55	0.42
1:B:73:GLY:O	1:B:254:PRO:HD3	2.19	0.42
1:A:269:LYS:NZ	1:A:274:ASP:OD1	2.53	0.42
1:B:85:MET:HB3	1:B:98:TYR:CD2	2.55	0.42
1:B:169:VAL:HG11	1:B:310:TYR:CG	2.54	0.42
1:C:126:LEU:HD22	1:C:131:ILE:CG2	2.50	0.42
1:A:97:ILE:HD13	1:A:97:ILE:HA	1.94	0.41
1:A:148:LYS:NZ	3:A:408:HOH:O	2.53	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:346:LEU:HD12	1:B:346:LEU:O	2.20	0.41
1:C:174:TRP:HB2	1:C:265:TRP:CH2	2.55	0.41
1:B:195:GLY:O	1:B:198:VAL:HG12	2.21	0.41
1:C:249:ARG:O	3:C:401:HOH:O	2.22	0.41
1:B:300:ILE:HG13	1:B:303:MET:HE3	2.03	0.41
1:A:197:ALA:O	1:A:198:VAL:C	2.64	0.41
1:B:7:GLU:OE1	1:B:7:GLU:N	2.50	0.41
1:C:41:LEU:HD11	1:C:66:ILE:CG2	2.49	0.40
1:A:165:TRP:CD1	3:A:421:HOH:O	2.74	0.40
1:C:86:GLU:OE2	1:C:99:THR:OG1	2.24	0.40
1:C:233:THR:HG22	1:C:236:GLU:OE1	2.21	0.40
1:C:300:ILE:O	1:C:303:MET:HG3	2.22	0.40
1:C:290:TYR:C	1:C:290:TYR:CD1	2.99	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	334/352 (95%)	315 (94%)	16 (5%)	3 (1%)	14	14
1	B	321/352 (91%)	307 (96%)	12 (4%)	2 (1%)	21	23
1	C	327/352 (93%)	310 (95%)	17 (5%)	0	100	100
All	All	982/1056 (93%)	932 (95%)	45 (5%)	5 (0%)	24	27

All (5) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	294	THR
1	A	228	ALA
1	A	230	MET
1	A	198	VAL

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	46	PRO

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	287/310 (93%)	287 (100%)	0	100	100
1	B	275/310 (89%)	274 (100%)	1 (0%)	84	92
1	C	262/310 (84%)	262 (100%)	0	100	100
All	All	824/930 (89%)	823 (100%)	1 (0%)	88	95

All (1) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	130	ASN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	192	GLN
1	B	192	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	GOL	B	401	-	5,5,5	0.33	0	5,5,5	0.34	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	GOL	B	401	-	-	2/4/4/4	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (2) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	B	401	GOL	C1-C2-C3-O3
2	B	401	GOL	O2-C2-C3-O3

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	340/352 (96%)	0.89	23 (6%) 23 20	44, 53, 70, 125	1 (0%)
1	B	327/352 (92%)	1.05	42 (12%) 7 5	41, 52, 78, 91	1 (0%)
1	C	333/352 (94%)	1.48	82 (24%) 2 1	50, 76, 98, 119	0
All	All	1000/1056 (94%)	1.14	147 (14%) 6 4	41, 58, 91, 125	2 (0%)

All (147) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	299	LEU	6.7
1	B	298	ALA	6.2
1	B	297	ILE	6.1
1	B	296	ASP	5.0
1	A	2	ALA	5.0
1	A	226	PRO	4.9
1	A	225	ASP	4.9
1	C	300	ILE	4.6
1	B	165	TRP	4.5
1	C	231	HIS	4.4
1	A	298	ALA	4.3
1	C	229	PRO	4.3
1	C	97	ILE	4.3
1	B	300	ILE	4.2
1	A	224	PRO	4.2
1	C	299	LEU	4.2
1	C	308	ASP	4.2
1	C	349	THR	4.2
1	B	349	THR	4.2
1	B	186	THR	3.9
1	C	228	ALA	3.7
1	B	1	MET	3.7
1	A	95	ILE	3.7

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	C	219	ASN	3.7
1	B	322	PRO	3.7
1	B	191	LEU	3.6
1	C	1	MET	3.6
1	C	288	TYR	3.5
1	B	194	GLY	3.5
1	B	321	VAL	3.5
1	B	122	MET	3.5
1	C	290	TYR	3.5
1	A	228	ALA	3.4
1	A	229	PRO	3.4
1	C	160	ILE	3.4
1	C	297	ILE	3.4
1	B	184	TRP	3.3
1	A	219	ASN	3.3
1	C	165	TRP	3.3
1	C	310	TYR	3.2
1	C	347	ASP	3.2
1	B	301	PRO	3.2
1	B	2	ALA	3.2
1	C	227	GLU	3.2
1	C	273	LEU	3.1
1	B	216	VAL	3.1
1	C	321	VAL	3.1
1	C	136	PHE	3.0
1	C	218	ALA	3.0
1	B	187	THR	3.0
1	B	294	THR	3.0
1	A	231	HIS	3.0
1	A	87	GLU	3.0
1	A	4	LEU	3.0
1	C	339	TYR	3.0
1	B	324	SER	3.0
1	C	346	LEU	2.9
1	B	295	ASN	2.9
1	C	66	ILE	2.8
1	B	290	TYR	2.8
1	B	193	PHE	2.8
1	C	320	HIS	2.8
1	C	340	VAL	2.8
1	C	327	ALA	2.8
1	B	169	VAL	2.8

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	C	12	PHE	2.7
1	C	135	THR	2.7
1	C	291	ILE	2.7
1	C	307	LEU	2.6
1	C	31	GLU	2.6
1	A	94	GLY	2.6
1	C	342	TRP	2.6
1	B	219	ASN	2.6
1	B	215	THR	2.6
1	C	317	THR	2.6
1	B	303	MET	2.6
1	C	305	MET	2.6
1	C	71	LEU	2.5
1	C	32	GLY	2.5
1	C	56	ILE	2.5
1	A	230	MET	2.5
1	B	97	ILE	2.5
1	C	6	LYS	2.5
1	B	325	HIS	2.5
1	C	326	TRP	2.5
1	C	226	PRO	2.5
1	A	92	GLU	2.4
1	C	131	ILE	2.4
1	C	129	LEU	2.4
1	C	34	ASP	2.4
1	A	227	GLU	2.4
1	A	325	HIS	2.4
1	B	320	HIS	2.4
1	C	312	GLU	2.4
1	C	319	GLU	2.4
1	B	180	PHE	2.4
1	B	346	LEU	2.4
1	C	10	GLU	2.4
1	C	230	MET	2.3
1	B	231	HIS	2.3
1	C	282	LEU	2.3
1	C	40	VAL	2.3
1	C	173	GLU	2.3
1	C	311	VAL	2.3
1	C	85	MET	2.3
1	A	45	PHE	2.2
1	C	285	ASN	2.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	C	20	GLY	2.2
1	A	299	LEU	2.2
1	C	155	GLU	2.2
1	C	171	GLU	2.2
1	C	84	ALA	2.2
1	C	279	LYS	2.2
1	C	166	TYR	2.2
1	C	177	LEU	2.2
1	B	185	PRO	2.2
1	C	172	PRO	2.2
1	C	145	TYR	2.2
1	C	184	TRP	2.2
1	B	302	GLU	2.2
1	C	163	ALA	2.2
1	A	208	PHE	2.1
1	B	188	LYS	2.1
1	C	306	HIS	2.1
1	C	233	THR	2.1
1	B	98	TYR	2.1
1	B	330	GLU	2.1
1	C	45	PHE	2.1
1	C	119	ALA	2.1
1	C	57	PRO	2.1
1	C	157	VAL	2.1
1	B	107	CYS	2.1
1	A	327	ALA	2.1
1	C	348	LYS	2.1
1	C	28	ILE	2.1
1	A	349	THR	2.0
1	C	331	ALA	2.0
1	C	130	ASN	2.0
1	A	138	GLY	2.0
1	C	183	LYS	2.0
1	B	326	TRP	2.0
1	C	150	TRP	2.0
1	C	192	GLN	2.0
1	B	4	LEU	2.0
1	C	101	LEU	2.0
1	C	289	LEU	2.0
1	C	30	PRO	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	GOL	B	401	6/6	0.71	0.16	80,81,83,87	0

6.5 Other polymers [i](#)

There are no such residues in this entry.