

Full wwPDB X-ray Structure Validation Report (i)

Dec 22, 2025 – 04:35 PM JST

PDB ID : 9L7A / pdb 00009l7a

Title: Serine Beta-Lactamase OXA-48 in complex with Dual MBL/SBL inhibitor

B32

Authors : Li, G.-B.; Liang, G.-Q.

Deposited on : 2024-12-26

Resolution : 1.50 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org
A user guide is available at

https://www.wwpdb.org/validation/2017/XrayValidationReportHelp with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0

Mogul : 1.8.5 (274361), CSD as541be (2020)

Xtriage (Phenix) : 2.0 EDS : 3.0

buster-report : 1.1.7 (2018)

Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)

CCP4 : 9.0.010 (Gargrove)

Density-Fitness : 1.0.12

Ideal geometry (proteins) : Engh & Huber (2001) Ideal geometry (DNA, RNA) : Parkinson et al. (1996)


Validation Pipeline (wwPDB-VP) : 2.47

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: X-RAY DIFFRACTION

The reported resolution of this entry is 1.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive	Similar resolution
Metric	$(\# \mathrm{Entries})$	$(\# ext{Entries}, ext{ resolution range}(ext{Å}))$
R_{free}	164625	3717 (1.50-1.50)
Clashscore	180529	4048 (1.50-1.50)
Ramachandran outliers	177936	3970 (1.50-1.50)
Sidechain outliers	177891	3967 (1.50-1.50)
RSRZ outliers	164620	3718 (1.50-1.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain						
1	A	242	93%	6% •					
1	В	242	7% 94%	5% •					
1	С	242	7% 90%	9% •					
1	D	242	92%	7%					



2 Entry composition (i)


There are 3 unique types of molecules in this entry. The entry contains 16149 atoms, of which 7776 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called Beta-lactamase.

Mol	Chain	Residues		Atoms						AltConf	Trace
1	Λ	242	Total	С	Н	N	О	S	24	0	0
1	A	242	3916	1262	1933	349	365	7	24	U	
1	В	242	Total	С	Н	N	О	S	24	0	0
1	Б	242	3916	1262	1933	349	365	7			
1	С	242	Total	С	Н	N	О	S	24	0	0
1		242	3916	1262	1933	349	365	7	24	U	
1	D	242	Total	С	Н	N	О	S	24	0	0
1		242	3916	1262	1933	349	365	7	24	U	U

• Molecule 2 is $(3 \{R\})$ -3- $[(3 \{R\})$ -4-fluoranyl-1-oxidanyl-3 $\{H\}$ -2,1-benzoxaborol-3-yl]-2-met hylidene-butanoic acid (CCD ID: A1EKC) (formula: $C_{12}H_{12}BFO_4$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf		
2	Λ	1	Total	В	С	F	Η	О	0	0	
	A	1	29	1	12	1	11	4	U		
2	D	1	Total	В	С	F	Н	О	0	0	
2	Б	1	29		12	1	11	4	U	U	



Continued from previous page...

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf			
2	C	1	Total	В	С	F	Н	О	0	0		
2		1	29	1	12	1	11	4	U			
9	2 D	D	D	1	Total	В	С	F	Н	О	0	0
2		$D \mid I \mid$	29	1	12	1	11	4	U	U		


• Molecule 3 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	110	Total O 110 110	0	0
3	В	88	Total O 88 88	0	0
3	С	111	Total O 111 111	0	0
3	D	60	Total O 60 60	0	0



3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.





4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants	58.40Å 106.33Å 94.10Å	Depositor
a, b, c, α , β , γ	90.00° 107.72° 90.00°	Depositor
Resolution (Å)	19.86 - 1.50	Depositor
Resolution (A)	19.86 - 1.50	EDS
% Data completeness	94.2 (19.86-1.50)	Depositor
(in resolution range)	94.7 (19.86-1.50)	EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) > 1$	26.41 (at 1.49Å)	Xtriage
Refinement program	PHENIX (1.10.1_2155: ???)	Depositor
R, R_{free}	0.208 , 0.225	Depositor
it, it free	0.208 , 0.225	DCC
R_{free} test set	2015 reflections (1.15%)	wwPDB-VP
Wilson B-factor (Å ²)	17.8	Xtriage
Anisotropy	0.238	Xtriage
Bulk solvent $k_{sol}(e/Å^3)$, $B_{sol}(Å^2)$	0.49, 38.7	EDS
L-test for twinning ²	$< L >=0.50, < L^2>=0.33$	Xtriage
Estimated twinning fraction	0.021 for h,-k,-h-l	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	16149	wwPDB-VP
Average B, all atoms (Å ²)	21.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 4.86% of the height of the origin peak. No significant pseudotranslation is detected.

²Theoretical values of <|L|>, $<L^2>$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



¹Intensities estimated from amplitudes.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: KCX, A1EKC

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bo	nd lengths	Bond angles		
MIOI		RMSZ	# Z > 5	RMSZ	# Z >5	
1	A	0.56	$2/2019 \ (0.1\%)$	0.70	4/2730 (0.1%)	
1	В	0.62	4/2019~(0.2%)	0.69	1/2730 (0.0%)	
1	С	0.60	4/2019 (0.2%)	0.75	3/2730 (0.1%)	
1	D	0.42	0/2019	0.59	0/2730	
All	All	0.56	10/8076 (0.1%)	0.68	8/10920 (0.1%)	

All (10) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(A)
1	A	202	ASP	CB-CG	-6.56	1.35	1.52
1	С	101	ASP	C-O	-6.29	1.16	1.24
1	A	118	SER	C-O	-6.16	1.15	1.23
1	С	138	MET	SD-CE	-5.89	1.64	1.79
1	В	214	ARG	CZ-NH2	-5.40	1.26	1.33
1	В	213	THR	C-O	-5.38	1.17	1.24
1	В	212	SER	C-O	-5.32	1.18	1.24
1	В	215	ILE	C-O	-5.29	1.18	1.23
1	С	182	HIS	N-CA	-5.24	1.39	1.46
1	С	183	VAL	N-CA	-5.09	1.40	1.46

All (8) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$Observed(^o)$	$\operatorname{Ideal}({}^{o})$
1	A	202	ASP	CB-CA-C	-8.05	95.99	109.03
1	С	182	HIS	CA-C-O	-7.93	108.73	119.12
1	С	138	MET	CG-SD-CE	-6.41	86.81	100.90
1	С	216	GLU	N-CA-C	5.67	117.03	109.83
1	В	216	GLU	C-N-CD	5.56	132.83	120.60
1	A	221	TRP	CA-CB-CG	5.38	123.81	113.60
1	A	138	MET	CG-SD-CE	-5.25	89.35	100.90



Continued from previous page...

Mol	Chain	Res	Type	Atoms	\mathbf{Z}	$Observed(^o)$	$\operatorname{Ideal}({}^{o})$
1	A	33	ALA	N-CA-C	-5.21	106.07	112.90

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1983	1933	1931	15	2
1	В	1983	1933	1931	20	0
1	С	1983	1933	1931	23	0
1	D	1983	1933	1931	14	2
2	A	18	11	0	2	0
2	В	18	11	0	2	0
2	С	18	11	0	2	0
2	D	18	11	0	2	0
3	A	110	0	0	5	0
3	В	88	0	0	6	0
3	С	111	0	0	4	0
3	D	60	0	0	1	0
All	All	8373	7776	7724	59	2

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (59) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	$\begin{array}{c} {\rm Interatomic} \\ {\rm distance} \ ({\rm \AA}) \end{array}$	$egin{aligned} ext{Clash} \ ext{overlap } (ext{Å}) \end{aligned}$
1:A:37:GLU:HG3	3:A:436:HOH:O	1.40	1.21
1:B:215:ILE:HG21	1:C:98:GLN:OE1	1.55	1.06
1:A:64:GLN:NE2	3:A:402:HOH:O	1.83	1.06
1:A:60:LYS:HD2	1:B:146:ASN:HA	1.50	0.94
1:B:215:ILE:CG2	1:C:98:GLN:OE1	2.18	0.91
1:B:215:ILE:HD11	3:B:455:HOH:O	1.71	0.89
1:C:94:LYS:NZ	1:C:108:ASP:OD2	2.13	0.81



Continued from previous page...

At any 1		Interatomic	Clash	
Atom-1	Atom-2	${\rm distance} \ (\mathring{\rm A})$	overlap(Å)	
1:A:60:LYS:HD2	1:B:146:ASN:CA	2.11	0.81	
1:C:134:ARG:O	1:C:138:MET:HG2	1.85	0.76	
1:C:51:LYS:HE2	3:C:482:HOH:O	1.87	0.72	
1:A:60:LYS:CD	1:B:146:ASN:HA	2.20	0.72	
1:A:64:GLN:HG3	3:A:452:HOH:O	1.91	0.70	
3:B:401:HOH:O	1:C:216:GLU:HG2	1.91	0.69	
1:C:138:MET:SD	3:C:404:HOH:O	2.52	0.67	
1:B:215:ILE:CD1	3:B:412:HOH:O	2.41	0.67	
1:B:98:GLN:OE1	3:B:402:HOH:O	2.12	0.65	
1:A:64:GLN:CG	3:A:452:HOH:O	2.45	0.64	
1:B:215:ILE:HD13	3:B:412:HOH:O	1.96	0.63	
1:A:60:LYS:HD2	1:B:146:ASN:CB	2.29	0.62	
1:B:94:LYS:HE3	3:B:435:HOH:O	2.02	0.60	
1:A:60:LYS:HD2	1:B:146:ASN:HB2	1.87	0.56	
1:D:39:LYS:HE3	1:D:39:LYS:HA	1.88	0.55	
1:D:104:THR:HG23	3:D:456:HOH:O	2.06	0.55	
1:D:118:SER:OG	1:D:208:LYS:NZ	2.22	0.55	
1:B:215:ILE:HG12	1:C:98:GLN:HG3	1.88	0.54	
1:D:118:SER:HG	1:D:208:LYS:HZ1	1.50	0.54	
1:B:215:ILE:HG23	1:C:98:GLN:OE1	2.08	0.53	
1:C:180:LYS:HE3	3:C:488:HOH:O	2.11	0.51	
1:C:24:GLU:O	1:C:24:GLU:HG2	2.10	0.51	
1:A:37:GLU:CG	3:A:436:HOH:O	2.20	0.51	
1:C:259:LYS:HE2	1:D:256:GLU:OE1	2.11	0.50	
1:A:70:SER:HB2	2:A:301:A1EKC:O06	2.13	0.48	
1:D:134:ARG:O	1:D:138:MET:HG2	2.14	0.47	
1:B:27:GLU:HA	1:B:56:THR:O	2.14	0.47	
1:B:215:ILE:HG12	1:C:98:GLN:CG	2.44	0.46	
1:C:49:GLU:CB	1:C:232:VAL:HG13	2.45	0.46	
1:A:70:SER:CB	2:A:301:A1EKC:O06	2.63	0.46	
1:C:70:SER:OG	2:C:301:A1EKC:C07	2.63	0.46	
1:D:49:GLU:CB	1:D:232:VAL:HG13	2.46	0.46	
1:D:39:LYS:HE3	1:D:39:LYS:CA	2.44	0.45	
1:A:31:TRP:HB2	1:A:57:ASN:HB3	1.98	0.45	
1:A:220:GLY:O	1:A:238:ASN:HA	2.16	0.45	
1:D:212:SER:O	1:D:218:LYS:HA	2.18	0.43	
1:C:94:LYS:CE	1:C:108:ASP:OD2	2.66	0.43	
1:D:33:ALA:O	1:D:37:GLU:HG3	2.18	0.43	
1:B:70:SER:CB	2:B:301:A1EKC:O06	2.65	0.43	
1:B:70:SER:HB2	2:B:301:A1EKC:O06	2.17	0.43	
1:B:99:THR:O	1:C:216:GLU:HB2	2.18	0.43	



Continued from previous page...

Atom-1	Atom-2	Interatomic	Clash
Atom-1	Atom-2	${\rm distance}({\rm \AA})$	overlap (Å)
1:B:101:ASP:OD1	1:C:216:GLU:HG2	2.19	0.43
1:C:27:GLU:HA	1:C:56:THR:O	2.18	0.43
1:C:220:GLY:O	1:C:238:ASN:HA	2.19	0.43
1:D:220:GLY:O	1:D:238:ASN:HA	2.20	0.42
1:D:70:SER:CB	2:D:301:A1EKC:O06	2.68	0.42
1:C:31:TRP:HB2	1:C:57:ASN:HB3	2.02	0.41
1:C:70:SER:OG	2:C:301:A1EKC:O04	2.38	0.41
1:D:70:SER:HB2	2:D:301:A1EKC:O06	2.21	0.41
1:C:107:ARG:HD3	3:C:435:HOH:O	2.20	0.40
1:A:211:TYR:CE1	1:A:213:THR:HG22	2.56	0.40
1:D:24:GLU:O	1:D:24:GLU:HG2	2.21	0.40

All (2) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	$\begin{array}{c} {\rm Interatomic} \\ {\rm distance} \ ({\rm \AA}) \end{array}$	$\begin{array}{c} \text{Clash} \\ \text{overlap } (\text{\AA}) \end{array}$
1:A:202:ASP:OD1	1:D:39:LYS:HZ3[2_658]	1.40	0.20
1:A:202:ASP:OD1	1:D:39:LYS:NZ[2_658]	2.02	0.18

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Perce	ntiles
1	A	$239/242 \ (99\%)$	233 (98%)	6 (2%)	0	100	100
1	В	$239/242 \ (99\%)$	233 (98%)	6 (2%)	0	100	100
1	C	$239/242 \ (99\%)$	232 (97%)	7 (3%)	0	100	100
1	D	$239/242 \ (99\%)$	236 (99%)	3 (1%)	0	100	100
All	All	956/968~(99%)	934 (98%)	22 (2%)	0	100	100

There are no Ramachandran outliers to report.



5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric Outliers		Percentiles		
1	A	$210/210 \; (100\%)$	208 (99%)	2 (1%)	73 53		
1	В	210/210 (100%)	208 (99%)	2 (1%)	73 53		
1	С	210/210 (100%)	210 (100%)	0	100 100		
1	D	210/210 (100%)	207 (99%)	3 (1%)	62 38		
All	All	840/840 (100%)	833 (99%)	7 (1%)	79 62		

All (7) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	60	LYS
1	A	82	ASP
1	В	82	ASP
1	В	101	ASP
1	D	60	LYS
1	D	87	LYS
1	D	232	VAL

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	D	64	GLN
1	D	90	HIS

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

4 non-standard protein/DNA/RNA residues are modelled in this entry.



In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol Type		Chain	Chain	Chain	n Res	Res Link	В	Bond lengths			Bond angles		
MIOI	Type	Chain	nes	LIIIK	Counts	RMSZ	# Z >2	Counts	RMSZ	# Z > 2			
1	KCX	D	73	1	9,11,12	0.71	0	5,12,14	0.71	0			
1	KCX	С	73	1	9,11,12	0.74	0	5,12,14	0.75	0			
1	KCX	A	73	1	9,11,12	1.28	2 (22%)	5,12,14	1.39	1 (20%)			
1	KCX	В	73	1	9,11,12	0.88	0	5,12,14	1.06	0			

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	KCX	D	73	1	-	0/9/10/12	-
1	KCX	С	73	1	-	0/9/10/12	-
1	KCX	A	73	1	-	0/9/10/12	-
1	KCX	В	73	1	-	0/9/10/12	-

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	73	KCX	OQ1-CX	-2.13	1.17	1.21
1	A	73	KCX	CX-NZ	-2.09	1.31	1.35

All (1) bond angle outliers are listed below:

\mathbf{Mol}	Chain	Res	Type	Atoms	\mathbf{Z}	$\mathbf{Observed}(^{o})$	$\mathbf{Ideal}(^{o})$
1	A	73	KCX	CE-NZ-CX	2.51	125.92	121.89

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.



5.5 Carbohydrates (i)

There are no oligosaccharides in this entry.

5.6 Ligand geometry (i)

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
	Type	Chain	nes	LIIIK	Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	A1EKC	D	301	-	17,19,19	1.98	5 (29%)	18,28,28	3.69	10 (55%)
2	A1EKC	С	301	-	17,19,19	2.34	4 (23%)	18,28,28	2.44	9 (50%)
2	A1EKC	A	301	-	17,19,19	1.92	4 (23%)	18,28,28	3.01	9 (50%)
2	A1EKC	В	301	-	17,19,19	1.79	5 (29%)	18,28,28	3.33	8 (44%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	A1EKC	D	301	-	-	2/12/24/24	0/2/2/2
2	A1EKC	С	301	-	-	2/12/24/24	0/2/2/2
2	A1EKC	A	301	-	-	2/12/24/24	0/2/2/2
2	A1EKC	В	301	-	-	2/12/24/24	0/2/2/2

All (18) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	\mathbf{Z}	$\operatorname{Observed}(\text{\AA})$	$\operatorname{Ideal}(ext{\AA})$
2	D	301	A1EKC	C02-C03	-5.21	1.47	1.54
2	С	301	A1EKC	B05-O04	4.85	1.42	1.38
2	A	301	A1EKC	C02-C03	-4.76	1.48	1.54
2	С	301	A1EKC	C02-C03	-4.58	1.48	1.54
2	С	301	A1EKC	C15-C14	-3.88	1.25	1.32
2	В	301	A1EKC	C15-C14	-3.81	1.25	1.32



Continued from previous page...

Mol	Chain	Res	Type	Atoms	\mathbf{Z}	Observed(A)	$\operatorname{Ideal}(\text{\AA})$
2	A	301	A1EKC	O17-C16	-3.35	1.20	1.30
2	С	301	A1EKC	O17-C16	-3.26	1.21	1.30
2	В	301	A1EKC	C02-C03	-3.21	1.50	1.54
2	A	301	A1EKC	C15-C14	-3.20	1.27	1.32
2	D	301	A1EKC	B05-O04	3.02	1.41	1.38
2	D	301	A1EKC	C15-C14	-2.88	1.27	1.32
2	В	301	A1EKC	B05-O04	2.62	1.40	1.38
2	D	301	A1EKC	C16-C14	2.37	1.51	1.49
2	В	301	A1EKC	O17-C16	-2.21	1.24	1.30
2	A	301	A1EKC	C09-C08	-2.11	1.34	1.38
2	В	301	A1EKC	C09-C10	-2.11	1.34	1.38
2	D	301	A1EKC	O04-C03	-2.06	1.42	1.45

All (36) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$\mathbf{Observed}(^o)$	$\operatorname{Ideal}({}^{o})$
2	D	301	A1EKC	O04-B05-C07	-9.81	102.85	108.43
2	В	301	A1EKC	O04-B05-C07	-9.21	103.20	108.43
2	A	301	A1EKC	O04-B05-C07	-6.76	104.59	108.43
2	С	301	A1EKC	O04-B05-C07	-5.69	105.19	108.43
2	В	301	A1EKC	O06-B05-O04	-5.64	110.20	121.96
2	D	301	A1EKC	B05-C07-C13	5.48	108.52	104.72
2	D	301	A1EKC	C10-C11-C13	-5.47	118.20	123.98
2	В	301	A1EKC	C10-C11-C13	-5.39	118.28	123.98
2	A	301	A1EKC	B05-C07-C13	5.29	108.39	104.72
2	A	301	A1EKC	O06-B05-O04	-5.13	111.27	121.96
2	A	301	A1EKC	C10-C11-C13	-4.82	118.89	123.98
2	В	301	A1EKC	B05-C07-C13	4.69	107.97	104.72
2	D	301	A1EKC	O17-C16-C14	4.19	119.48	114.44
2	D	301	A1EKC	O18-C16-C14	-4.17	117.84	122.79
2	С	301	A1EKC	O06-B05-O04	-4.17	113.27	121.96
2	D	301	A1EKC	O06-B05-O04	-4.00	113.61	121.96
2	D	301	A1EKC	C11-C13-C07	3.92	122.14	119.22
2	С	301	A1EKC	O18-C16-C14	-3.22	118.97	122.79
2	С	301	A1EKC	C10-C11-C13	-3.15	120.65	123.98
2	A	301	A1EKC	O18-C16-C14	-2.92	119.33	122.79
2	В	301	A1EKC	O18-C16-C14	-2.77	119.50	122.79
2	A	301	A1EKC	O17-C16-C14	2.70	117.68	114.44
2	D	301	A1EKC	C09-C08-C07	-2.69	118.35	121.69
2	С	301	A1EKC	B05-C07-C13	2.57	106.50	104.72
2	A	301	A1EKC	B05-C07-C08	-2.54	130.46	135.01
2	В	301	A1EKC	C11-C13-C07	2.43	121.03	119.22



Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	$Observed(^o)$	$\operatorname{Ideal}({}^{o})$
2	D	301	A1EKC	B05-C07-C08	-2.43	130.66	135.01
2	С	301	A1EKC	F12-C11-C13	-2.36	115.82	118.13
2	С	301	A1EKC	B05-O04-C03	-2.34	103.85	106.52
2	D	301	A1EKC	F12-C11-C10	2.28	123.68	118.59
2	A	301	A1EKC	F12-C11-C10	2.25	123.60	118.59
2	С	301	A1EKC	F12-C11-C10	2.21	123.53	118.59
2	В	301	A1EKC	F12-C11-C13	2.21	120.29	118.13
2	С	301	A1EKC	C09-C08-C07	-2.18	118.99	121.69
2	В	301	A1EKC	C09-C08-C07	-2.10	119.08	121.69
2	A	301	A1EKC	C08-C07-C13	2.06	121.02	119.09

There are no chirality outliers.

All (8) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	301	A1EKC	C01-C02-C14-C16
2	A	301	A1EKC	C03-C02-C14-C16
2	В	301	A1EKC	C01-C02-C14-C16
2	С	301	A1EKC	C01-C02-C14-C16
2	D	301	A1EKC	C01-C02-C14-C16
2	В	301	A1EKC	C03-C02-C14-C16
2	С	301	A1EKC	C03-C02-C14-C16
2	D	301	A1EKC	C03-C02-C14-C16

There are no ring outliers.


4 monomers are involved in 8 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	D	301	A1EKC	2	0
2	С	301	A1EKC	2	0
2	A	301	A1EKC	2	0
2	В	301	A1EKC	2	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and




any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.
















5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ>2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95^{th} percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<rsrz></rsrz>	$\#\mathrm{RSRZ}{>}2$	$OWAB(A^2)$	Q<0.9
1	A	241/242 (99%)	-0.09	10 (4%) 42 45	13, 17, 33, 41	0
1	В	241/242 (99%)	0.04	18 (7%) 22 22	13, 19, 35, 44	0
1	С	241/242 (99%)	-0.03	16 (6%) 26 27	12, 17, 33, 43	0
1	D	241/242 (99%)	0.33	13 (5%) 32 34	16, 23, 37, 44	0
All	All	964/968 (99%)	0.07	57 (5%) 29 30	12, 19, 35, 44	0

All (57) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	С	215	ILE	8.0
1	В	25	TRP	5.8
1	С	25	TRP	5.7
1	D	37	GLU	4.8
1	С	213	THR	4.5
1	D	39	LYS	4.4
1	В	101	ASP	4.4
1	В	26	GLN	4.3
1	С	24	GLU	4.0
1	В	37	GLU	3.9
1	D	60	LYS	3.9
1	A	33	ALA	3.9
1	В	36	THR	3.9
1	A	25	TRP	3.8
1	D	36	THR	3.7
1	A	102	ILE	3.7
1	A	24	GLU	3.7
1	D	98	GLN	3.4
1	В	102	ILE	3.4
1	С	216	GLU	3.3
1	A	64	GLN	3.2



Continued from previous page...

Mol	$\frac{nuea\ jron}{\mathbf{Chain}}$	Res	Type	RSRZ
1	D	25	TRP	3.2
1	С	214	ARG	3.2
1	В	24	GLU	3.2
1	С	39	LYS	3.1
1	A	37	GLU	3.1
1	С	36	THR	3.0
1	С	37	GLU	3.0
1	С	182	HIS	2.8
1	С	107	ARG	2.8
1	D	24	GLU	2.8
1	A	60	LYS	2.8
1	D	102	ILE	2.8
1	В	98	GLN	2.8
1	В	215	ILE	2.7
1	В	216	GLU	2.7
1	A	202	ASP	2.6
1	С	230	ASP	2.6
1	В	213	THR	2.5
1	С	217	PRO	2.5
1	В	230	ASP	2.5
1	В	39	LYS	2.4
1	D	107	ARG	2.4
1	A	39	LYS	2.4
1	D	182	HIS	2.4
1	A	98	GLN	2.3
1	D	95	TRP	2.3
1	В	232	VAL	2.3
1	С	26	GLN	2.3
1	В	107	ARG	2.3
1	С	98	GLN	2.3
1	С	93	PHE	2.2
1	В	182	HIS	2.2
1	D	41	GLN	2.2
1	В	214	ARG	2.1
1	В	217	PRO	2.1
1	D	103	ALA	2.0

6.2 Non-standard residues in protein, DNA, RNA chains (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column



labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	$\mathbf{B} ext{-}\mathbf{factors}(\mathbf{\mathring{A}}^2)$	Q < 0.9
1	KCX	A	73	12/13	-	-	12,15,17,18	24
1	KCX	В	73	12/13	-	-	12,15,17,18	24
1	KCX	С	73	12/13	-	-	12,14,17,17	24
1	KCX	D	73	12/13	-	-	16,18,22,22	24

6.3 Carbohydrates (i)

There are no oligosaccharides in this entry.


6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	$\operatorname{B-factors}(\mathring{\mathbf{A}}^2)$	Q < 0.9
2	A1EKC	D	301	18/18	0.92	0.09	20,20,20,20	0
2	A1EKC	С	301	18/18	0.95	0.08	20,20,20,20	0
2	A1EKC	A	301	18/18	0.95	0.07	20,20,20,20	0
2	A1EKC	В	301	18/18	0.96	0.08	20,20,20,20	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



















6.5 Other polymers (i)

There are no such residues in this entry.

