



Full wwPDB X-ray Structure Validation Report ⓘ

Dec 22, 2025 – 08:57 PM JST

PDB ID : 9L8X / pdb_00009l8x
Title : The crystal structure of Halogenases XhnO11 complex with vanadate
Authors : Wu, L.; Fang, S.Q.; Nie, Q.Y.; Tang, G.L.
Deposited on : 2024-12-28
Resolution : 2.40 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 2.0
EDS : 3.0
buster-report : 1.1.7 (2018)
Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.47


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.40 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	164625	4642 (2.40-2.40)
Clashscore	180529	5218 (2.40-2.40)
Ramachandran outliers	177936	5158 (2.40-2.40)
Sidechain outliers	177891	5159 (2.40-2.40)
RSRZ outliers	164620	4642 (2.40-2.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	528	
1	B	528	

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 7511 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Vanadium-dependent haloperoxidase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	B	458	Total	C	N	O	S	0	0	0
			3565	2230	631	689	15			
1	A	458	Total	C	N	O	S	0	0	0
			3565	2230	631	689	15			

There are 56 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
B	-20	MET	-	initiating methionine	UNP A0A420V7T7
B	-19	GLY	-	expression tag	UNP A0A420V7T7
B	-18	SER	-	expression tag	UNP A0A420V7T7
B	-17	SER	-	expression tag	UNP A0A420V7T7
B	-16	HIS	-	expression tag	UNP A0A420V7T7
B	-15	HIS	-	expression tag	UNP A0A420V7T7
B	-14	HIS	-	expression tag	UNP A0A420V7T7
B	-13	HIS	-	expression tag	UNP A0A420V7T7
B	-12	HIS	-	expression tag	UNP A0A420V7T7
B	-11	HIS	-	expression tag	UNP A0A420V7T7
B	-10	SER	-	expression tag	UNP A0A420V7T7
B	-9	SER	-	expression tag	UNP A0A420V7T7
B	-8	GLY	-	expression tag	UNP A0A420V7T7
B	-7	LEU	-	expression tag	UNP A0A420V7T7
B	-6	VAL	-	expression tag	UNP A0A420V7T7
B	-5	GLY	-	expression tag	UNP A0A420V7T7
B	-4	ARG	-	expression tag	UNP A0A420V7T7
B	-3	PRO	-	expression tag	UNP A0A420V7T7
B	-2	SER	-	expression tag	UNP A0A420V7T7
B	-1	HIS	-	expression tag	UNP A0A420V7T7
B	0	MET	-	expression tag	UNP A0A420V7T7
B	78	VAL	ILE	conflict	UNP A0A420V7T7
B	254	LYS	GLU	conflict	UNP A0A420V7T7
B	262	LYS	GLU	conflict	UNP A0A420V7T7
B	285	GLU	ASP	conflict	UNP A0A420V7T7

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	377	LYS	ARG	conflict	UNP A0A420V7T7
B	423	ARG	LYS	conflict	UNP A0A420V7T7
B	463	LYS	ARG	conflict	UNP A0A420V7T7
A	-20	MET	-	initiating methionine	UNP A0A420V7T7
A	-19	GLY	-	expression tag	UNP A0A420V7T7
A	-18	SER	-	expression tag	UNP A0A420V7T7
A	-17	SER	-	expression tag	UNP A0A420V7T7
A	-16	HIS	-	expression tag	UNP A0A420V7T7
A	-15	HIS	-	expression tag	UNP A0A420V7T7
A	-14	HIS	-	expression tag	UNP A0A420V7T7
A	-13	HIS	-	expression tag	UNP A0A420V7T7
A	-12	HIS	-	expression tag	UNP A0A420V7T7
A	-11	HIS	-	expression tag	UNP A0A420V7T7
A	-10	SER	-	expression tag	UNP A0A420V7T7
A	-9	SER	-	expression tag	UNP A0A420V7T7
A	-8	GLY	-	expression tag	UNP A0A420V7T7
A	-7	LEU	-	expression tag	UNP A0A420V7T7
A	-6	VAL	-	expression tag	UNP A0A420V7T7
A	-5	GLY	-	expression tag	UNP A0A420V7T7
A	-4	ARG	-	expression tag	UNP A0A420V7T7
A	-3	PRO	-	expression tag	UNP A0A420V7T7
A	-2	SER	-	expression tag	UNP A0A420V7T7
A	-1	HIS	-	expression tag	UNP A0A420V7T7
A	0	MET	-	expression tag	UNP A0A420V7T7
A	78	VAL	ILE	conflict	UNP A0A420V7T7
A	254	LYS	GLU	conflict	UNP A0A420V7T7
A	262	LYS	GLU	conflict	UNP A0A420V7T7
A	285	GLU	ASP	conflict	UNP A0A420V7T7
A	377	LYS	ARG	conflict	UNP A0A420V7T7
A	423	ARG	LYS	conflict	UNP A0A420V7T7
A	463	LYS	ARG	conflict	UNP A0A420V7T7

- Molecule 2 is VANADATE ION (CCD ID: VO4) (formula: O₄V) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	B	1	Total	O	V	0	0
			5	4	1		
2	A	1	Total	O	V	0	0
			5	4	1		

- Molecule 3 is CHLORIDE ION (CCD ID: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	B	1	Total	Cl	0	0
			1	1		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	B	201	Total	O	0	0
			201	201		
4	A	169	Total	O	0	0
			169	169		


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Vanadium-dependent haloperoxidase



• Molecule 1: Vanadium-dependent haloperoxidase



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	87.89Å 94.03Å 126.03Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	47.02 – 2.40 47.02 – 2.40	Depositor EDS
% Data completeness (in resolution range)	99.9 (47.02-2.40) 99.9 (47.02-2.40)	Depositor EDS
R_{merge}	0.16	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.37 (at 2.39Å)	Xtriage
Refinement program	PHENIX 1.12_2829	Depositor
R, R_{free}	0.190 , 0.247 0.192 , 0.247	Depositor DCC
R_{free} test set	2107 reflections (5.07%)	wwPDB-VP
Wilson B-factor (Å ²)	27.6	Xtriage
Anisotropy	0.704	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 33.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.34$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	7511	wwPDB-VP
Average B, all atoms (Å ²)	31.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.44% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: CL, VO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.34	0/3653	0.54	0/4953
1	B	0.35	0/3653	0.53	0/4953
All	All	0.35	0/7306	0.54	0/9906

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	319	ILE	Peptide

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3565	0	3419	36	0
1	B	3565	0	3418	39	0
2	A	5	0	0	0	0
2	B	5	0	0	0	0
3	B	1	0	0	0	0
4	A	169	0	0	4	0
4	B	201	0	0	2	0
All	All	7511	0	6837	73	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (73) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:218:THR:HG22	1:A:220:TYR:H	1.25	1.01
1:B:80:MET:HE3	1:B:306:ASN:HD21	1.45	0.81
1:A:80:MET:HE2	1:A:306:ASN:HD21	1.47	0.79
1:A:466:VAL:HG21	1:A:473:LYS:HD2	1.72	0.71
1:A:507:ARG:O	4:A:701:HOH:O	2.09	0.69
1:A:432:LYS:HE2	1:A:445:LYS:H	1.59	0.67
1:B:260:ASP:OD1	1:B:262:LYS:HD3	1.97	0.65
1:A:80:MET:CE	1:A:306:ASN:HD21	2.11	0.64
1:A:372:TRP:CZ3	1:A:391:TYR:HA	2.36	0.61
1:B:67:VAL:HG21	1:B:92:MET:HE1	1.83	0.60
1:B:427:ARG:HH11	1:B:450:ARG:HH21	1.48	0.59
1:B:488:GLU:HG3	1:B:489:SER:N	2.20	0.57
1:B:265:LEU:HD11	1:B:350:LYS:HD3	1.87	0.56
1:A:64:TYR:OH	1:A:93:GLU:OE1	2.24	0.55
1:B:429:ASP:OD1	1:B:448:SER:OG	2.16	0.55
1:B:80:MET:HE1	1:B:393:GLN:HB3	1.90	0.53
1:A:237:ASN:HB3	1:A:239:ILE:H	1.73	0.53
1:A:99:ALA:HB1	1:A:130:ALA:HB1	1.91	0.53
1:A:139:MET:HB3	1:A:150:TRP:CE2	2.45	0.51
1:A:332:ALA:HA	1:A:494:VAL:HG21	1.92	0.51
1:B:242:ALA:HB3	1:A:242:ALA:HB3	1.93	0.50
1:B:307:ASP:HB3	1:B:310:ILE:HB	1.92	0.50
1:B:429:ASP:CG	1:B:450:ARG:HH12	2.20	0.50
1:B:92:MET:HE2	1:B:135:THR:HG23	1.95	0.49
1:A:218:THR:HG22	1:A:220:TYR:N	2.09	0.49
1:B:181:ALA:O	1:B:185:LYS:HG2	2.12	0.49
1:A:225:PRO:HG2	1:A:274:HIS:CG	2.48	0.49

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:320:SER:O	1:A:321:ASP:HB2	2.12	0.49
1:A:237:ASN:ND2	4:A:717:HOH:O	2.47	0.48
1:A:78:VAL:HG11	1:A:80:MET:HE3	1.94	0.48
1:B:133:TYR:CE2	1:B:172:VAL:HG22	2.48	0.48
1:B:82:GLY:C	1:B:341:MET:HE2	2.38	0.48
1:B:82:GLY:O	1:B:86:LEU:HB2	2.14	0.48
1:B:151:ARG:NH2	1:B:162:ASP:OD1	2.44	0.48
1:A:302:ALA:HA	1:A:464:SER:HB2	1.97	0.46
1:B:413:GLU:HG2	1:B:484:ALA:O	2.16	0.46
1:B:219:PRO:HG2	1:B:220:TYR:CE2	2.51	0.45
1:B:87:ARG:NH1	1:B:191:MET:HE3	2.32	0.45
1:B:67:VAL:HG11	1:B:92:MET:HE1	1.98	0.45
1:A:82:GLY:C	1:A:341:MET:HE2	2.43	0.44
1:B:99:ALA:HB1	1:B:130:ALA:HB1	2.00	0.43
1:A:228:TRP:CD1	1:A:398:PRO:HA	2.53	0.43
1:A:185:LYS:NZ	4:A:725:HOH:O	2.50	0.43
1:A:424:THR:O	1:A:452:THR:HA	2.18	0.43
1:A:282:ARG:O	1:A:286:GLU:HG3	2.18	0.43
1:B:111:HIS:HB2	4:B:763:HOH:O	2.19	0.43
1:B:111:HIS:ND1	1:B:182:VAL:HG22	2.34	0.43
1:A:490:SER:O	1:A:494:VAL:HG23	2.19	0.43
1:B:78:VAL:O	1:B:80:MET:N	2.52	0.43
1:B:139:MET:HB3	1:B:150:TRP:CE2	2.54	0.42
1:B:269:PRO:HB3	1:B:400:TYR:CG	2.54	0.42
1:B:241:THR:HB	1:A:241:THR:HB	2.02	0.42
1:A:368:LYS:HD2	1:A:382:ASP:C	2.44	0.42
1:B:281:LYS:HE3	1:B:281:LYS:HB2	1.89	0.42
1:B:136:TYR:OH	4:B:702:HOH:O	2.13	0.42
1:B:253:ALA:O	1:B:342:LEU:HD11	2.20	0.42
1:B:302:ALA:HA	1:B:464:SER:HB2	2.01	0.42
1:B:191:MET:HE1	1:B:338:PHE:CE1	2.54	0.41
1:A:340:VAL:HG11	1:A:407:PHE:HA	2.02	0.41
1:B:347:ASN:HB3	1:B:401:PRO:HD3	2.02	0.41
1:B:372:TRP:CZ3	1:B:391:TYR:HA	2.55	0.41
1:B:427:ARG:HG2	1:B:450:ARG:HD2	2.03	0.41
1:A:395:SER:HB3	1:A:397:HIS:NE2	2.35	0.41
1:A:399:GLU:O	1:A:471:HIS:CE1	2.74	0.41
1:B:79:ALA:HB3	1:B:307:ASP:OD1	2.21	0.41
1:A:193:GLN:O	4:A:702:HOH:O	2.22	0.41
1:A:83:THR:N	1:A:341:MET:HE2	2.36	0.41
1:A:224:ASN:OD1	1:A:226:SER:HB2	2.21	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:464:SER:HA	1:A:467:TRP:CE3	2.56	0.41
1:B:150:TRP:O	1:B:153:MET:HB3	2.21	0.40
1:A:409:ALA:HB3	1:A:483:GLY:HA3	2.03	0.40
1:A:186:ARG:HA	1:A:186:ARG:HD3	1.90	0.40
1:B:356:PRO:HA	1:B:472:PHE:CZ	2.57	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	456/528 (86%)	443 (97%)	12 (3%)	1 (0%)	44	59
1	B	456/528 (86%)	441 (97%)	15 (3%)	0	100	100
All	All	912/1056 (86%)	884 (97%)	27 (3%)	1 (0%)	48	65

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	445	LYS

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	372/425 (88%)	364 (98%)	8 (2%)	47	67
1	B	372/425 (88%)	368 (99%)	4 (1%)	70	84
All	All	744/850 (88%)	732 (98%)	12 (2%)	58	76

All (12) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	268	LYS
1	B	300	LEU
1	B	488	GLU
1	B	505	SER
1	A	58	SER
1	A	166	THR
1	A	267	SER
1	A	300	LEU
1	A	308	LYS
1	A	488	GLU
1	A	492	ARG
1	A	501	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	471	HIS
1	A	149	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 3 ligands modelled in this entry, 1 is monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	VO4	B	601	4	1,4,4	4.85	1 (100%)	-		
2	VO4	A	601	1	1,4,4	4.66	1 (100%)	-		

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	601	VO4	O1-V	4.85	1.91	1.63
2	A	601	VO4	O1-V	4.66	1.90	1.63

There are no bond angle outliers.

There are no chirality outliers.


There are no torsion outliers.


There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier.

The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	458/528 (86%)	-0.09	3 (0%) 84 81	19, 31, 50, 71	0
1	B	458/528 (86%)	-0.33	2 (0%) 89 87	18, 27, 42, 66	0
All	All	916/1056 (86%)	-0.21	5 (0%) 87 85	18, 29, 47, 71	0

All (5) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	316	MET	3.2
1	A	316	MET	2.5
1	A	50	ALA	2.2
1	B	319	ILE	2.0
1	A	444	ALA	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	CL	B	602	1/1	0.90	0.10	47,47,47,47	0
2	VO4	B	601	5/5	0.97	0.10	27,29,33,36	0
2	VO4	A	601	5/5	0.98	0.07	26,28,31,31	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around VO4 B 601:

2mF_o-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around VO4 A 601:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.