



# Full wwPDB X-ray Structure Validation Report i

Jan 5, 2026 – 04:11 PM JST

PDB ID : 9LGR / pdb\_00009lgr  
Title : Y-degron fused ZZ-domain of the *Arabidopsis thaliana* E3 ubiquitin-protein ligase BIG  
Authors : Yang, W.S.; Lee, J.; Song, H.K.  
Deposited on : 2025-01-10  
Resolution : 2.05 Å (reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the i symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references](#) i) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20231227.v01 (using entries in the PDB archive December 27th 2023)
CCP4	:	9.0.011 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.47


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## X-RAY DIFFRACTION

The reported resolution of this entry is 2.05 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R <sub>free</sub>	164625	2096 (2.04-2.04)
Clashscore	180529	2229 (2.04-2.04)
Ramachandran outliers	177936	2217 (2.04-2.04)
Sidechain outliers	177891	2217 (2.04-2.04)
RSRZ outliers	164620	2096 (2.04-2.04)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5%. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.



## 2 Entry composition (i)

There are 4 unique types of molecules in this entry. The entry contains 5562 atoms, of which 2593 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Auxin transport protein BIG.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	59	Total	C	H	N	O	S	0	0	0
			905	293	436	82	86	8			
1	B	59	Total	C	H	N	O	S	0	0	0
			905	293	436	82	86	8			
1	C	59	Total	C	H	N	O	S	0	0	0
			905	293	436	82	86	8			
1	D	59	Total	C	H	N	O	S	0	0	0
			905	293	436	82	86	8			
1	E	57	Total	C	H	N	O	S	0	0	0
			883	286	427	80	82	8			
1	F	57	Total	C	H	N	O	S	0	0	0
			877	283	422	80	84	8			

There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	2611	TYR	-	expression tag	UNP Q9SRU2
B	2611	TYR	-	expression tag	UNP Q9SRU2
C	2611	TYR	-	expression tag	UNP Q9SRU2
D	2611	TYR	-	expression tag	UNP Q9SRU2
E	2611	TYR	-	expression tag	UNP Q9SRU2
F	2611	TYR	-	expression tag	UNP Q9SRU2

- Molecule 2 is ZINC ION (CCD ID: ZN) (formula: Zn) (labeled as "Ligand of Interest" by depositor).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	2	Total Zn 2 2	0	0
2	B	2	Total Zn 2 2	0	0
2	C	2	Total Zn 2 2	0	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	D	2	Total Zn 2 2	0	0
2	E	2	Total Zn 2 2	0	0
2	F	2	Total Zn 2 2	0	0

- Molecule 3 is SODIUM ION (CCD ID: NA) (formula: Na) (labeled as "Ligand of Interest" by depositor).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	D	1	Total Na 1 1	0	0
3	E	1	Total Na 1 1	0	0


- Molecule 4 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	26	Total O 26 26	0	0
4	B	36	Total O 36 36	0	0
4	C	33	Total O 33 33	0	0
4	D	27	Total O 27 27	0	0
4	E	27	Total O 27 27	0	0
4	F	19	Total O 19 19	0	0


### 3 Residue-property plots [\(i\)](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Auxin transport protein BIG




- Molecule 1: Auxin transport protein BIG




- Molecule 1: Auxin transport protein BIG





- Molecule 1: Auxin transport protein BIG



- Molecule 1: Auxin transport protein BIG



- Molecule 1: Auxin transport protein BIG



## 4 Data and refinement statistics i

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	23.78Å 93.83Å 80.88Å 90.00° 92.86° 90.00°	Depositor
Resolution (Å)	30.61 – 2.05 30.61 – 2.05	Depositor EDS
% Data completeness (in resolution range)	96.8 (30.61-2.05) 96.8 (30.61-2.05)	Depositor EDS
$R_{merge}$	0.15	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle^1$	1.98 (at 2.06Å)	Xtriage
Refinement program	PHENIX 1.18rc7_3834	Depositor
$R$ , $R_{free}$	0.192, 0.219 0.193, 0.220	Depositor DCC
$R_{free}$ test set	1993 reflections (9.00%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	18.8	Xtriage
Anisotropy	0.339	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.51, 60.1	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	0.067 for h,-k,-l	Xtriage
$F_o, F_c$ correlation	0.93	EDS
Total number of atoms	5562	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	28.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 7.19% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality i

### 5.1 Standard geometry i

Bond lengths and bond angles in the following residue types are not validated in this section: ZN, NA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.26	0/484	0.54	0/664
1	B	0.19	0/484	0.43	0/664
1	C	0.17	0/484	0.42	0/664
1	D	0.19	0/484	0.48	0/664
1	E	0.27	0/470	0.52	0/643
1	F	0.33	0/467	0.56	0/636
All	All	0.24	0/2873	0.49	0/3935

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts i

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	469	436	436	3	0
1	B	469	436	436	1	0
1	C	469	436	436	1	0
1	D	469	436	436	1	0
1	E	456	427	426	3	0
1	F	455	422	422	3	0
2	A	2	0	0	0	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	B	2	0	0	0	0
2	C	2	0	0	0	0
2	D	2	0	0	0	0
2	E	2	0	0	0	0
2	F	2	0	0	0	0
3	D	1	0	0	0	0
3	E	1	0	0	0	0
4	A	26	0	0	1	0
4	B	36	0	0	0	0
4	C	33	0	0	0	0
4	D	27	0	0	0	0
4	E	27	0	0	0	0
4	F	19	0	0	0	0
All	All	2969	2593	2592	11	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (11) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:E:2629:ARG:HD3	1:E:2666:ALA:HB1	1.34	1.07
1:A:2644:ALA:O	1:A:2648:VAL:HG22	1.92	0.68
1:F:2633:CYS:SG	1:F:2662:HIS:HB3	2.46	0.55
1:E:2629:ARG:CD	1:E:2666:ALA:HB1	2.24	0.52
1:E:2629:ARG:HD3	1:E:2666:ALA:CB	2.25	0.51
1:F:2659:THR:OG1	1:F:2661:ASP:HB2	2.10	0.51
1:A:2651:ALA:HA	1:A:2654:LEU:HD12	1.98	0.46
1:A:2635:VAL:HG22	4:A:2816:HOH:O	2.16	0.45
1:B:2648:VAL:HG21	1:F:2668:PRO:HG3	2.02	0.42
1:C:2627:LEU:O	1:C:2628:ARG:HB3	2.19	0.41
1:D:2651:ALA:HB3	1:D:2660:ARG:CA	2.51	0.41

There are no symmetry-related clashes.

## 5.3 Torsion angles (i)

### 5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	57/59 (97%)	56 (98%)	1 (2%)	0	100 100
1	B	57/59 (97%)	55 (96%)	2 (4%)	0	100 100
1	C	57/59 (97%)	55 (96%)	2 (4%)	0	100 100
1	D	57/59 (97%)	56 (98%)	1 (2%)	0	100 100
1	E	53/59 (90%)	52 (98%)	1 (2%)	0	100 100
1	F	53/59 (90%)	51 (96%)	2 (4%)	0	100 100
All	All	334/354 (94%)	325 (97%)	9 (3%)	0	100 100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	55/55 (100%)	55 (100%)	0	100 100
1	B	55/55 (100%)	55 (100%)	0	100 100
1	C	55/55 (100%)	55 (100%)	0	100 100
1	D	55/55 (100%)	55 (100%)	0	100 100
1	E	54/55 (98%)	54 (100%)	0	100 100
1	F	53/55 (96%)	53 (100%)	0	100 100
All	All	327/330 (99%)	327 (100%)	0	100 100

There are no protein residues with a non-rotameric sidechain to report.

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	D	2615	GLN

### 5.3.3 RNA [\(i\)](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [\(i\)](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [\(i\)](#)

There are no oligosaccharides in this entry.

### 5.6 Ligand geometry [\(i\)](#)

Of 14 ligands modelled in this entry, 14 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

### 5.7 Other polymers [\(i\)](#)

There are no such residues in this entry.

### 5.8 Polymer linkage issues [\(i\)](#)

There are no chain breaks in this entry.

## 6 Fit of model and data i

### 6.1 Protein, DNA and RNA chains i

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	59/59 (100%)	0.10	2 (3%) 48 50	15, 23, 44, 63	0
1	B	59/59 (100%)	-0.21	2 (3%) 48 50	12, 19, 47, 53	0
1	C	59/59 (100%)	0.03	3 (5%) 34 37	14, 23, 57, 67	0
1	D	59/59 (100%)	0.23	4 (6%) 25 26	14, 25, 45, 59	0
1	E	57/59 (96%)	0.35	3 (5%) 33 35	15, 29, 51, 65	0
1	F	57/59 (96%)	0.57	6 (10%) 13 13	19, 30, 49, 62	0
All	All	350/354 (98%)	0.18	20 (5%) 30 32	12, 25, 49, 67	0

All (20) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	2651	ALA	4.3
1	C	2650	ASP	3.8
1	E	2650	ASP	3.7
1	F	2657	PRO	3.6
1	D	2651	ALA	3.3
1	F	2635	VAL	3.1
1	F	2661	ASP	3.1
1	A	2652	ASP	2.9
1	E	2654	LEU	2.9
1	D	2650	ASP	2.7
1	C	2652	ASP	2.7
1	D	2648	VAL	2.6
1	B	2650	ASP	2.5
1	A	2648	VAL	2.4
1	F	2633	CYS	2.4
1	B	2653	ARG	2.3
1	F	2634	THR	2.3
1	E	2648	VAL	2.2
1	D	2654	LEU	2.1

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	F	2653	ARG	2.1

## 6.2 Non-standard residues in protein, DNA, RNA chains [\(i\)](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [\(i\)](#)


There are no oligosaccharides in this entry.


## 6.4 Ligands [\(i\)](#)


In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
2	ZN	F	2702	1/1	0.96	0.06	38,38,38,38	0
2	ZN	A	2702	1/1	0.99	0.01	15,15,15,15	0
2	ZN	B	2702	1/1	0.99	0.01	14,14,14,14	0
2	ZN	C	2702	1/1	0.99	0.02	16,16,16,16	0
2	ZN	D	2702	1/1	0.99	0.02	18,18,18,18	0
2	ZN	E	2701	1/1	0.99	0.03	26,26,26,26	0
2	ZN	E	2702	1/1	0.99	0.02	18,18,18,18	0
2	ZN	A	2701	1/1	0.99	0.03	16,16,16,16	0
3	NA	D	2703	1/1	0.99	0.10	7,7,7,7	0
3	NA	E	2703	1/1	0.99	0.09	2,2,2,2	0
2	ZN	F	2701	1/1	1.00	0.01	19,19,19,19	0
2	ZN	C	2701	1/1	1.00	0.01	17,17,17,17	0
2	ZN	B	2701	1/1	1.00	0.01	15,15,15,15	0
2	ZN	D	2701	1/1	1.00	0.02	21,21,21,21	0


The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.






































## 6.5 Other polymers [\(i\)](#)

There are no such residues in this entry.