



# Full wwPDB X-ray Structure Validation Report ⓘ

Feb 16, 2026 – 02:30 PM JST

PDB ID : 9LYY / pdb\_00009lyy  
Title : Crystal structure of glycerol kinase from Entamoeba histolytica complexed with ADP and G3P.  
Authors : Balogun, E.O.; Jeelani, G.; Hane, E.; Kondo, H.; Hasegawa, Y.; Kojima, C.; Chishima, T.; Harada, S.; Kishikawa, J.; Nozaki, T.; Shiba, T.  
Deposited on : 2025-02-21  
Resolution : 1.60 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 2.0  
EDS : 3.0  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)  
CCP4 : 9.0.010 (Gargrove)  
Density-Fitness : 1.0.12  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.48


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	164625	4274 (1.60-1.60)
Clashscore	180529	4682 (1.60-1.60)
Ramachandran outliers	177936	4583 (1.60-1.60)
Sidechain outliers	177891	4582 (1.60-1.60)
RSRZ outliers	164620	4272 (1.60-1.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	487	 2% (red), 87% (green), 11% (yellow), 0% (orange), 0% (grey)
1	B	487	 0% (red), 93% (green), 7% (yellow), 0% (orange), 0% (grey)

## 2 Entry composition [i](#)

There are 6 unique types of molecules in this entry. The entry contains 7967 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called glycerol kinase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	485	Total 3789	C 2416	N 625	O 726	S 22	0	2	0
1	B	487	Total 3823	C 2440	N 630	O 733	S 20	0	5	0

There are 4 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	-1	ALA	-	expression tag	UNP A0A5K1V6Z1
A	0	ALA	-	expression tag	UNP A0A5K1V6Z1
B	-1	ALA	-	expression tag	UNP A0A5K1V6Z1
B	0	ALA	-	expression tag	UNP A0A5K1V6Z1

- Molecule 2 is ADENOSINE-5'-DIPHOSPHATE (CCD ID: ADP) (formula: C<sub>10</sub>H<sub>15</sub>N<sub>5</sub>O<sub>10</sub>P<sub>2</sub>) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	N	O	P		
2	A	1	27	10	5	10	2	0	0
2	B	1	27	10	5	10	2	0	0

- Molecule 3 is SN-GLYCEROL-3-PHOSPHATE (CCD ID: G3P) (formula: C<sub>3</sub>H<sub>9</sub>O<sub>6</sub>P) (labeled as "Ligand of Interest" by depositor).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	O	P		
3	A	1	10	3	6	1	0	0
3	B	1	10	3	6	1	0	0

- Molecule 4 is MAGNESIUM ION (CCD ID: MG) (formula: Mg) (labeled as "Ligand of Interest" by depositor).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	Mg		
4	A	1	1	1	0	0
4	B	1	1	1	0	0

- Molecule 5 is TETRAETHYLENE GLYCOL (CCD ID: PG4) (formula: C<sub>8</sub>H<sub>18</sub>O<sub>5</sub>) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	B	1	Total	C	O	0	0
			13	8	5		


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	130	Total	O	0	0
			130	130		
6	B	136	Total	O	0	0
			136	136		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: glycerol kinase



- Molecule 1: glycerol kinase



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	57.35Å 79.95Å 205.04Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	19.99 – 1.60 19.99 – 1.60	Depositor EDS
% Data completeness (in resolution range)	97.9 (19.99-1.60) 98.0 (19.99-1.60)	Depositor EDS
$R_{merge}$	0.08	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.43 (at 1.60Å)	Xtrriage
Refinement program	REFMAC 5.8.0103	Depositor
R, $R_{free}$	0.186 , 0.206 0.194 , 0.213	Depositor DCC
$R_{free}$ test set	6103 reflections (4.85%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	24.9	Xtrriage
Anisotropy	0.119	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.36 , 28.2	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.97	EDS
Total number of atoms	7967	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	29.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.20% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: MG, PG4, ADP, G3P

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.70	5/3861 (0.1%)	0.84	3/5221 (0.1%)
1	B	0.65	3/3895 (0.1%)	0.81	2/5270 (0.0%)
All	All	0.67	8/7756 (0.1%)	0.83	5/10491 (0.0%)

All (8) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	77	ILE	C-O	-6.54	1.17	1.24
1	A	80	GLN	C-O	-6.50	1.16	1.23
1	B	163	ILE	C-O	-6.25	1.16	1.24
1	A	77	ILE	C-O	-6.10	1.18	1.24
1	A	163	ILE	C-O	-6.10	1.17	1.24
1	B	79	ASN	C-O	-5.70	1.17	1.23
1	A	159	LEU	C-O	-5.59	1.16	1.24
1	A	160	PHE	C-O	-5.12	1.17	1.23

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	448	ASP	N-CA-C	9.50	121.64	111.28
1	B	79	ASN	N-CA-C	6.97	119.46	108.79
1	A	79	ASN	N-CA-C	6.33	118.47	108.79
1	A	162	THR	N-CA-C	-5.39	102.92	110.35
1	B	162	THR	N-CA-C	-5.36	102.95	110.35

There are no chirality outliers.

There are no planarity outliers.

## 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3789	0	3791	53	0
1	B	3823	0	3832	15	0
2	A	27	0	12	0	0
2	B	27	0	12	0	0
3	A	10	0	7	0	0
3	B	10	0	7	0	0
4	A	1	0	0	0	0
4	B	1	0	0	0	0
5	B	13	0	18	0	0
6	A	130	0	0	0	0
6	B	136	0	0	0	0
All	All	7967	0	7679	66	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (66) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:443:LYS:HD3	1:A:444:GLU:N	1.63	1.12
1:A:443:LYS:CD	1:A:444:GLU:OE2	2.03	1.06
1:A:443:LYS:HD3	1:A:444:GLU:OE2	1.59	1.01
1:A:443:LYS:HD2	1:A:444:GLU:OE2	1.72	0.89
1:A:443:LYS:HD3	1:A:444:GLU:H	1.32	0.87
1:A:443:LYS:HD2	1:A:444:GLU:CD	1.98	0.87
1:A:443:LYS:CD	1:A:444:GLU:H	1.88	0.86
1:A:443:LYS:HZ3	1:A:444:GLU:HB3	1.40	0.84
1:B:45:GLN:HE22	1:B:80:GLN:HE22	1.28	0.82
1:A:447:LEU:HD12	1:A:451:ARG:HH22	1.44	0.81
1:A:447:LEU:HG	1:A:448:ASP:N	1.95	0.81
1:A:443:LYS:CD	1:A:444:GLU:CD	2.52	0.81
1:B:217:GLN:HE22	1:B:292:GLY:H	1.30	0.80
1:A:45:GLN:HE22	1:A:80:GLN:HE22	1.30	0.78
1:A:443:LYS:CE	1:A:444:GLU:H	1.97	0.77
1:A:443:LYS:NZ	1:A:444:GLU:HB3	2.01	0.76

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:443:LYS:CD	1:A:444:GLU:N	2.46	0.69
1:A:382:ALA:HB2	1:A:413:ILE:HD13	1.81	0.61
1:A:447:LEU:HD23	1:A:448:ASP:H	1.66	0.60
1:A:444:GLU:HG2	1:A:445:ASP:N	2.16	0.60
1:A:443:LYS:HE2	1:A:444:GLU:H	1.66	0.59
1:A:111:SER:HA	1:A:114:MET:HE3	1.84	0.58
1:A:323:GLU:HG3	1:A:327:ILE:HG13	1.88	0.55
1:A:444:GLU:CG	1:A:445:ASP:N	2.71	0.54
1:A:444:GLU:O	1:A:447:LEU:HD22	2.09	0.53
1:B:111:SER:HA	1:B:114:MET:HE3	1.89	0.52
1:A:413:ILE:HD12	1:A:414:CYS:N	2.25	0.51
1:A:437:LEU:HD11	1:A:443:LYS:HE2	1.92	0.51
1:B:220:ASP:O	1:B:237:THR:HA	2.11	0.50
1:B:62:LEU:HD21	1:B:72:ILE:HD11	1.93	0.50
1:B:350:ASP:OD2	1:B:485:HIS:HE1	1.94	0.50
1:A:443:LYS:HD3	1:A:443:LYS:C	2.27	0.50
1:A:444:GLU:O	1:A:447:LEU:HB3	2.12	0.50
1:A:447:LEU:CG	1:A:448:ASP:N	2.72	0.50
1:A:217:GLN:HE22	1:A:292:GLY:H	1.59	0.49
1:A:443:LYS:HD2	1:A:444:GLU:OE1	2.11	0.49
1:A:443:LYS:HD3	1:A:444:GLU:CD	2.28	0.48
1:B:110:THR:HG22	1:B:114:MET:HE2	1.96	0.48
1:B:217:GLN:NE2	1:B:292:GLY:H	2.05	0.47
1:B:267:VAL:HB	1:B:384:VAL:HG11	1.97	0.46
1:A:220:ASP:O	1:A:237:THR:HA	2.15	0.46
1:A:267:VAL:HB	1:A:384:VAL:HG11	1.97	0.46
1:A:443:LYS:HZ2	1:A:444:GLU:CD	2.24	0.45
1:B:5[B]:LEU:C	1:B:5[B]:LEU:HD13	2.42	0.45
1:B:332:ASN:C	1:B:332:ASN:HD22	2.24	0.45
1:A:217:GLN:NE2	1:A:292:GLY:H	2.14	0.45
1:A:444:GLU:O	1:A:447:LEU:N	2.30	0.45
1:A:441:LEU:HD12	1:A:441:LEU:HA	1.84	0.45
1:A:447:LEU:CD2	1:A:448:ASP:H	2.29	0.44
1:A:410:GLN:NE2	1:A:413:ILE:HD11	2.31	0.44
1:A:332:ASN:HD22	1:A:334:GLY:H	1.65	0.44
1:A:78:THR:OG1	1:A:242:ASP:HA	2.19	0.43
1:A:433:MET:HE1	1:A:451:ARG:NH1	2.33	0.43
1:A:361:THR:CG2	1:B:354:ARG:HD2	2.49	0.43
1:A:17:ILE:HD13	1:A:428:ALA:CB	2.49	0.43
1:B:181:ILE:HG21	1:B:289:GLU:HB2	2.01	0.42
1:A:258:LYS:HD2	1:A:258:LYS:C	2.45	0.42

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:248:PHE:CE2	1:A:433:MET:HE2	2.55	0.42
1:A:265:CYS:HB3	1:A:302:VAL:HB	2.01	0.41
1:A:444:GLU:C	1:A:446:GLU:N	2.76	0.41
1:B:5[A]:LEU:HD22	1:B:231:PHE:CE1	2.55	0.41
1:A:160:PHE:CG	1:A:161:GLY:N	2.88	0.41
1:A:110:THR:HG22	1:A:114:MET:HE2	2.02	0.40
1:A:27:HIS:HE1	1:A:29:GLU:OE2	2.05	0.40
1:A:90:ARG:HG3	1:A:157:VAL:CG2	2.52	0.40
1:A:361:THR:HG22	1:B:354:ARG:HD2	2.02	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	485/487 (100%)	475 (98%)	9 (2%)	1 (0%)	44	25
1	B	490/487 (101%)	484 (99%)	5 (1%)	1 (0%)	44	25
All	All	975/974 (100%)	959 (98%)	14 (1%)	2 (0%)	44	25

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	389	GLY
1	B	389	GLY

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	414/412 (100%)	395 (95%)	19 (5%)	23	6
1	B	417/412 (101%)	405 (97%)	12 (3%)	37	15
All	All	831/824 (101%)	800 (96%)	31 (4%)	28	9

All (31) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	26	ILE
1	A	61	CYS
1	A	120	GLU
1	A	162	THR
1	A	180	ASP
1	A	327	ILE
1	A	332	ASN
1	A	362	SER
1	A	386	GLU
1	A	390	LYS
1	A	422	LYS
1	A	437	LEU
1	A	441	LEU
1	A	443	LYS
1	A	444	GLU
1	A	454	GLU
1	A	458	LYS
1	A	463	GLU
1	A	478	GLU
1	B	24	GLN
1	B	26	ILE
1	B	157	VAL
1	B	162	THR
1	B	180	ASP
1	B	309	LEU
1	B	332	ASN
1	B	390	LYS
1	B	422	LYS
1	B	458	LYS
1	B	472	VAL
1	B	475	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (27)

such sidechains are listed below:

Mol	Chain	Res	Type
1	A	27	HIS
1	A	45	GLN
1	A	123	GLN
1	A	170	ASN
1	A	206	ASN
1	A	217	GLN
1	A	232	ASN
1	A	322	ASN
1	A	332	ASN
1	A	408	GLN
1	A	440	HIS
1	A	455	HIS
1	A	468	GLN
1	B	27	HIS
1	B	45	GLN
1	B	170	ASN
1	B	175	GLN
1	B	206	ASN
1	B	217	GLN
1	B	232	ASN
1	B	291	ASN
1	B	322	ASN
1	B	332	ASN
1	B	342	GLN
1	B	455	HIS
1	B	468	GLN
1	B	485	HIS

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

## 5.6 Ligand geometry

Of 7 ligands modelled in this entry, 2 are monoatomic - leaving 5 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
3	G3P	B	502	4	9,9,9	0.43	0	11,12,12	0.96	0
5	PG4	B	504	-	12,12,12	0.44	0	11,11,11	0.40	0
2	ADP	B	501	-	24,29,29	0.94	1 (4%)	29,45,45	1.50	5 (17%)
3	G3P	A	502	4	9,9,9	0.53	0	11,12,12	0.82	0
2	ADP	A	501	-	24,29,29	1.04	2 (8%)	29,45,45	1.64	6 (20%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	G3P	B	502	4	-	0/8/8/8	-
5	PG4	B	504	-	-	0/10/10/10	-
2	ADP	B	501	-	-	4/12/32/32	0/3/3/3
3	G3P	A	502	4	-	0/8/8/8	-
2	ADP	A	501	-	-	2/12/32/32	0/3/3/3

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	501	ADP	C2-N3	2.41	1.36	1.32
2	A	501	ADP	C5-C4	2.29	1.47	1.40
2	A	501	ADP	O4'-C1'	2.15	1.44	1.41

All (11) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	501	ADP	N3-C2-N1	-4.42	121.77	128.68
2	B	501	ADP	N3-C2-N1	-4.22	122.08	128.68

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	501	ADP	C3'-C2'-C1'	3.65	106.48	100.98
2	B	501	ADP	C3'-C2'-C1'	3.32	105.98	100.98
2	A	501	ADP	C2-N1-C6	2.70	123.37	118.75
2	B	501	ADP	C1'-N9-C4	-2.67	121.95	126.64
2	A	501	ADP	O3B-PB-O2B	2.60	117.58	107.64
2	B	501	ADP	C4-C5-N7	-2.42	106.88	109.40
2	A	501	ADP	C1'-N9-C4	-2.29	122.61	126.64
2	A	501	ADP	O3B-PB-O3A	-2.07	97.69	104.64
2	B	501	ADP	PA-O3A-PB	2.06	139.90	132.83

There are no chirality outliers.


All (6) torsion outliers are listed below:



Mol	Chain	Res	Type	Atoms
2	A	501	ADP	PA-O3A-PB-O3B
2	A	501	ADP	PB-O3A-PA-O5'
2	B	501	ADP	PB-O3A-PA-O5'
2	B	501	ADP	PA-O3A-PB-O1B
2	B	501	ADP	PA-O3A-PB-O2B
2	B	501	ADP	PA-O3A-PB-O3B


There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.







## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	485/487 (99%)	-0.01	11 (2%) 61 63	13, 26, 47, 71	2 (0%)
1	B	487/487 (100%)	-0.04	5 (1%) 79 82	11, 28, 45, 70	5 (1%)
All	All	972/974 (99%)	-0.02	16 (1%) 70 73	11, 27, 46, 71	7 (0%)

All (16) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	447	LEU	5.8
1	A	450	MET	4.3
1	A	449	SER	3.9
1	A	67	ILE	3.0
1	A	463	GLU	2.7
1	B	391	ILE	2.7
1	A	448	ASP	2.7
1	B	389	GLY	2.4
1	A	445	ASP	2.4
1	B	394	ILE	2.3
1	A	68	ASN	2.2
1	A	1	MET	2.2
1	B	26	ILE	2.1
1	A	444	GLU	2.1
1	A	389	GLY	2.1
1	B	147	GLU	2.0

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

### 6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
5	PG4	B	504	13/13	0.87	0.11	39,41,46,47	0
2	ADP	B	501	27/27	0.91	0.10	28,34,48,52	0
2	ADP	A	501	27/27	0.94	0.08	26,28,36,39	0
3	G3P	B	502	10/10	0.97	0.06	20,22,25,25	0
4	MG	A	503	1/1	0.97	0.06	25,25,25,25	0
3	G3P	A	502	10/10	0.97	0.06	17,19,24,24	0
4	MG	B	503	1/1	0.99	0.07	26,26,26,26	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

**Electron density around PG4 B 504:**




$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)








**Electron density around G3P B 502:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)






**Electron density around MG A 503:**


$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)



**Electron density around G3P A 502:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)





## 6.5 Other polymers ⓘ

There are no such residues in this entry.