



Full wwPDB X-ray Structure Validation Report i

Jan 6, 2026 – 01:50 PM EST

PDB ID : 9MNT / pdb_00009mnt
Title : Structure of Hepatitis C virus envelope glycoprotein E2 core from genotype 6a bound to broadly neutralizing antibody RM1-73
Authors : Nguyen, T.K.Y.; Stanfield, R.L.; Wilson, I.A.
Deposited on : 2024-12-23
Resolution : 2.55 Å (reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>
with specific help available everywhere you see the i symbol.

The types of validation reports are described at
<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references](#) i) were used in the production of this report:

MolProbity : 4.5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
buster-report : 1.1.7 (2018)
Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.47


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION


The reported resolution of this entry is 2.55 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	164625	1004 (2.54-2.54)
Clashscore	180529	1055 (2.54-2.54)
Ramachandran outliers	177936	1048 (2.54-2.54)
Sidechain outliers	177891	1048 (2.54-2.54)
RSRZ outliers	164620	1004 (2.54-2.54)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.



The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
5	NAG	B	1	-	-	X	-

2 Entry composition (i)

There are 8 unique types of molecules in this entry. The entry contains 4734 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called RM1-73 Fab light chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	L	212	Total	C	N	O	S	0	0	0

1629 1015 272 336 6

- Molecule 2 is a protein called RM1-73 Fab heavy chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	H	230	Total	C	N	O	S	0	0	0

1671 1049 282 334 6

- Molecule 3 is a protein called Hepatitis C virus envelope glycoprotein E2 core from genotype 6a.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	E	169	Total	C	N	O	S	0	0	0

1299 826 222 238 13

- Molecule 4 is an oligosaccharide called 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
4	A	2	Total	C	N	O		0	0	0


28 16 2 10

- Molecule 5 is an oligosaccharide called beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.




Mol	Chain	Residues	Atoms	ZeroOcc	AltConf	Trace
5	B	3	Total C N O 39 22 2 15	0	0	0

- Molecule 6 is SULFATE ION (CCD ID: SO4) (formula: O₄S) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	L	1	Total O S 5 4 1	0	0

- Molecule 7 is 2-acetamido-2-deoxy-beta-D-glucopyranose (CCD ID: NAG) (formula: C₈H₁₅NO₆) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
7	E	1	Total C N O 14 8 1 5	0	0
7	E	1	Total C N O 14 8 1 5	0	0
7	E	1	Total C N O 14 8 1 5	0	0


- Molecule 8 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
8	L	9	Total O 9 9	0	0
8	H	11	Total O 11 11	0	0
8	E	1	Total O 1 1	0	0


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: RM1-73 Fab light chain



- Molecule 2: RM1-73 Fab heavy chain



- Molecule 3: Hepatitis C virus envelope glycoprotein E2 core from genotype 6a



- Molecule 4: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose



- Molecule 5: beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose



MAG1
MAG2
BMA3

4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	60.63 \AA 73.22 \AA 208.91 \AA 90.00° 90.00° 90.00°	Depositor
Resolution (\AA)	46.70 – 2.55 46.70 – 2.55	Depositor EDS
% Data completeness (in resolution range)	99.0 (46.70-2.55) 99.6 (46.70-2.55)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle^{\text{1}}$	1.51 (at 2.54 \AA)	Xtriage
Refinement program	PHENIX 1.19.2_4158, PHENIX 1.19.2_4158	Depositor
R, R_{free}	0.231, 0.252 0.230, 0.250	Depositor DCC
R_{free} test set	1493 reflections (4.76%)	wwPDB-VP
Wilson B-factor (\AA^2)	51.3	Xtriage
Anisotropy	0.313	Xtriage
Bulk solvent k_{sol} (e/ \AA^3), B_{sol} (\AA^2)	0.33, 34.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.49, \langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	4734	wwPDB-VP
Average B, all atoms (\AA^2)	57.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.96% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [\(i\)](#)

5.1 Standard geometry [\(i\)](#)

Bond lengths and bond angles in the following residue types are not validated in this section: SO4, BMA, NAG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	L	0.09	0/1662	0.28	0/2255
2	H	0.18	0/1705	0.34	0/2322
3	E	0.25	0/1344	0.47	1/1840 (0.1%)
All	All	0.18	0/4711	0.36	1/6417 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
3	E	423	ASN	CA-CB-CG	-7.63	104.97	112.60

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [\(i\)](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	L	1629	0	1577	7	0
2	H	1671	0	1658	11	1
3	E	1299	0	1211	31	0
4	A	28	0	25	0	0
5	B	39	0	34	10	0
6	L	5	0	0	0	0
7	E	42	0	39	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
8	E	1	0	0	0	0
8	H	11	0	0	0	0
8	L	9	0	0	0	0
All	All	4734	0	4544	48	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (48) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:E:527:TYR:HB3	5:B:1:NAG:O7	1.55	1.06
3:E:423:ASN:ND2	5:B:1:NAG:C1	2.31	0.93
3:E:527:TYR:CB	5:B:1:NAG:O7	2.17	0.93
2:H:94:ARG:HG2	2:H:102:SER:HB2	1.70	0.72
3:E:423:ASN:HD21	5:B:1:NAG:C1	2.04	0.69
3:E:601:PRO:HG2	3:E:608:LEU:HD11	1.74	0.68
3:E:527:TYR:CG	5:B:1:NAG:H81	2.30	0.65
3:E:421:HIS:ND1	3:E:423:ASN:HB2	2.16	0.61
1:L:37:GLN:HB2	1:L:47:LEU:HD11	1.82	0.61
2:H:54:VAL:HG11	3:E:438:ILE:HG21	1.83	0.59
2:H:30:SER:HB2	2:H:53:LEU:HB2	1.85	0.59
3:E:603:ILE:HG23	3:E:604:THR:HG23	1.86	0.58
2:H:123:PRO:HD3	2:H:209:LYS:HE3	1.85	0.58
3:E:527:TYR:HB2	5:B:1:NAG:O7	2.02	0.56
3:E:608:LEU:HD22	3:E:621:THR:HG21	1.88	0.56
1:L:166:GLN:HG3	1:L:173:TYR:CZ	2.41	0.56
3:E:555:MET:HG2	3:E:617:HIS:HD2	1.72	0.54
3:E:422:ILE:O	3:E:425:THR:HG22	2.10	0.51
3:E:509:PHE:CZ	3:E:601:PRO:HB3	2.47	0.50
2:H:6:GLN:HB3	2:H:107:VAL:HG23	1.95	0.48
3:E:527:TYR:HB3	5:B:1:NAG:C7	2.40	0.47
3:E:555:MET:HG2	3:E:617:HIS:CD2	2.48	0.47
3:E:632:PHE:CZ	3:E:637:GLU:HB2	2.48	0.47
3:E:423:ASN:ND2	3:E:529:TRP:CE2	2.82	0.47
3:E:527:TYR:CB	5:B:1:NAG:C7	2.92	0.47
3:E:527:TYR:O	5:B:1:NAG:O7	2.32	0.47
3:E:421:HIS:CE1	3:E:423:ASN:HB2	2.50	0.47
2:H:66:ARG:NH2	2:H:86:ASP:OD2	2.48	0.46
2:H:89:VAL:HG22	2:H:108:VAL:HG22	1.96	0.46
3:E:495:ASP:HA	3:E:540:GLU:HB2	1.97	0.46

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:E:423:ASN:ND2	3:E:529:TRP:CZ2	2.84	0.46
2:H:2:VAL:HG13	2:H:27:PHE:CD1	2.51	0.45
1:L:28:ASN:OD1	1:L:68:GLY:HA2	2.18	0.44
3:E:440:SER:HB2	3:E:616:TRP:CD1	2.51	0.44
1:L:113:PRO:HB3	1:L:139:PHE:HB3	2.00	0.44
1:L:89:GLN:HB2	1:L:98:PHE:CD2	2.54	0.43
2:H:8:GLY:HA2	2:H:201:LYS:HD3	2.01	0.43
3:E:448:ASP:O	3:E:451:GLY:N	2.51	0.43
3:E:527:TYR:CB	5:B:1:NAG:H81	2.48	0.43
3:E:607:CYS:HA	3:E:644:CYS:HA	2.01	0.42
3:E:546:THR:HG21	3:E:634:GLY:HA3	2.02	0.42
1:L:4:MET:HE1	1:L:25:ALA:HB2	2.02	0.42
3:E:433:LEU:HD11	3:E:558:THR:O	2.20	0.42
3:E:509:PHE:CE2	3:E:601:PRO:HB3	2.54	0.42
2:H:119:PRO:HB3	2:H:145:TYR:HB3	2.02	0.41
1:L:125:LEU:O	1:L:183:LYS:HD2	2.20	0.41
3:E:567:PRO:HA	3:E:568:PRO:HD3	1.98	0.41
2:H:87:THR:HA	2:H:109:VAL:O	2.22	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:H:9:ALA:O	2:H:129:LYS:NZ[3_454]	2.15	0.05

5.3 Torsion angles [\(i\)](#)

5.3.1 Protein backbone [\(i\)](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	L	210/212 (99%)	203 (97%)	7 (3%)	0	100 100
2	H	228/235 (97%)	226 (99%)	2 (1%)	0	100 100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
3	E	165/189 (87%)	154 (93%)	9 (6%)	2 (1%)	11 14
All	All	603/636 (95%)	583 (97%)	18 (3%)	2 (0%)	37 46

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
3	E	445	LYS
3	E	510	THR

5.3.2 Protein sidechains [\(i\)](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	L	187/187 (100%)	186 (100%)	1 (0%)	86 93
2	H	186/191 (97%)	184 (99%)	2 (1%)	70 82
3	E	144/159 (91%)	140 (97%)	4 (3%)	38 55
All	All	517/537 (96%)	510 (99%)	7 (1%)	62 77

All (7) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	L	33	LEU
2	H	11	VAL
2	H	83	ARG
3	E	433	LEU
3	E	434	GLN
3	E	536	VAL
3	E	604	THR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
2	H	64	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
3	E	421	HIS
3	E	423	ASN

5.3.3 RNA [\(i\)](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [\(i\)](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [\(i\)](#)

5 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	NAG	A	1	3,4	14,14,15	0.71	0	17,19,21	1.21	2 (11%)
4	NAG	A	2	4	14,14,15	0.69	0	17,19,21	0.91	0
5	NAG	B	1	5	14,14,15	0.38	0	17,19,21	0.66	0
5	NAG	B	2	5	14,14,15	0.37	0	17,19,21	0.65	0
5	BMA	B	3	5	11,11,12	0.25	0	15,15,17	0.56	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	NAG	A	1	3,4	-	1/6/23/26	0/1/1/1
4	NAG	A	2	4	-	2/6/23/26	0/1/1/1
5	NAG	B	1	5	-	4/6/23/26	0/1/1/1

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	NAG	B	2	5	-	0/6/23/26	0/1/1/1
5	BMA	B	3	5	-	0/2/19/22	0/1/1/1

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
4	A	1	NAG	C1-O5-C5	3.14	116.39	112.19
4	A	1	NAG	O4-C4-C3	-2.01	105.63	110.38

There are no chirality outliers.

All (7) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
5	B	1	NAG	C1-C2-N2-C7
4	A	2	NAG	C8-C7-N2-C2
4	A	2	NAG	O7-C7-N2-C2
4	A	1	NAG	O5-C5-C6-O6
5	B	1	NAG	C3-C2-N2-C7
5	B	1	NAG	C8-C7-N2-C2
5	B	1	NAG	O7-C7-N2-C2


There are no ring outliers.

1 monomer is involved in 10 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	B	1	NAG	10	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.





5.6 Ligand geometry (i)

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
6	SO4	L	301	-	4,4,4	0.66	0	6,6,6	0.12	0
7	NAG	E	703	3	14,14,15	0.70	0	17,19,21	0.85	0
7	NAG	E	701	3	14,14,15	0.69	0	17,19,21	0.88	0
7	NAG	E	702	3	14,14,15	0.68	0	17,19,21	0.87	1 (5%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
7	NAG	E	703	3	-	1/6/23/26	0/1/1/1
7	NAG	E	701	3	-	2/6/23/26	0/1/1/1
7	NAG	E	702	3	-	2/6/23/26	0/1/1/1

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
7	E	702	NAG	C2-N2-C7	2.31	125.99	122.90

There are no chirality outliers.


All (5) torsion outliers are listed below:



Mol	Chain	Res	Type	Atoms
7	E	701	NAG	C8-C7-N2-C2
7	E	701	NAG	O7-C7-N2-C2
7	E	702	NAG	C8-C7-N2-C2
7	E	702	NAG	O7-C7-N2-C2
7	E	703	NAG	O5-C5-C6-O6


There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.







5.7 Other polymers [\(i\)](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [\(i\)](#)

There are no chain breaks in this entry.

6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å ²)	Q<0.9
1	L	212/212 (100%)	0.22	2 (0%)	81	83	31, 52, 70, 95
2	H	230/235 (97%)	0.25	6 (2%)	57	61	31, 49, 70, 83
3	E	169/189 (89%)	1.08	27 (15%)	6	7	47, 67, 97, 177
All	All	611/636 (96%)	0.47	35 (5%)	30	34	31, 55, 83, 177

All (35) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
3	E	453	PRO	7.1
3	E	450	SER	6.0
1	L	212	CYS	5.4
2	H	7	SER	4.8
3	E	606	ARG	4.6
3	E	447	VAL	4.3
3	E	449	SER	4.2
3	E	423	ASN	4.0
3	E	448	ASP	3.9
3	E	595	THR	3.9
3	E	433	LEU	3.6
3	E	446	ASN	3.5
3	E	451	GLY	3.5
3	E	452	CYS	3.4
2	H	8	GLY	3.2
3	E	491	PRO	3.2
3	E	596	ASP	3.2
3	E	545	PRO	3.1
2	H	214	LYS	2.9
2	H	2	VAL	2.7
3	E	498	SER	2.5
3	E	417	ASN	2.5
3	E	432	SER	2.4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
3	E	495	ASP	2.4
3	E	522	LEU	2.4
1	L	190	LYS	2.4
3	E	608	LEU	2.3
3	E	492	ARG	2.3
3	E	422	ILE	2.3
3	E	540	GLU	2.3
2	H	127	SER	2.2
3	E	645	ASN	2.2
2	H	23	LYS	2.1
3	E	425	THR	2.0
3	E	444	ALA	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [\(i\)](#)


There are no non-standard protein/DNA/RNA residues in this entry.


6.3 Carbohydrates [\(i\)](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
4	NAG	A	1	14/15	-	-	74,85,99,107	0
4	NAG	A	2	14/15	-	-	98,108,111,114	0
5	NAG	B	1	14/15	-	-	30,30,30,30	0
5	NAG	B	2	14/15	-	-	30,30,30,30	0
5	BMA	B	3	11/12	-	-	96,96,96,96	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.







6.4 Ligands (i)


In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
7	NAG	E	701	14/15	0.24	0.19	105,117,123,124	0
7	NAG	E	702	14/15	0.44	0.21	95,113,128,129	0
7	NAG	E	703	14/15	0.60	0.20	116,126,131,132	0
6	SO4	L	301	5/5	0.64	0.12	97,114,123,176	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.







6.5 Other polymers [\(i\)](#)

There are no such residues in this entry.