# Full wwPDB X-ray Structure Validation Report (i)

Jan 6, 2026 – 01:50 PM EST

| | | |
|---|---|---|
| PDB ID | : | 9MSC / pdb_00009msc |
| Title | : | Structure of Hepatitis C Virus Envelope Glycoprotein HCV-1 E2ecto from genotype 1a bound to neutralizing antibody RM5-16 |
| Authors | : | Nguyen, T.K.Y.; Stanfield, R.L.; Wilson, I.A. |
| Deposited on | : | 2025-01-09 |
| Resolution | : | 3.39 Å(reported) |

The following versions of software and data (see references (i)) were used in the production of this report:

| | | |
|---|---|---|
| MolProbity | : | 4-5-2 with Phenix2.0 |
| Mogul | : | 2022.3.0, CSD as543be (2022) |
| Xtriage (Phenix) | : | 2.0 |
| EDS | : | 3.0 |
| buster-report | : | 1.1.7 (2018) |
| Percentile statistics | : | 20231227.v01 (using entries in the PDB archive December 27th 2023) |
| CCP4 | : | 9.0.010 (Gargrove) |
| Density-Fitness | : | 1.0.12 |
| Ideal geometry (proteins) | : | Engh & Huber (2001) |
| Ideal geometry (DNA, RNA) | : | Parkinson et al. (1996) |
| Validation Pipeline (wwPDB-VP) | : | 2.47 |

# 1  Overall quality at a glance  ⓘ

The following experimental techniques were used to determine the structure:
*X-RAY DIFFRACTION*

The reported resolution of this entry is 3.39 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



| Metric | Whole archive (#Entries) | Similar resolution (#Entries, resolution range(Å)) |
|---|---|---|
| $R_{free}$ | 164625 | 1140 (3.46-3.34) |
| Clashscore | 180529 | 1172 (3.46-3.34) |
| Ramachandran outliers | 177936 | 1172 (3.46-3.34) |
| Sidechain outliers | 177891 | 1172 (3.46-3.34) |
| RSRZ outliers | 164620 | 1140 (3.46-3.34) |

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

| Mol | Chain | Length | Quality of chain |
|---|---|---|---|
| 1 | L | 216 | 86%  10%  • |
| 2 | H | 224 | 86%  12%  •• |
| 3 | B | 217 | 4%  89%  11% |
| 4 | A | 223 | 78%  19%  •• |
| 5 | E | 262 | 3%  61%  24%  •  12% |

*Continued on next page...*

*Continued from previous page...*

| Mol | Chain | Length | Quality of chain |
|-----|-------|--------|------------------|
| 6 | C | 2 | 100% |
| 6 | D | 2 | 50% 50% |
| 7 | F | 3 | 100% |

## 2 Entry composition ⓘ

There are 9 unique types of molecules in this entry. The entry contains 8425 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called RM5-16 fab LC.

| Mol | Chain | Residues | Atoms | | | | | ZeroOcc | AltConf | Trace |
|-----|-------|----------|-------|---|---|---|---|---------|---------|-------|
| 1 | L | 208 | Total | C | N | O | S | 0 | 0 | 0 |
| | | | 1561 | 976 | 261 | 320 | 4 | | | |

- Molecule 2 is a protein called RM5-16 fan HC.

| Mol | Chain | Residues | Atoms | | | | | ZeroOcc | AltConf | Trace |
|-----|-------|----------|-------|---|---|---|---|---------|---------|-------|
| 2 | H | 222 | Total | C | N | O | S | 0 | 0 | 0 |
| | | | 1657 | 1045 | 278 | 328 | 6 | | | |

- Molecule 3 is a protein called HEPC46 fab LC.

| Mol | Chain | Residues | Atoms | | | | | ZeroOcc | AltConf | Trace |
|-----|-------|----------|-------|---|---|---|---|---------|---------|-------|
| 3 | B | 217 | Total | C | N | O | S | 0 | 0 | 0 |
| | | | 1632 | 1015 | 274 | 338 | 5 | | | |

- Molecule 4 is a protein called HEPC46 fab HC.

| Mol | Chain | Residues | Atoms | | | | | ZeroOcc | AltConf | Trace |
|-----|-------|----------|-------|---|---|---|---|---------|---------|-------|
| 4 | A | 220 | Total | C | N | O | S | 0 | 0 | 0 |
| | | | 1642 | 1032 | 279 | 324 | 7 | | | |

- Molecule 5 is a protein called Envelope glycoprotein E2.

| Mol | Chain | Residues | Atoms | | | | | ZeroOcc | AltConf | Trace |
|-----|-------|----------|-------|---|---|---|---|---------|---------|-------|
| 5 | E | 230 | Total | C | N | O | S | 0 | 0 | 0 |
| | | | 1785 | 1132 | 312 | 323 | 18 | | | |

- Molecule 6 is an oligosaccharide called 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.

| Mol | Chain | Residues | Atoms | | | | ZeroOcc | AltConf | Trace |
|-----|-------|----------|-------|---|---|---|---------|---------|-------|
| 6 | C | 2 | Total 28 | C 16 | N 2 | O 10 | 0 | 0 | 0 |
| 6 | D | 2 | Total 28 | C 16 | N 2 | O 10 | 0 | 0 | 0 |

- Molecule 7 is an oligosaccharide called beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.



| Mol | Chain | Residues | Atoms | | | | ZeroOcc | AltConf | Trace |
|-----|-------|----------|-------|---|---|---|---------|---------|-------|
| 7 | F | 3 | Total 39 | C 22 | N 2 | O 15 | 0 | 0 | 0 |

- Molecule 8 is 2-acetamido-2-deoxy-beta-D-glucopyranose (CCD ID: NAG) (formula: $C_8H_{15}NO_6$) (labeled as "Ligand of Interest" by depositor).



| Mol | Chain | Residues | Atoms | | | | ZeroOcc | AltConf |
|-----|-------|----------|-------|---|---|---|---------|---------|
| 8 | E | 1 | Total 14 | C 8 | N 1 | O 5 | 0 | 0 |

*Continued on next page...*

*Continued from previous page...*

| Mol | Chain | Residues | Atoms | | | | ZeroOcc | AltConf |
| --- | --- | --- | --- | --- | --- | --- | --- | --- |
| 8 | E | 1 | Total<br>14 | C<br>8 | N<br>1 | O<br>5 | 0 | 0 |
| 8 | E | 1 | Total<br>14 | C<br>8 | N<br>1 | O<br>5 | 0 | 0 |

- Molecule 9 is alpha-D-mannopyranose (CCD ID: MAN) (formula: $C_6H_{12}O_6$) (labeled as "Ligand of Interest" by depositor).



| Mol | Chain | Residues | Atoms | | | ZeroOcc | AltConf |
| --- | --- | --- | --- | --- | --- | --- | --- |
| 9 | E | 1 | Total<br>11 | C<br>6 | O<br>5 | 0 | 0 |

# 3 Residue-property plots ⓘ

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: RM5-16 fab LC

Chain L:

• Molecule 2: RM5-16 fan HC

Chain H:

• Molecule 3: HEPC46 fab LC

Chain B:

• Molecule 4: HEPC46 fab HC

Chain A:

• Molecule 5: Envelope glycoprotein E2

Chain E:

GLU THR HIS VAL THR GLY GLY SER ALA GLY HIS THR VAL SER GLY PHE VAL SER LEU LEU ALA PRO GLY ALA LYS GLN ASN VAL GLN LEU ILE ASN T416 S419 W420 L427 N428 L441 F442 N448 S449 S450 G451 C452 P453 A457 D464 G470 P471 I472 S473 N476

G479 P491 K492 I496 V497 P498 A499 V502 P505 F509 T510 P511 S512 P513 V514 D520 R521 S522 G523 A524 W529 F537 V538 R543 P544 N548 W549 F550 G551 C552 T553 W554 M555 N556 S557 V563 C564 G565 A566 P567 P568 C569 V570 L579 T583 D584

S595 G598 S599 G600 P601 W602 I603 T604 P605 R606 C607 L608 V609 W616 H617 C620 M631 Y632 V633 V636 R639 L640 N645

• Molecule 6: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain C:

100%

NAG1 NAG2

• Molecule 6: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain D:

50% 50%

NAG1 NAG2

• Molecule 7: beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain F:

100%

NAG1 NAG2 BMA3

# 4 Data and refinement statistics ⓘ

| Property | Value | Source |
|---|---|---|
| Space group | P 32 2 1 | Depositor |
| Cell constants<br>a, b, c, $\alpha$, $\beta$, $\gamma$ | 72.71Å    72.71Å    564.78Å<br>90.00°    90.00°    120.00° | Depositor |
| Resolution (Å) | 49.64   –   3.39<br>49.64   –   3.40 | Depositor<br>EDS |
| % Data completeness<br>(in resolution range) | 95.4 (49.64-3.39)<br>95.7 (49.64-3.40) | Depositor<br>EDS |
| $R_{merge}$ | (Not available) | Depositor |
| $R_{sym}$ | (Not available) | Depositor |
| $< I/\sigma(I) >$ [1] | 1.11 (at 3.40Å) | Xtriage |
| Refinement program | PHENIX 1.21rc1_5127 | Depositor |
| R, $R_{free}$ | 0.278   ,   0.317<br>0.282   ,   0.320 | Depositor<br>DCC |
| $R_{free}$ test set | 1244 reflections (4.88%) | wwPDB-VP |
| Wilson B-factor (Å$^2$) | 111.3 | Xtriage |
| Anisotropy | 0.175 | Xtriage |
| Bulk solvent $k_{sol}$(e/Å$^3$), $B_{sol}$(Å$^2$) | 0.32 , 115.4 | EDS |
| L-test for twinning[2] | $< |L| > = 0.39, < L^2 > = 0.22$ | Xtriage |
| Estimated twinning fraction | 0.118 for -h,-k,l | Xtriage |
| Reported twinning fraction | 0.110 for -h,-k,l | Depositor |
| Outliers | 0 of 24365 reflections | Xtriage |
| $F_o$,$F_c$ correlation | 0.89 | EDS |
| Total number of atoms | 8425 | wwPDB-VP |
| Average B, all atoms (Å$^2$) | 162.0 | wwPDB-VP |

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.20% of the height of the origin peak. No significant pseudotranslation is detected.*

---

[1]Intensities estimated from amplitudes.
[2]Theoretical values of $< |L| >$, $< L^2 >$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

# 5 Model quality ⓘ

## 5.1 Standard geometry ⓘ

Bond lengths and bond angles in the following residue types are not validated in this section: BMA, MAN, PCA, NAG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

| Mol | Chain | Bond lengths | | Bond angles | |
|-----|-------|------|---------|------|---------|
| | | RMSZ | #$|Z| > 5$ | RMSZ | #$|Z| > 5$ |
| 1 | L | 0.09 | 0/1597 | 0.26 | 0/2179 |
| 2 | H | 0.10 | 0/1693 | 0.26 | 0/2313 |
| 3 | B | 0.11 | 0/1670 | 0.28 | 0/2275 |
| 4 | A | 0.15 | 0/1680 | 0.44 | 0/2290 |
| 5 | E | 0.14 | 0/1854 | 0.47 | 0/2543 |
| All | All | 0.12 | 0/8494 | 0.36 | 0/11600 |

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

## 5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

| Mol | Chain | Non-H | H(model) | H(added) | Clashes | Symm-Clashes |
|-----|-------|-------|----------|----------|---------|--------------|
| 1 | L | 1561 | 0 | 1511 | 12 | 0 |
| 2 | H | 1657 | 0 | 1625 | 16 | 0 |
| 3 | B | 1632 | 0 | 1553 | 16 | 0 |
| 4 | A | 1642 | 0 | 1618 | 36 | 0 |
| 5 | E | 1785 | 0 | 1640 | 41 | 0 |
| 6 | C | 28 | 0 | 25 | 1 | 0 |
| 6 | D | 28 | 0 | 25 | 0 | 0 |
| 7 | F | 39 | 0 | 34 | 0 | 0 |

*Continued on next page...*

*Continued from previous page...*

| Mol | Chain | Non-H | H(model) | H(added) | Clashes | Symm-Clashes |
|-----|-------|-------|----------|----------|---------|--------------|
| 8 | E | 42 | 0 | 39 | 1 | 0 |
| 9 | E | 11 | 0 | 10 | 0 | 0 |
| All | All | 8425 | 0 | 8080 | 114 | 0 |

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

All (114) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

| Atom-1 | Atom-2 | Interatomic distance (Å) | Clash overlap (Å) |
|--------|--------|--------------------------|-------------------|
| 5:E:604:THR:HG22 | 5:E:606:ARG:H | 1.43 | 0.83 |
| 4:A:87:THR:HG23 | 4:A:110:THR:HA | 1.64 | 0.79 |
| 5:E:551:GLY:HA3 | 5:E:600:GLY:HA2 | 1.68 | 0.75 |
| 2:H:11:LEU:HB2 | 2:H:147:PRO:HG3 | 1.70 | 0.73 |
| 5:E:553:THR:HG22 | 5:E:563:VAL:HG22 | 1.73 | 0.70 |
| 3:B:113:PRO:HB3 | 3:B:139:PHE:HB3 | 1.75 | 0.69 |
| 4:A:36:TRP:HB3 | 4:A:48:MET:HE3 | 1.75 | 0.68 |
| 1:L:83:GLU:HG3 | 1:L:106:VAL:HG23 | 1.77 | 0.66 |
| 3:B:95:LEU:O | 4:A:61:GLN:NE2 | 2.29 | 0.65 |
| 1:L:39:LEU:HD23 | 1:L:84:ALA:HB2 | 1.80 | 0.64 |
| 5:E:448:ASN:O | 5:E:450:SER:N | 2.31 | 0.64 |
| 5:E:583:THR:OG1 | 5:E:584:ASP:N | 2.31 | 0.63 |
| 5:E:464:ASP:HB3 | 5:E:579:LEU:HD13 | 1.81 | 0.62 |
| 4:A:28:ILE:HD11 | 8:E:701:NAG:H83 | 1.81 | 0.62 |
| 5:E:452:CYS:H | 5:E:453:PRO:HD2 | 1.64 | 0.61 |
| 3:B:120:PRO:HD3 | 3:B:132:VAL:HG22 | 1.82 | 0.60 |
| 4:A:119:PRO:HB3 | 4:A:145:TYR:HB3 | 1.83 | 0.60 |
| 4:A:168:ALA:HB2 | 4:A:178:LEU:HD23 | 1.83 | 0.60 |
| 1:L:132:LEU:HD12 | 1:L:178:LEU:HD23 | 1.84 | 0.59 |
| 5:E:476:ASN:HA | 5:E:567:PRO:HB3 | 1.83 | 0.59 |
| 4:A:52(A):VAL:HG22 | 4:A:71:THR:HG21 | 1.83 | 0.59 |
| 5:E:520:ASP:HB3 | 5:E:524:ALA:HB3 | 1.85 | 0.59 |
| 3:B:91:TRP:HZ3 | 4:A:50:TRP:CH2 | 2.21 | 0.58 |
| 3:B:28:ILE:O | 3:B:66:LYS:NZ | 2.35 | 0.58 |
| 5:E:502:VAL:HG21 | 5:E:537:PHE:HE2 | 1.68 | 0.57 |
| 5:E:491:PRO:HB3 | 5:E:563:VAL:HG12 | 1.86 | 0.57 |
| 5:E:479:GLY:HA3 | 5:E:492:LYS:HG3 | 1.86 | 0.56 |
| 2:H:39:GLN:HB2 | 2:H:45:LEU:HD23 | 1.88 | 0.56 |
| 4:A:168:ALA:HA | 4:A:178:LEU:HB3 | 1.87 | 0.56 |
| 5:E:520:ASP:C | 5:E:522:SER:H | 2.14 | 0.56 |
| 4:A:51:ILE:HD12 | 4:A:57:THR:HG22 | 1.88 | 0.55 |

*Continued on next page...*

*Continued from previous page...*

| Atom-1 | Atom-2 | Interatomic distance (Å) | Clash overlap (Å) |
|---|---|---|---|
| 4:A:68:THR:HG23 | 4:A:82(A):ARG:HH22 | 1.71 | 0.55 |
| 5:E:639:ARG:HG3 | 5:E:639:ARG:HH11 | 1.72 | 0.55 |
| 4:A:128:SER:OG | 4:A:214:LYS:O | 2.22 | 0.54 |
| 5:E:520:ASP:O | 5:E:522:SER:N | 2.36 | 0.54 |
| 2:H:51:ILE:HD13 | 2:H:71:THR:HG23 | 1.89 | 0.54 |
| 4:A:193:THR:HG23 | 4:A:210:LYS:HE3 | 1.90 | 0.53 |
| 5:E:427:LEU:HD23 | 5:E:441:LEU:HD21 | 1.91 | 0.52 |
| 3:B:46:LEU:HD21 | 3:B:49:TYR:HB3 | 1.92 | 0.52 |
| 5:E:510:THR:HG22 | 5:E:548:ASN:O | 2.09 | 0.52 |
| 5:E:509:PHE:HB3 | 5:E:511:PRO:O | 2.11 | 0.51 |
| 1:L:31:TYR:O | 1:L:66:LYS:NZ | 2.44 | 0.50 |
| 2:H:117:LYS:N | 2:H:146:PHE:O | 2.34 | 0.50 |
| 3:B:91:TRP:HZ3 | 4:A:50:TRP:HH2 | 1.58 | 0.50 |
| 5:E:556:ASN:OD1 | 5:E:557:SER:N | 2.44 | 0.50 |
| 4:A:6:GLN:H | 4:A:105:GLN:HE22 | 1.60 | 0.49 |
| 1:L:14:ALA:HA | 1:L:106(A):LEU:HB2 | 1.94 | 0.49 |
| 4:A:48:MET:HE1 | 4:A:90:TYR:HD2 | 1.78 | 0.49 |
| 5:E:543:ARG:HH12 | 5:E:598:GLY:C | 2.21 | 0.49 |
| 3:B:39:PHE:HB2 | 3:B:42:THR:HB | 1.95 | 0.48 |
| 5:E:544:PRO:HD2 | 5:E:550:PHE:CE1 | 2.49 | 0.48 |
| 1:L:183:GLU:OE1 | 1:L:183:GLU:N | 2.37 | 0.48 |
| 5:E:496:ILE:HG22 | 5:E:538:VAL:HG12 | 1.94 | 0.48 |
| 1:L:38:GLN:NE2 | 2:H:39:GLN:OE1 | 2.47 | 0.48 |
| 4:A:139:GLY:HA2 | 4:A:154:TRP:CH2 | 2.49 | 0.48 |
| 5:E:509:PHE:CD2 | 5:E:601:PRO:HB3 | 2.49 | 0.47 |
| 5:E:555:MET:HG2 | 5:E:617:HIS:CD2 | 2.49 | 0.47 |
| 2:H:98:SER:HB2 | 5:E:529:TRP:CZ3 | 2.50 | 0.47 |
| 3:B:80:SER:HA | 3:B:106:VAL:HG11 | 1.96 | 0.47 |
| 5:E:603:ILE:HD11 | 5:E:609:VAL:HG12 | 1.96 | 0.47 |
| 6:C:1:NAG:O4 | 6:C:2:NAG:O7 | 2.32 | 0.46 |
| 2:H:131:THR:O | 2:H:186:SER:HB2 | 2.15 | 0.46 |
| 5:E:633:VAL:O | 5:E:636:VAL:HG12 | 2.16 | 0.46 |
| 2:H:186:SER:HA | 2:H:189:LEU:HG | 1.96 | 0.46 |
| 4:A:13:LYS:HB2 | 4:A:14:PRO:CD | 2.46 | 0.46 |
| 4:A:159:LEU:HD21 | 4:A:182:VAL:HG21 | 1.96 | 0.46 |
| 4:A:13:LYS:HB2 | 4:A:14:PRO:HD3 | 1.97 | 0.46 |
| 5:E:472:ILE:HD11 | 5:E:569:CYS:SG | 2.55 | 0.46 |
| 5:E:608:LEU:HD23 | 5:E:609:VAL:O | 2.16 | 0.46 |
| 4:A:36:TRP:HD1 | 4:A:69:MET:HE3 | 1.81 | 0.46 |
| 2:H:139:GLY:HA2 | 2:H:154:TRP:CH2 | 2.51 | 0.45 |
| 4:A:36:TRP:CE2 | 4:A:80:MET:HB2 | 2.51 | 0.45 |

*Continued on next page...*

*Continued from previous page...*

| Atom-1 | Atom-2 | Interatomic distance (Å) | Clash overlap (Å) |
|---|---|---|---|
| 5:E:428:ASN:ND2 | 5:E:499:ALA:O | 2.45 | 0.45 |
| 3:B:120:PRO:HG3 | 3:B:130:ALA:HB1 | 1.98 | 0.45 |
| 1:L:19:VAL:HG21 | 1:L:104:LEU:HD11 | 1.99 | 0.44 |
| 3:B:83:GLU:HG3 | 3:B:105:THR:HA | 1.97 | 0.44 |
| 5:E:602:TRP:CE2 | 5:E:608:LEU:HD12 | 2.53 | 0.44 |
| 4:A:33:GLY:O | 4:A:34:ILE:HD13 | 2.18 | 0.44 |
| 5:E:639:ARG:HG3 | 5:E:639:ARG:NH1 | 2.33 | 0.44 |
| 4:A:6:GLN:NE2 | 4:A:104:GLY:HA3 | 2.33 | 0.44 |
| 5:E:470:GLY:O | 5:E:471:PRO:C | 2.61 | 0.44 |
| 1:L:79:GLN:O | 1:L:106:VAL:HG21 | 2.18 | 0.44 |
| 2:H:50:TYR:CE2 | 2:H:58:ASN:HB3 | 2.52 | 0.43 |
| 2:H:116:THR:HG21 | 2:H:202:PRO:C | 2.43 | 0.43 |
| 4:A:97:GLN:CD | 5:E:543:ARG:HD2 | 2.43 | 0.43 |
| 3:B:121:SER:HA | 4:A:214:LYS:NZ | 2.33 | 0.43 |
| 5:E:452:CYS:N | 5:E:453:PRO:HD2 | 2.31 | 0.43 |
| 2:H:123:PRO:HD3 | 2:H:209:LYS:HE2 | 1.99 | 0.43 |
| 4:A:186:SER:O | 4:A:189:LEU:HG | 2.17 | 0.43 |
| 5:E:512:SER:HA | 5:E:640:LEU:HD22 | 1.99 | 0.43 |
| 5:E:419:SER:HB2 | 5:E:420:TRP:CE3 | 2.54 | 0.43 |
| 5:E:473:SER:HB2 | 5:E:570:VAL:HB | 2.01 | 0.43 |
| 5:E:565:GLY:HA3 | 5:E:599:SER:O | 2.18 | 0.43 |
| 1:L:18:ARG:HD3 | 1:L:76:THR:HG22 | 2.01 | 0.42 |
| 3:B:84:ALA:O | 3:B:104:VAL:HG22 | 2.20 | 0.42 |
| 5:E:554:TRP:CH2 | 5:E:564:CYS:HB2 | 2.53 | 0.42 |
| 5:E:441:LEU:HB2 | 5:E:616:TRP:CZ2 | 2.54 | 0.42 |
| 4:A:51:ILE:HB | 4:A:69:MET:HE2 | 2.02 | 0.42 |
| 1:L:37:GLN:HB2 | 1:L:47:LEU:HD11 | 2.02 | 0.42 |
| 2:H:35:SER:HB3 | 2:H:95:TYR:CE1 | 2.54 | 0.42 |
| 2:H:8:GLY:HA3 | 2:H:20:LEU:HD23 | 2.01 | 0.42 |
| 4:A:36:TRP:CD1 | 4:A:69:MET:HE3 | 2.54 | 0.41 |
| 4:A:19:LYS:HA | 4:A:80:MET:O | 2.20 | 0.41 |
| 4:A:94:ARG:O | 4:A:100(A):VAL:HA | 2.20 | 0.41 |
| 3:B:145:LYS:HB2 | 3:B:197:THR:HB | 2.02 | 0.41 |
| 1:L:18:ARG:HG2 | 1:L:76:THR:HA | 2.02 | 0.41 |
| 2:H:47:TRP:HZ2 | 2:H:50:TYR:HD2 | 1.69 | 0.41 |
| 2:H:23:ALA:HA | 2:H:77:GLN:HG2 | 2.02 | 0.40 |
| 3:B:60:ASP:OD1 | 3:B:60:ASP:N | 2.54 | 0.40 |
| 3:B:95(C):PRO:HG2 | 4:A:50:TRP:CZ3 | 2.56 | 0.40 |
| 4:A:6:GLN:NE2 | 4:A:92:CYS:HB3 | 2.36 | 0.40 |
| 4:A:114:ALA:HB3 | 4:A:146:PHE:CZ | 2.56 | 0.40 |
| 4:A:61:GLN:H | 4:A:61:GLN:HG2 | 1.61 | 0.40 |

*Continued on next page...*

*Continued from previous page...*

| Atom-1 | Atom-2 | Interatomic distance (Å) | Clash overlap (Å) |
|--------|--------|--------------------------|-------------------|
| 4:A:112:SER:C | 4:A:114:ALA:H | 2.28 | 0.40 |

There are no symmetry-related clashes.

## 5.3   Torsion angles ⓘ

### 5.3.1   Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

| Mol | Chain | Analysed | Favoured | Allowed | Outliers | Percentiles | |
|-----|-------|----------|----------|---------|----------|-------------|---|
| 1 | L | 206/216 (95%) | 195 (95%) | 9 (4%) | 2 (1%) | 13 | 39 |
| 2 | H | 220/224 (98%) | 216 (98%) | 3 (1%) | 1 (0%) | 25 | 54 |
| 3 | B | 215/217 (99%) | 198 (92%) | 17 (8%) | 0 | 100 | 100 |
| 4 | A | 218/223 (98%) | 204 (94%) | 8 (4%) | 6 (3%) | 4 | 20 |
| 5 | E | 228/262 (87%) | 192 (84%) | 30 (13%) | 6 (3%) | 4 | 22 |
| All | All | 1087/1142 (95%) | 1005 (92%) | 67 (6%) | 15 (1%) | 9 | 31 |

All (15) Ramachandran outliers are listed below:

| Mol | Chain | Res | Type |
|-----|-------|-----|------|
| 5 | E | 449 | SER |
| 5 | E | 450 | SER |
| 4 | A | 128 | SER |
| 5 | E | 471 | PRO |
| 1 | L | 28 | ILE |
| 1 | L | 151 | ASP |
| 2 | H | 132 | SER |
| 4 | A | 113 | SER |
| 5 | E | 598 | GLY |
| 4 | A | 14 | PRO |
| 4 | A | 146 | PHE |
| 4 | A | 13 | LYS |
| 4 | A | 190 | GLY |

*Continued on next page...*

*Continued from previous page...*

| Mol | Chain | Res | Type |
|-----|-------|-----|------|
| 5 | E | 470 | GLY |
| 5 | E | 510 | THR |

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

| Mol | Chain | Analysed | Rotameric | Outliers | Percentiles | |
|-----|-------|----------|-----------|----------|-----|-----|
| 1 | L | 177/185 (96%) | 174 (98%) | 3 (2%) | 56 | 74 |
| 2 | H | 189/191 (99%) | 187 (99%) | 2 (1%) | 70 | 81 |
| 3 | B | 184/184 (100%) | 184 (100%) | 0 | 100 | 100 |
| 4 | A | 185/187 (99%) | 184 (100%) | 1 (0%) | 86 | 91 |
| 5 | E | 195/219 (89%) | 191 (98%) | 4 (2%) | 48 | 69 |
| All | All | 930/966 (96%) | 920 (99%) | 10 (1%) | 70 | 81 |

All (10) residues with a non-rotameric sidechain are listed below:

| Mol | Chain | Res | Type |
|-----|-------|-----|------|
| 1 | L | 26 | HIS |
| 1 | L | 28 | ILE |
| 1 | L | 171 | LYS |
| 2 | H | 132 | SER |
| 2 | H | 186 | SER |
| 4 | A | 100(A) | VAL |
| 5 | E | 471 | PRO |
| 5 | E | 512 | SER |
| 5 | E | 514 | VAL |
| 5 | E | 620 | CYS |

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (9) such sidechains are listed below:

| Mol | Chain | Res | Type |
|-----|-------|-----|------|
| 1 | L | 37 | GLN |
| 1 | L | 170 | ASN |

*Continued on next page...*

*Continued from previous page...*

| Mol | Chain | Res | Type |
|-----|-------|-----|------|
| 2 | H | 58 | ASN |
| 3 | B | 31 | ASN |
| 3 | B | 53 | GLN |
| 4 | A | 6 | GLN |
| 4 | A | 171 | GLN |
| 4 | A | 199 | ASN |
| 5 | E | 577 | ASN |

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

1 non-standard protein/DNA/RNA residue is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

| Mol | Type | Chain | Res | Link | Bond lengths | | | Bond angles | | |
|-----|------|-------|-----|------|--------|------|--------|--------|------|--------|
| | | | | | Counts | RMSZ | #$|Z| > 2$ | Counts | RMSZ | #$|Z| > 2$ |
| 2 | PCA | H | 1 | 2 | 7,8,9 | 0.64 | 0 | 9,10,12 | 1.24 | 1 (11%) |

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

| Mol | Type | Chain | Res | Link | Chirals | Torsions | Rings |
|-----|------|-------|-----|------|---------|----------|-------|
| 2 | PCA | H | 1 | 2 | - | 0/0/11/13 | 0/1/1/1 |

There are no bond length outliers.

All (1) bond angle outliers are listed below:

| Mol | Chain | Res | Type | Atoms | Z | Observed($^o$) | Ideal($^o$) |
|-----|-------|-----|------|-------|---|----------------|-------------|
| 2 | H | 1 | PCA | CB-CA-N | 2.17 | 109.22 | 103.24 |

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.5 Carbohydrates ⓘ

7 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

| Mol | Type | Chain | Res | Link | Bond lengths | | | Bond angles | | |
|-----|------|-------|-----|------|--------|------|--------|--------|------|--------|
| | | | | | Counts | RMSZ | #$|Z|$ > 2 | Counts | RMSZ | #$|Z|$ > 2 |
| 6 | NAG | C | 1 | 5,6 | 14,14,15 | 0.68 | 0 | 17,19,21 | 0.98 | 1 (5%) |
| 6 | NAG | C | 2 | 6 | 14,14,15 | 0.69 | 0 | 17,19,21 | 2.57 | 5 (29%) |
| 6 | NAG | D | 1 | 5,6 | 14,14,15 | 0.56 | 0 | 17,19,21 | 2.68 | 3 (17%) |
| 6 | NAG | D | 2 | 6 | 14,14,15 | 0.68 | 0 | 17,19,21 | 0.92 | 0 |
| 7 | NAG | F | 1 | 7,5 | 14,14,15 | 0.71 | 0 | 17,19,21 | 1.04 | 1 (5%) |
| 7 | NAG | F | 2 | 7 | 14,14,15 | 0.68 | 0 | 17,19,21 | 0.88 | 1 (5%) |
| 7 | BMA | F | 3 | 7 | 11,11,12 | 0.78 | 0 | 15,15,17 | 1.79 | 1 (6%) |

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

| Mol | Type | Chain | Res | Link | Chirals | Torsions | Rings |
|-----|------|-------|-----|------|---------|----------|-------|
| 6 | NAG | C | 1 | 5,6 | - | 3/6/23/26 | 0/1/1/1 |
| 6 | NAG | C | 2 | 6 | - | 1/6/23/26 | 0/1/1/1 |
| 6 | NAG | D | 1 | 5,6 | - | 0/6/23/26 | 0/1/1/1 |
| 6 | NAG | D | 2 | 6 | - | 0/6/23/26 | 0/1/1/1 |
| 7 | NAG | F | 1 | 7,5 | - | 0/6/23/26 | 0/1/1/1 |
| 7 | NAG | F | 2 | 7 | - | 0/6/23/26 | 0/1/1/1 |
| 7 | BMA | F | 3 | 7 | - | 0/2/19/22 | 0/1/1/1 |

There are no bond length outliers.

All (12) bond angle outliers are listed below:

| Mol | Chain | Res | Type | Atoms | Z | Observed($^o$) | Ideal($^o$) |
|---|---|---|---|---|---|---|---|
| 6 | D | 1 | NAG | C1-O5-C5 | 9.93 | 125.49 | 112.19 |
| 6 | C | 2 | NAG | C2-N2-C7 | 7.18 | 132.52 | 122.90 |
| 7 | F | 3 | BMA | C1-O5-C5 | 5.80 | 119.96 | 112.19 |
| 6 | C | 2 | NAG | C1-C2-N2 | 5.42 | 118.98 | 110.43 |
| 6 | C | 2 | NAG | O5-C1-C2 | -3.24 | 106.28 | 111.29 |
| 7 | F | 1 | NAG | O5-C1-C2 | -2.43 | 107.53 | 111.29 |
| 6 | D | 1 | NAG | C2-N2-C7 | 2.38 | 126.08 | 122.90 |
| 6 | D | 1 | NAG | O5-C5-C4 | 2.36 | 116.56 | 110.83 |
| 6 | C | 1 | NAG | O5-C1-C2 | -2.22 | 107.86 | 111.29 |
| 6 | C | 2 | NAG | C1-O5-C5 | 2.20 | 115.13 | 112.19 |
| 6 | C | 2 | NAG | O7-C7-N2 | 2.19 | 125.84 | 121.98 |
| 7 | F | 2 | NAG | O5-C1-C2 | -2.05 | 108.12 | 111.29 |

There are no chirality outliers.

All (4) torsion outliers are listed below:

| Mol | Chain | Res | Type | Atoms |
|---|---|---|---|---|
| 6 | C | 2 | NAG | C1-C2-N2-C7 |
| 6 | C | 1 | NAG | O5-C5-C6-O6 |
| 6 | C | 1 | NAG | C4-C5-C6-O6 |
| 6 | C | 1 | NAG | C1-C2-N2-C7 |

There are no ring outliers.

2 monomers are involved in 1 short contact:

| Mol | Chain | Res | Type | Clashes | Symm-Clashes |
|---|---|---|---|---|---|
| 6 | C | 1 | NAG | 1 | 0 |
| 6 | C | 2 | NAG | 1 | 0 |

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.

## Oligosaccharide Chain C


Bond lengths


Bond angles


Torsions


Rings

## Oligosaccharide Chain D


Bond lengths


Bond angles


Torsions


Rings

**Oligosaccharide Chain F**



Bond lengths

Bond angles

Torsions

Rings

## 5.6 Ligand geometry (i)

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

| Mol | Type | Chain | Res | Link | Bond lengths | | | Bond angles | | |
| --- | --- | --- | --- | --- | --- | --- | --- | --- | --- | --- |
| | | | | | Counts | RMSZ | #$|Z| > 2$ | Counts | RMSZ | #$|Z| > 2$ |
| 8 | NAG | E | 701 | 5 | 14,14,15 | 0.77 | 0 | 17,19,21 | 1.34 | 2 (11%) |
| 8 | NAG | E | 702 | 5 | 14,14,15 | 0.73 | 0 | 17,19,21 | 0.79 | 0 |
| 8 | NAG | E | 703 | 5 | 14,14,15 | 0.67 | 0 | 17,19,21 | 1.47 | 2 (11%) |
| 9 | MAN | E | 704 | - | 11,11,12 | 0.76 | 0 | 15,15,17 | 1.05 | 1 (6%) |

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

| Mol | Type | Chain | Res | Link | Chirals | Torsions | Rings |
|-----|------|-------|-----|------|---------|----------|-------|
| 8 | NAG | E | 701 | 5 | - | 3/6/23/26 | 0/1/1/1 |
| 8 | NAG | E | 702 | 5 | - | 2/6/23/26 | 0/1/1/1 |
| 8 | NAG | E | 703 | 5 | - | 3/6/23/26 | 0/1/1/1 |
| 9 | MAN | E | 704 | - | - | 0/2/19/22 | 0/1/1/1 |

There are no bond length outliers.

All (5) bond angle outliers are listed below:

| Mol | Chain | Res | Type | Atoms | Z | Observed($^o$) | Ideal($^o$) |
|-----|-------|-----|------|-------|---|-----------|--------|
| 8 | E | 703 | NAG | C2-N2-C7 | 4.01 | 128.27 | 122.90 |
| 8 | E | 701 | NAG | C2-N2-C7 | 3.29 | 127.31 | 122.90 |
| 9 | E | 704 | MAN | C1-O5-C5 | 2.93 | 116.11 | 112.19 |
| 8 | E | 701 | NAG | C1-O5-C5 | 2.43 | 115.44 | 112.19 |
| 8 | E | 703 | NAG | O5-C1-C2 | -2.33 | 107.69 | 111.29 |

There are no chirality outliers.

All (8) torsion outliers are listed below:

| Mol | Chain | Res | Type | Atoms |
|-----|-------|-----|------|-------|
| 8 | E | 703 | NAG | C1-C2-N2-C7 |
| 8 | E | 701 | NAG | O5-C5-C6-O6 |
| 8 | E | 701 | NAG | C4-C5-C6-O6 |
| 8 | E | 702 | NAG | O5-C5-C6-O6 |
| 8 | E | 702 | NAG | C4-C5-C6-O6 |
| 8 | E | 703 | NAG | O5-C5-C6-O6 |
| 8 | E | 701 | NAG | C3-C2-N2-C7 |
| 8 | E | 703 | NAG | C3-C2-N2-C7 |

There are no ring outliers.

1 monomer is involved in 1 short contact:

| Mol | Chain | Res | Type | Clashes | Symm-Clashes |
|-----|-------|-----|------|---------|--------------|
| 8 | E | 701 | NAG | 1 | 0 |

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring

in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



**Ligand NAG E 701**

Bond lengths

Bond angles

Torsions

Rings

## Ligand NAG E 702


Bond lengths


Bond angles


Torsions


Rings

## Ligand NAG E 703


Bond lengths


Bond angles


Torsions


Rings

**Ligand MAN E 704**



Bond lengths

Bond angles

Torsions

Rings

## 5.7 Other polymers ⓘ

There are no such residues in this entry.

## 5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

# 6 Fit of model and data ⓘ

## 6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, $95^{th}$ percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

| Mol | Chain | Analysed | <RSRZ> | #RSRZ>2 | | | OWAB(Å²) | Q<0.9 |
|-----|-------|----------|--------|---------|------|------|----------|-------|
| 1 | L | 208/216 (96%) | -0.03 | 0 | 100 | 100 | 143, 198, 237, 260 | 0 |
| 2 | H | 221/224 (98%) | 0.02 | 1 (0%) | 87 | 83 | 149, 193, 235, 261 | 0 |
| 3 | B | 217/217 (100%) | -0.14 | 9 (4%) | 42 | 36 | 72, 124, 169, 210 | 0 |
| 4 | A | 220/223 (98%) | -0.26 | 0 | 100 | 100 | 68, 114, 151, 217 | 0 |
| 5 | E | 230/262 (87%) | 0.11 | 9 (3%) | 44 | 37 | 118, 167, 229, 264 | 0 |
| All | All | 1096/1142 (95%) | -0.06 | 19 (1%) | 69 | 59 | 68, 165, 227, 264 | 0 |

All (19) RSRZ outliers are listed below:

| Mol | Chain | Res | Type | RSRZ |
|-----|-------|-----|------|------|
| 3 | B | 37 | GLN | 5.2 |
| 2 | H | 102 | VAL | 5.1 |
| 5 | E | 457 | ALA | 4.0 |
| 5 | E | 420 | TRP | 3.7 |
| 3 | B | 96 | TRP | 3.1 |
| 5 | E | 498 | PRO | 3.1 |
| 3 | B | 36 | TYR | 3.0 |
| 3 | B | 181 | LEU | 2.9 |
| 3 | B | 86 | TYR | 2.8 |
| 3 | B | 185 | ASP | 2.6 |
| 5 | E | 595 | SER | 2.6 |
| 3 | B | 27(A) | SER | 2.5 |
| 5 | E | 427 | LEU | 2.4 |
| 3 | B | 135 | LEU | 2.3 |
| 5 | E | 442 | PHE | 2.2 |
| 3 | B | 47 | LEU | 2.2 |
| 5 | E | 505 | PRO | 2.1 |
| 5 | E | 631 | MET | 2.1 |
| 5 | E | 553 | THR | 2.0 |

## 6.2　Non-standard residues in protein, DNA, RNA chains ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, $95^{th}$ percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

| Mol | Type | Chain | Res | Atoms | RSCC | RSR | B-factors($\text{Å}^2$) | Q<0.9 |
|-----|------|-------|-----|-------|------|-----|-------------------------|-------|
| 2 | PCA | H | 1 | 8/9 | 0.37 | 0.10 | 213,240,252,260 | 0 |

## 6.3　Carbohydrates ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, $95^{th}$ percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

| Mol | Type | Chain | Res | Atoms | RSCC | RSR | B-factors($\text{Å}^2$) | Q<0.9 |
|-----|------|-------|-----|-------|------|-----|-------------------------|-------|
| 6 | NAG | C | 1 | 14/15 | - | - | 165,184,196,220 | 0 |
| 6 | NAG | C | 2 | 14/15 | - | - | 220,236,250,255 | 0 |
| 6 | NAG | D | 1 | 14/15 | - | - | 111,141,169,182 | 0 |
| 6 | NAG | D | 2 | 14/15 | - | - | 168,188,198,201 | 0 |
| 7 | NAG | F | 2 | 14/15 | 0.54 | 0.09 | 119,175,183,196 | 0 |
| 7 | NAG | F | 1 | 14/15 | 0.79 | 0.07 | 139,176,203,209 | 0 |
| 7 | BMA | F | 3 | 11/12 | - | - | 154,178,199,203 | 0 |

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.

**Electron density around Chain C:**

$2mF_o$-$DF_c$ (at 0.7 rmsd) in gray
$mF_o$-$DF_c$ (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around Chain D:**

2mF$_o$-DF$_c$ (at 0.7 rmsd) in gray
mF$_o$-DF$_c$ (at 3 rmsd) in purple (negative)
and green (positive)

Electron density around Chain F:

$2mF_o$-$DF_c$ (at 0.7 rmsd) in gray
$mF_o$-$DF_c$ (at 3 rmsd) in purple (negative)
and green (positive)

## 6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, $95^{th}$ percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

| Mol | Type | Chain | Res | Atoms | RSCC | RSR | B-factors($\text{Å}^2$) | Q<0.9 |
|-----|------|-------|-----|-------|------|-----|-------------------------|-------|
| 9 | MAN | E | 704 | 11/12 | 0.36 | 0.10 | 305,334,349,361 | 0 |
| 8 | NAG | E | 703 | 14/15 | 0.64 | 0.10 | 197,219,227,238 | 0 |
| 8 | NAG | E | 702 | 14/15 | 0.74 | 0.07 | 173,213,224,228 | 0 |
| 8 | NAG | E | 701 | 14/15 | 0.78 | 0.10 | 161,185,198,205 | 0 |

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

**Electron density around MAN E 704:**

$2mF_o$-$DF_c$ (at 0.7 rmsd) in gray
$mF_o$-$DF_c$ (at 3 rmsd) in purple (negative)
and green (positive)



**Electron density around NAG E 703:**

$2mF_o$-$DF_c$ (at 0.7 rmsd) in gray
$mF_o$-$DF_c$ (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around NAG E 702:**

$2mF_o\text{-}DF_c$ (at 0.7 rmsd) in gray
$mF_o\text{-}DF_c$ (at 3 rmsd) in purple (negative)
and green (positive)



**Electron density around NAG E 701:**

$2mF_o\text{-}DF_c$ (at 0.7 rmsd) in gray
$mF_o\text{-}DF_c$ (at 3 rmsd) in purple (negative)
and green (positive)

## 6.5 Other polymers ⓘ

There are no such residues in this entry.