Full wwPDB X-ray Structure Validation Report ⓘ

Aug 31, 2026 – 03:01 PM EDT

PDB ID	:	9MUY / pdb_00009muy
Title	:	anti-IL6 designed Fab
Authors	:	Kiefer, J.R.; Frey, N.C.; Alberstein, R.G.; Seeger, F.; Watkins, A.M.; Gligorijevic, V.; Dou, Y.; Tang, Y.; Regev, A.; Bonneau, R.
Deposited on	:	2025-01-14
Resolution	:	1.97 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org*
A user guide is available at
https://www.wwpdb.org/validation/2017/XrayValidationReportHelp
with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at
http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references ⓘ) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.015 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.50

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:
X-RAY DIFFRACTION

The reported resolution of this entry is 1.97 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1506 (1.98-1.98)
Clashscore	190562	1534 (1.98-1.98)
Ramachandran outliers	187476	1518 (1.98-1.98)
Sidechain outliers	187428	1518 (1.98-1.98)
RSRZ outliers	180081	1506 (1.98-1.98)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	H	228	8% 92%
2	L	213	22% 87% 13%

2 Entry composition (i)

There are 6 unique types of molecules in this entry. The entry contains 6776 atoms, of which 3191 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called anti-IL6 designed Fab heavy chain.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	H	218	Total	C	H	N	O	S	0	4	0
			3318	1083	1624	274	331	6			

- Molecule 2 is a protein called anti-IL6 designed Fab light chain.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
2	L	213	Total	C	H	N	O	S	0	2	0
			3197	1030	1549	277	335	6			

- Molecule 3 is ZINC ION (CCD ID: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	H	1	Total	Zn	0	0
			1	1		
3	L	1	Total	Zn	0	0
			1	1		

- Molecule 4 is 1,2-ETHANEDIOL (CCD ID: EDO) (formula: $C_2H_6O_2$).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	H	1	Total C H O 10 2 6 2	0	0
4	H	1	Total C H O 10 2 6 2	0	0
4	H	1	Total C H O 10 2 6 2	0	0

- Molecule 5 is CHLORIDE ION (CCD ID: CL) (formula: Cl).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	H	4	Total Cl 4 4	0	0

- Molecule 6 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	H	133	Total O 135 135	0	2
6	L	90	Total O 90 90	0	0

3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: anti-IL6 designed Fab heavy chain

Chain H:



• Molecule 2: anti-IL6 designed Fab light chain

Chain L:



4 Data and refinement statistics (i)

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α, β, γ	91.51Å 64.35Å 104.57Å 90.00° 114.66° 90.00°	Depositor
Resolution (Å)	33.33 – 1.97 33.33 – 1.97	Depositor EDS
% Data completeness (in resolution range)	99.5 (33.33-1.97) 99.6 (33.33-1.97)	Depositor EDS
R_{merge}	0.08	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) >$ [1]	1.70 (at 1.97Å)	Xtriage
Refinement program	PHENIX (1.21.2_5419: ???)	Depositor
R, R_{free}	0.208 , 0.223 0.208 , 0.225	Depositor DCC
R_{free} test set	1925 reflections (4.93%)	wwPDB-VP
Wilson B-factor (Å^2)	34.6	Xtriage
Anisotropy	0.119	Xtriage
Bulk solvent k_{sol}(e/Å^3), B_{sol}(Å^2)	0.38 , 45.6	EDS
L-test for twinning[2]	$< \|L\| > = 0.50$, $< L^2 > = 0.34$	Xtriage
Estimated twinning fraction	0.012 for h,-k,-h-l	Xtriage
F_o,F_c correlation	0.95	EDS
Total number of atoms	6776	wwPDB-VP
Average B, all atoms (Å^2)	59.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.58% of the height of the origin peak. No significant pseudotranslation is detected.*

[1] Intensities estimated from amplitudes.

[2] Theoretical values of $< |L| >$, $< L^2 >$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: CL, ZN, EDO

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	#$\|Z\|>5$	RMSZ	#$\|Z\|>5$
1	H	0.30	1/1758 (0.1%)	0.48	0/2399
2	L	0.29	0/1693	0.51	1/2305 (0.0%)
All	All	0.30	1/3451 (0.0%)	0.50	1/4704 (0.0%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	H	100(G)	PHE	C-O	-7.32	1.15	1.23

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})
2	L	170	ASP	CA-CB-CG	5.33	117.93	112.60

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	H	1694	1624	1611	5	0
2	L	1648	1549	1537	24	1
3	H	1	0	0	0	1
3	L	1	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	H	12	18	18	0	0
5	H	4	0	0	0	0
6	H	135	0	0	1	0
6	L	90	0	0	1	0
All	All	3585	3191	3166	26	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (26) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:L:4:MET:HE3	2:L:23:CYS:SG	2.20	0.82
2:L:148:TRP:CE3	2:L:179:LEU:HD22	2.27	0.69
2:L:45:LYS:HE2	6:L:431:HOH:O	1.98	0.63
2:L:4:MET:CE	2:L:23:CYS:SG	2.94	0.56
2:L:132:VAL:HG12	2:L:148:TRP:HH2	1.73	0.53
2:L:211:ARG:HG2	2:L:211:ARG:HH11	1.74	0.52
2:L:80:PRO:HA	2:L:106:LEU:HD13	1.93	0.51
1:H:143:LYS:HA	1:H:177:SER:OG	2.11	0.50
2:L:132:VAL:HB	2:L:148:TRP:CH2	2.46	0.50
2:L:132:VAL:HG12	2:L:148:TRP:CH2	2.48	0.49
2:L:146:VAL:HG12	2:L:196:VAL:HG22	1.94	0.48
2:L:186:TYR:O	2:L:192:TYR:OH	2.28	0.48
2:L:83:VAL:HG13	2:L:104:LEU:O	2.12	0.48
2:L:132:VAL:HB	2:L:148:TRP:CZ3	2.50	0.47
2:L:1:ASP:O	2:L:1:ASP:CG	2.57	0.46
1:H:100(F):ARG:CZ	2:L:49:TYR:CE2	2.99	0.46
2:L:112:ALA:HB1	2:L:201:LEU:HD21	2.00	0.44
2:L:112:ALA:HB1	2:L:201:LEU:CD2	2.48	0.43
2:L:132:VAL:CG1	2:L:148:TRP:CH2	3.01	0.43
1:H:123:PRO:HD2	2:L:121:SER:HB2	2.01	0.43
1:H:64:LYS:NZ	6:H:403:HOH:O	2.42	0.42
2:L:175:LEU:HD13	2:L:176:SER:N	2.35	0.41
2:L:132:VAL:CG1	2:L:148:TRP:CZ3	3.03	0.41
2:L:136:LEU:HD13	2:L:175:LEU:HD12	2.03	0.41
1:H:100(E):ASP:OD1	2:L:91:ARG:HD2	2.21	0.41
2:L:201:LEU:HD13	2:L:205:VAL:CG2	2.51	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:L:52:HIS:HE2	3:H:301:ZN:ZN[2_556]	1.35	0.25

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	H	218/228 (96%)	210 (96%)	7 (3%)	1 (0%)	24 14
2	L	213/213 (100%)	206 (97%)	7 (3%)	0	100 100
All	All	431/441 (98%)	416 (96%)	14 (3%)	1 (0%)	43 35

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	H	189	LEU

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	H	188/194 (97%)	188 (100%)	0	100 100
2	L	184/189 (97%)	184 (100%)	0	100 100
All	All	372/383 (97%)	372 (100%)	0	100 100

There are no protein residues with a non-rotameric sidechain to report.

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (6) such sidechains are listed below:

Mol	Chain	Res	Type
1	H	3	GLN
1	H	192	GLN
2	L	30	ASN
2	L	53	ASN
2	L	92	ASN
2	L	147	GLN

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no oligosaccharides in this entry.

5.6 Ligand geometry (i)

Of 9 ligands modelled in this entry, 6 are monoatomic - leaving 3 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\#\|Z\| > 2$	Counts	RMSZ	$\#\|Z\| > 2$
4	EDO	H	303	-	3,3,3	0.46	0	2,2,2	0.30	0
4	EDO	H	304	-	3,3,3	0.43	0	2,2,2	0.46	0
4	EDO	H	302	-	3,3,3	0.46	0	2,2,2	0.27	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	EDO	H	303	-	-	0/1/1/1	-
4	EDO	H	304	-	-	1/1/1/1	-
4	EDO	H	302	-	-	0/1/1/1	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (1) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	H	304	EDO	O1-C1-C2-O2

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95^{th} percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2			OWAB(Å^2)	Q<0.9
1	H	218/228 (95%)	0.40	19 (8%)	16	22	16, 48, 94, 116	2 (0%)
2	L	213/213 (100%)	1.02	47 (22%)	2	2	24, 62, 116, 138	1 (0%)
All	All	431/441 (97%)	0.70	66 (15%)	5	7	16, 55, 112, 138	3 (0%)

All (66) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	L	194	CYS	5.6
1	H	1	TRP	5.1
2	L	10[A]	PHE	5.0
2	L	146	VAL	4.9
2	L	148	TRP	4.8
2	L	150	VAL	4.8
2	L	193	ALA	4.8
1	H	127	SER	4.7
1	H	100(D)	TYR	4.6
2	L	191	VAL	4.3
1	H	99	ASP	4.1
2	L	189	HIS	4.0
2	L	205	VAL	4.0
1	H	214	LYS	3.8
2	L	149	LYS	3.8
1	H	189	LEU	3.7
2	L	181	LEU	3.7
2	L	201	LEU	3.6
2	L	186	TYR	3.5
2	L	153	ALA	3.2
1	H	191	THR	3.2
2	L	154	LEU	3.2
2	L	196	VAL	3.2
2	L	200	GLY	3.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
2	L	125	LEU	3.1
2	L	179	LEU	3.1
2	L	126	LYS	3.1
2	L	192	TYR	3.1
2	L	130	ALA	3.1
2	L	214	CYS	3.0
2	L	152	ASN	3.0
1	H	135	THR	2.9
1	H	100	GLY	2.9
2	L	112	ALA	2.9
1	H	100(C)	TYR	2.8
2	L	117	ILE	2.8
2	L	190	LYS	2.8
1	H	190	GLY	2.7
2	L	1	ASP	2.7
1	H	100(B)	TYR	2.7
2	L	183	LYS	2.7
2	L	110	VAL	2.7
2	L	115	VAL	2.7
2	L	209	PHE	2.6
2	L	184	ALA	2.6
1	H	124	LEU	2.4
2	L	212	GLY	2.4
1	H	16	ARG	2.4
2	L	197	THR	2.4
2	L	187	GLU	2.4
2	L	169	LYS	2.4
2	L	204	PRO	2.3
2	L	132	VAL	2.3
2	L	151	ASP	2.3
1	H	100(A)	SER	2.3
1	H	188	SER	2.2
2	L	195	GLU	2.1
2	L	199	GLN	2.1
2	L	188	LYS	2.1
2	L	116	PHE	2.1
2	L	49	TYR	2.1
1	H	136	ALA	2.1
1	H	98	TYR	2.1
2	L	157	GLY	2.0
1	H	184	VAL	2.0
2	L	133	VAL	2.0

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

There are no oligosaccharides in this entry.

6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å^2)	Q<0.9
3	ZN	L	301	1/1	0.57	0.23	195,195,195,195	0
4	EDO	H	303	4/4	0.58	0.17	60,72,85,99	0
4	EDO	H	304	4/4	0.88	0.17	51,62,69,74	0
4	EDO	H	302	4/4	0.91	0.09	41,49,55,65	0
5	CL	H	306	1/1	0.92	0.14	62,62,62,62	0
5	CL	H	307	1/1	0.94	0.09	79,79,79,79	0
5	CL	H	308	1/1	0.95	0.15	70,70,70,70	0
3	ZN	H	301	1/1	0.98	0.05	27,27,27,27	1
5	CL	H	305	1/1	0.99	0.07	27,27,27,27	0

6.5 Other polymers (i)

There are no such residues in this entry.