



Full wwPDB X-ray Structure Validation Report ⓘ

Aug 31, 2026 – 03:11 PM EDT

PDB ID : 9MVJ / pdb_00009mvj
Title : anti-EGFR designed Fab
Authors : Kiefer, J.R.; Frey, N.C.; Alberstein, R.G.; Seeger, F.; Watkins, A.M.; Gligorijevic, V.; Dou, Y.; Tang, Y.; Regev, A.; Bonneau, R.
Deposited on : 2025-01-15
Resolution : 1.95 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.015 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.50


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.95 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	3494 (1.96-1.96)
Clashscore	190562	3612 (1.96-1.96)
Ramachandran outliers	187476	3587 (1.96-1.96)
Sidechain outliers	187428	3587 (1.96-1.96)
RSRZ outliers	180081	3495 (1.96-1.96)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	H	225	14% 92%
2	L	213	18% 92% 8%

2 Entry composition [i](#)

There are 7 unique types of molecules in this entry. The entry contains 6924 atoms, of which 3209 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called anti-EGFR designed Fab heavy chain.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	H	216	Total	C	H	N	O	S	0	6	0
			3240	1049	1591	270	324	6			

- Molecule 2 is a protein called anti-EGFR designed Fab light chain.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
2	L	213	Total	C	H	N	O	S	0	2	0
			3129	1003	1516	266	339	5			

- Molecule 3 is 1,2-ETHANEDIOL (CCD ID: EDO) (formula: C₂H₆O₂).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	H	1	Total	C	H	O	0	0
			10	2	6	2		
3	H	1	Total	C	H	O	0	0
			10	2	6	2		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	H	1	Total	C	H	O	0	0
			10	2	6	2		
3	H	1	Total	C	H	O	0	0
			10	2	6	2		
3	H	1	Total	C	H	O	0	0
			10	2	6	2		
3	H	1	Total	C	H	O	0	0
			10	2	6	2		
3	H	1	Total	C	H	O	0	0
			10	2	6	2		
3	H	1	Total	C	H	O	0	0
			10	2	6	2		
3	H	1	Total	C	H	O	0	0
			10	2	6	2		
3	L	1	Total	C	H	O	0	0
			10	2	6	2		
3	L	1	Total	C	H	O	0	0
			10	2	6	2		
3	L	1	Total	C	H	O	0	0
			10	2	6	2		
3	L	1	Total	C	H	O	0	0
			10	2	6	2		
3	L	1	Total	C	H	O	0	0
			10	2	6	2		

- Molecule 4 is SULFATE ION (CCD ID: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	H	1	Total	O	S	0	0
			5	4	1		
4	H	1	Total	O	S	0	0
			5	4	1		
4	H	1	Total	O	S	0	0
			5	4	1		
4	H	1	Total	O	S	0	0
			5	4	1		
4	H	1	Total	O	S	0	0
			5	4	1		
4	H	1	Total	O	S	0	0
			5	4	1		
4	L	1	Total	O	S	0	0
			5	4	1		
4	L	1	Total	O	S	0	0
			5	4	1		

- Molecule 5 is CHLORIDE ION (CCD ID: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	H	2	Total	Cl	0	0
			2	2		
5	L	4	Total	Cl	0	0
			4	4		

- Molecule 6 is SODIUM ION (CCD ID: NA) (formula: Na).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	L	1	Total 1	Na 1	0	0


- Molecule 7 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	H	169	Total 169	O 169	0	1
7	L	159	Total 159	O 159	0	1


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: anti-EGFR designed Fab heavy chain



- Molecule 2: anti-EGFR designed Fab light chain



4 Data and refinement statistics

Property	Value	Source
Space group	H 3 2	Depositor
Cell constants a, b, c, α , β , γ	197.09Å 197.09Å 108.96Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	39.73 – 1.95 39.73 – 1.95	Depositor EDS
% Data completeness (in resolution range)	100.0 (39.73-1.95) 100.0 (39.73-1.95)	Depositor EDS
R_{merge}	0.11	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.65 (at 1.95Å)	Xtriage
Refinement program	PHENIX (1.21.2_5419: ???)	Depositor
R, R_{free}	0.180 , 0.198 0.181 , 0.201	Depositor DCC
R_{free} test set	2939 reflections (4.97%)	wwPDB-VP
Wilson B-factor (Å ²)	34.7	Xtriage
Anisotropy	0.543	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.42 , 63.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.51$, $\langle L^2 \rangle = 0.34$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	6924	wwPDB-VP
Average B, all atoms (Å ²)	52.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.25% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: NA, SO4, CL, EDO

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	H	0.21	0/1712	0.38	0/2334
2	L	0.20	0/1648	0.40	0/2251
All	All	0.21	0/3360	0.39	0/4585

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	H	1649	1591	1568	5	0
2	L	1613	1516	1514	7	0
3	H	44	66	66	1	0
3	L	24	36	36	0	0
4	H	40	0	0	0	0
4	L	10	0	0	0	0
5	H	2	0	0	0	0
5	L	4	0	0	0	0
6	L	1	0	0	0	0
7	H	169	0	0	0	0
7	L	159	0	0	0	0
All	All	3715	3209	3184	12	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (12) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:L:149:LYS:HG3	2:L:154:LEU:HD23	1.82	0.60
1:H:12:VAL:HG11	1:H:82(C):LEU:HD12	1.84	0.60
1:H:148:GLU:OE1	3:H:311:EDO:O1	2.25	0.52
1:H:178:LEU:C	1:H:178:LEU:HD12	2.37	0.49
2:L:61:ARG:HD2	2:L:77:PRO:O	2.16	0.46
2:L:158:ASN:O	2:L:179:LEU:HD12	2.16	0.46
2:L:135:LEU:C	2:L:136:LEU:HD12	2.41	0.46
2:L:147:GLN:O	2:L:194:CYS:HA	2.18	0.43
2:L:151:ASP:HA	2:L:191:VAL:HB	2.00	0.43
1:H:195:ILE:HA	1:H:209:LYS:O	2.19	0.43
2:L:189:HIS:O	2:L:211:ARG:NE	2.50	0.42
1:H:184:VAL:HG11	1:H:194:TYR:CZ	2.56	0.41

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	H	218/225 (97%)	214 (98%)	4 (2%)	0	100	100
2	L	212/213 (100%)	205 (97%)	7 (3%)	0	100	100
All	All	430/438 (98%)	419 (97%)	11 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	H	183/192 (95%)	181 (99%)	2 (1%)	65	64
2	L	182/186 (98%)	178 (98%)	4 (2%)	45	40
All	All	365/378 (97%)	359 (98%)	6 (2%)	53	52

All (6) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	H	193	THR
1	H	206	LYS
2	L	114	SER
2	L	127	SER
2	L	132	VAL
2	L	211	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	H	39	GLN
2	L	38	GLN
2	L	43	GLN
2	L	199	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 34 ligands modelled in this entry, 7 are monoatomic - leaving 27 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	SO4	H	317	-	4,4,4	0.67	0	6,6,6	0.08	0
3	EDO	H	309	-	3,3,3	0.28	0	2,2,2	0.30	0
3	EDO	H	302	-	3,3,3	0.30	0	2,2,2	0.43	0
4	SO4	H	314	-	4,4,4	0.66	0	6,6,6	0.10	0
3	EDO	H	305	-	3,3,3	0.27	0	2,2,2	0.43	0
4	SO4	L	306	-	4,4,4	0.68	0	6,6,6	0.13	0
3	EDO	L	301	-	3,3,3	0.27	0	2,2,2	0.52	0
3	EDO	H	310	-	3,3,3	0.24	0	2,2,2	0.43	0
3	EDO	H	311	-	3,3,3	0.26	0	2,2,2	0.30	0
4	SO4	H	315	-	4,4,4	0.67	0	6,6,6	0.15	0
3	EDO	L	307	-	3,3,3	0.23	0	2,2,2	0.27	0
3	EDO	H	308	-	3,3,3	0.25	0	2,2,2	0.41	0
4	SO4	H	321	-	4,4,4	0.67	0	6,6,6	0.07	0
3	EDO	L	303	-	3,3,3	0.26	0	2,2,2	0.32	0
3	EDO	H	301	-	3,3,3	0.28	0	2,2,2	0.46	0
3	EDO	L	304	-	3,3,3	0.28	0	2,2,2	0.47	0
4	SO4	H	316	-	4,4,4	0.68	0	6,6,6	0.13	0
4	SO4	L	308	-	4,4,4	0.67	0	6,6,6	0.10	0
3	EDO	H	303	-	3,3,3	0.29	0	2,2,2	0.28	0
3	EDO	H	306	-	3,3,3	0.25	0	2,2,2	0.39	0
3	EDO	L	302	-	3,3,3	0.21	0	2,2,2	0.46	0
4	SO4	H	312	-	4,4,4	0.67	0	6,6,6	0.08	0
4	SO4	H	318	-	4,4,4	0.69	0	6,6,6	0.09	0
3	EDO	L	305	-	3,3,3	0.25	0	2,2,2	0.56	0
4	SO4	H	313	-	4,4,4	0.65	0	6,6,6	0.11	0
3	EDO	H	304	-	3,3,3	0.23	0	2,2,2	0.34	0
3	EDO	H	307	-	3,3,3	0.26	0	2,2,2	0.31	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	EDO	L	301	-	-	0/1/1/1	-
3	EDO	H	310	-	-	0/1/1/1	-
3	EDO	L	303	-	-	1/1/1/1	-
3	EDO	L	305	-	-	0/1/1/1	-
3	EDO	H	301	-	-	0/1/1/1	-
3	EDO	H	311	-	-	1/1/1/1	-
3	EDO	H	309	-	-	0/1/1/1	-
3	EDO	L	302	-	-	0/1/1/1	-
3	EDO	L	307	-	-	0/1/1/1	-
3	EDO	H	304	-	-	0/1/1/1	-
3	EDO	L	304	-	-	0/1/1/1	-
3	EDO	H	308	-	-	0/1/1/1	-
3	EDO	H	302	-	-	1/1/1/1	-
3	EDO	H	303	-	-	0/1/1/1	-
3	EDO	H	306	-	-	0/1/1/1	-
3	EDO	H	305	-	-	1/1/1/1	-
3	EDO	H	307	-	-	0/1/1/1	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (4) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	H	311	EDO	O1-C1-C2-O2
3	H	302	EDO	O1-C1-C2-O2
3	L	303	EDO	O1-C1-C2-O2
3	H	305	EDO	O1-C1-C2-O2

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	H	311	EDO	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	H	216/225 (96%)	0.44	32 (14%) 5 6	19, 39, 103, 128	4 (1%)
2	L	213/213 (100%)	0.64	38 (17%) 4 4	17, 45, 111, 153	1 (0%)
All	All	429/438 (97%)	0.54	70 (16%) 4 5	17, 43, 109, 153	5 (1%)

All (70) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	L	125	LEU	6.1
1	H	133	GLY	5.8
2	L	154	LEU	5.5
2	L	156	SER	5.4
2	L	157	GLY	5.1
2	L	186	TYR	5.0
2	L	193	ALA	4.5
2	L	192	TYR	4.5
1	H	189	LEU	4.2
2	L	126	LYS	4.2
2	L	190	LYS	4.2
1	H	214	LYS	4.2
2	L	214	CYS	3.9
2	L	209	PHE	3.9
1	H	207	VAL	3.9
1	H	191	THR	3.8
2	L	120	PRO	3.7
2	L	184	ALA	3.7
2	L	150	VAL	3.7
2	L	129	THR	3.7
2	L	188	LYS	3.7
1	H	126	PRO	3.5
1	H	202	PRO	3.4
1	H	158	ALA	3.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
2	L	181	LEU	3.2
1	H	213	PRO	3.2
1	H	204	ASN	3.2
1	H	212	GLU	3.2
2	L	213	GLU	3.2
2	L	147	GLN	3.2
2	L	158	ASN	3.1
2	L	205	VAL	3.1
2	L	189	HIS	3.0
1	H	134	GLY	3.0
1	H	190	GLY	3.0
1	H	140	CYS	3.0
2	L	153	ALA	3.0
2	L	155	GLN	2.9
1	H	135	THR	2.8
1	H	203	SER	2.8
1	H	206	LYS	2.8
1	H	194	TYR	2.7
1	H	84[A]	PHE	2.7
2	L	191	VAL	2.7
2	L	152	ASN	2.7
2	L	212	GLY	2.6
1	H	210	LYS	2.6
1	H	187	SER	2.6
2	L	121	SER	2.5
2	L	116	PHE	2.5
1	H	172	SER	2.5
2	L	179	LEU	2.5
1	H	156	SER	2.5
1	H	122	PHE	2.3
2	L	210	ASN	2.3
1	H	121	VAL	2.3
2	L	187	GLU	2.2
2	L	118	PHE	2.2
1	H	211	VAL	2.2
1	H	160	THR	2.2
2	L	194	CYS	2.2
1	H	193	THR	2.2
2	L	145	LYS	2.2
2	L	183	LYS	2.2
2	L	146	VAL	2.1
2	L	127	SER	2.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	H	185	PRO	2.1
1	H	155	ASN	2.1
1	H	125	ALA	2.1
1	H	157	GLY	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
4	SO4	L	306	5/5	0.61	0.11	81,85,89,111	0
4	SO4	H	312	5/5	0.70	0.12	66,76,85,92	0
4	SO4	H	318	5/5	0.76	0.11	49,49,57,66	5
3	EDO	H	306	4/4	0.82	0.17	60,72,80,90	0
4	SO4	H	317	5/5	0.86	0.11	55,67,80,80	0
4	SO4	L	308	5/5	0.86	0.12	66,67,72,73	5
3	EDO	L	303	4/4	0.87	0.15	50,61,78,78	0
3	EDO	L	304	4/4	0.87	0.19	51,61,63,63	0
3	EDO	H	302	4/4	0.87	0.16	45,54,60,60	0
4	SO4	H	315	5/5	0.87	0.12	58,65,72,75	5
3	EDO	H	311	4/4	0.88	0.16	40,53,64,64	0
3	EDO	L	307	4/4	0.88	0.13	49,61,74,83	0
3	EDO	L	302	4/4	0.88	0.17	44,53,62,65	0
3	EDO	H	305	4/4	0.88	0.14	44,53,57,68	0
3	EDO	H	308	4/4	0.89	0.17	47,60,72,73	0
5	CL	L	312	1/1	0.90	0.16	80,80,80,80	0
6	NA	L	313	1/1	0.90	0.26	69,69,69,69	0
3	EDO	L	305	4/4	0.92	0.13	41,49,60,73	0
3	EDO	H	307	4/4	0.92	0.17	45,58,70,70	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	EDO	H	304	4/4	0.92	0.14	45,55,63,64	0
3	EDO	H	309	4/4	0.92	0.13	42,57,69,74	0
3	EDO	H	310	4/4	0.92	0.14	46,56,68,81	0
5	CL	L	309	1/1	0.93	0.19	66,66,66,66	1
3	EDO	L	301	4/4	0.93	0.14	35,42,50,50	0
3	EDO	H	301	4/4	0.93	0.13	42,59,71,71	0
5	CL	L	311	1/1	0.94	0.11	63,63,63,63	0
4	SO4	H	316	5/5	0.94	0.08	51,55,61,71	0
4	SO4	H	314	5/5	0.94	0.10	35,51,61,62	5
4	SO4	H	313	5/5	0.95	0.08	49,59,64,73	0
4	SO4	H	321	5/5	0.95	0.08	62,64,65,66	5
5	CL	L	310	1/1	0.96	0.10	55,55,55,55	1
3	EDO	H	303	4/4	0.97	0.09	34,42,56,56	0
5	CL	H	320	1/1	0.98	0.24	67,67,67,67	1
5	CL	H	319	1/1	0.99	0.07	46,46,46,46	0

6.5 Other polymers [i](#)

There are no such residues in this entry.