



Full wwPDB X-ray Structure Validation Report ⓘ

Aug 31, 2026 – 02:45 PM EDT

PDB ID : 9NIW / pdb_00009niw
Title : Fab1550 in complex with the C-terminal alpha-TSR domain of the *P. falciparum* circumsporozoite protein
Authors : Moskovitz, R.; Wilson, I.A.
Deposited on : 2025-02-26
Resolution : 2.00 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.015 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.50


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	10052 (2.00-2.00)
Clashscore	190562	11152 (2.00-2.00)
Ramachandran outliers	187476	11031 (2.00-2.00)
Sidechain outliers	187428	11029 (2.00-2.00)
RSRZ outliers	180081	10067 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	L	214	94% 6%
2	H	225	96% .
3	C	73	7% 84% 12%
4	A	56	14% 95% 5%

2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 8642 atoms, of which 4105 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Fab1550 light chain.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	L	214	Total	C	H	N	O	S	1	1	0
			3236	1029	1586	274	341	6			

- Molecule 2 is a protein called Fab1550 heavy chain.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
2	H	225	Total	C	H	N	O	S	0	0	0
			3283	1055	1609	282	331	6			

- Molecule 3 is a protein called Circumsporozoite protein.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
3	C	64	Total	C	H	N	O	S	0	0	0
			1004	313	499	87	101	4			

There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
C	377	HIS	-	expression tag	UNP Q7K740
C	378	HIS	-	expression tag	UNP Q7K740
C	379	HIS	-	expression tag	UNP Q7K740
C	380	HIS	-	expression tag	UNP Q7K740
C	381	HIS	-	expression tag	UNP Q7K740
C	382	HIS	-	expression tag	UNP Q7K740

- Molecule 4 is a protein called Immunoglobulin G-binding protein G.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
4	A	56	Total	C	H	N	O		0	0	0
			843	269	411	67	96				

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	29	ALA	VAL	conflict	UNP P19909

- Molecule 5 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	L	109	Total O 109 109	0	0
5	H	147	Total O 147 147	0	0
5	C	11	Total O 11 11	0	0
5	A	9	Total O 9 9	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.



- Molecule 1: Fab1550 light chain



- Molecule 2: Fab1550 heavy chain



- Molecule 3: Circumsporozoite protein



- Molecule 4: Immunoglobulin G-binding protein G



4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	147.33Å 61.01Å 70.45Å 90.00° 106.50° 90.00°	Depositor
Resolution (Å)	37.27 – 2.00 37.27 – 2.00	Depositor EDS
% Data completeness (in resolution range)	90.1 (37.27-2.00) 90.1 (37.27-2.00)	Depositor EDS
R_{merge}	0.13	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	5.14 (at 2.00Å)	Xtriage
Refinement program	PHENIX (1.21.2_5419: ???)	Depositor
R, R_{free}	0.165 , 0.208 0.165 , 0.208	Depositor DCC
R_{free} test set	1861 reflections (4.57%)	wwPDB-VP
Wilson B-factor (Å ²)	21.2	Xtriage
Anisotropy	0.191	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.44 , 45.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	8642	wwPDB-VP
Average B, all atoms (Å ²)	29.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.75% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: PCA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	L	0.41	0/1684	0.48	0/2285
2	H	0.47	0/1710	0.53	0/2330
3	C	0.32	0/513	0.36	0/688
4	A	0.30	0/438	0.33	0/594
All	All	0.42	0/4345	0.47	0/5897

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	L	1650	1586	1595	9	0
2	H	1674	1609	1620	5	0
3	C	505	499	505	2	0
4	A	432	411	411	3	0
5	A	9	0	0	1	0
5	C	11	0	0	0	0
5	H	147	0	0	0	0
5	L	109	0	0	0	0
All	All	4537	4105	4131	17	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (17) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
4:A:36:ASP:OD2	5:A:101:HOH:O	2.06	0.72
2:H:12:VAL:HG11	2:H:82(C):LEU:HD13	1.79	0.65
1:L:31:LYS:HE2	1:L:67:SER:HB3	1.88	0.56
4:A:19:GLU:OE1	4:A:19:GLU:N	2.41	0.53
3:C:320:LEU:HD11	3:C:364:ILE:HG21	1.91	0.52
1:L:80:PRO:O	1:L:83:ILE:HG12	2.11	0.49
2:H:64:LYS:HE3	2:H:65:GLY:H	1.77	0.49
1:L:11:LEU:HD12	1:L:11:LEU:C	2.39	0.47
2:H:178:LEU:C	2:H:178:LEU:HD12	2.39	0.47
1:L:4:MET:HE1	1:L:25:ALA:HB2	1.99	0.44
1:L:83:ILE:HD11	1:L:106:ILE:HG12	2.00	0.43
4:A:22:ASP:N	4:A:22:ASP:OD1	2.49	0.43
2:H:100(B):PRO:HG2	3:C:347:LYS:HE2	2.01	0.42
1:L:12:SER:OG	1:L:107:LYS:HG3	2.21	0.41
1:L:34:ASN:ND2	2:H:100(F):TYR:HB3	2.36	0.40
1:L:83:ILE:HD11	1:L:106:ILE:CG1	2.51	0.40
1:L:119:PRO:HG3	1:L:214:CYS:SG	2.62	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	L	213/214 (100%)	206 (97%)	7 (3%)	0	100	100
2	H	223/225 (99%)	218 (98%)	5 (2%)	0	100	100
3	C	62/73 (85%)	62 (100%)	0	0	100	100
4	A	54/56 (96%)	53 (98%)	1 (2%)	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
All	All	552/568 (97%)	539 (98%)	13 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	L	190/189 (100%)	190 (100%)	0	100	100
2	H	186/186 (100%)	186 (100%)	0	100	100
3	C	59/68 (87%)	59 (100%)	0	100	100
4	A	45/45 (100%)	45 (100%)	0	100	100
All	All	480/488 (98%)	480 (100%)	0	100	100

There are no protein residues with a non-rotameric sidechain to report.

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (9) such sidechains are listed below:

Mol	Chain	Res	Type
1	L	38	GLN
1	L	79	GLN
1	L	93	ASN
1	L	138	ASN
1	L	155	GLN
1	L	158	ASN
2	H	39	GLN
2	H	171	GLN
3	C	321	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

1 non-standard protein/DNA/RNA residue is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	PCA	H	3	2	7,8,9	0.74	0	9,10,12	1.42	1 (11%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	PCA	H	3	2	-	0/0/11/13	0/1/1/1

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
2	H	3	PCA	CB-CA-N	2.22	109.35	103.24

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	L	214/214 (100%)	-0.22	3 (1%) 73 73	13, 28, 43, 59	1 (0%)
2	H	224/225 (99%)	-0.56	0 100 100	14, 21, 35, 48	0
3	C	64/73 (87%)	0.46	5 (7%) 19 18	22, 40, 75, 86	0
4	A	56/56 (100%)	0.71	8 (14%) 6 5	23, 43, 65, 78	0
All	All	558/568 (98%)	-0.19	16 (2%) 53 52	13, 26, 54, 86	1 (0%)

All (16) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	L	214	CYS	5.7
3	C	349	GLY	3.3
4	A	1	THR	3.2
1	L	212	GLY	3.1
3	C	373	LYS	3.0
3	C	354	PRO	2.9
4	A	18	THR	2.8
4	A	17	THR	2.7
3	C	351	ALA	2.4
1	L	213	GLU	2.4
4	A	20	ALA	2.3
4	A	21	VAL	2.3
3	C	356	ASP	2.3
4	A	42	GLU	2.1
4	A	19	GLU	2.1
4	A	2	THR	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum,

median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	PCA	H	3	8/9	0.92	0.09	16,24,34,34	0

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.