



Full wwPDB X-ray Structure Validation Report ⓘ

Aug 31, 2026 – 02:49 PM EDT

PDB ID : 9NJ4 / pdb_00009nj4
Title : Fab1437 in complex with the C-terminal alpha-TSR domain of the *P. falciparum* circumsporozoite protein
Authors : Moskovitz, R.; Wilson, I.A.
Deposited on : 2025-02-26
Resolution : 1.95 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.015 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.50


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.95 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	3494 (1.96-1.96)
Clashscore	190562	3612 (1.96-1.96)
Ramachandran outliers	187476	3587 (1.96-1.96)
Sidechain outliers	187428	3587 (1.96-1.96)
RSRZ outliers	180081	3495 (1.96-1.96)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	H	225	32% 96%
2	L	216	2% 95%
3	C	73	36% 88% 11%

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 7814 atoms, of which 3751 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Fab1437 heavy chain.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	H	225	Total	C	H	N	O	S	0	11	0
			3373	1066	1672	290	338	7			

- Molecule 2 is a protein called Fab1437 light chain.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
2	L	214	Total	C	H	N	O	S	0	3	0
			3142	996	1555	265	321	5			

- Molecule 3 is a protein called Circumsporozoite protein.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
3	C	65	Total	C	H	N	O	S	0	2	0
			1033	319	516	89	104	5			

There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
C	377	HIS	-	expression tag	UNP Q7K740
C	378	HIS	-	expression tag	UNP Q7K740
C	379	HIS	-	expression tag	UNP Q7K740
C	380	HIS	-	expression tag	UNP Q7K740
C	381	HIS	-	expression tag	UNP Q7K740
C	382	HIS	-	expression tag	UNP Q7K740

- Molecule 4 is GLYCEROL (CCD ID: GOL) (formula: C₃H₈O₃).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	L	1	Total	C	H	O	0	0
			14	3	8	3		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	H	85	Total	O	0	0
			85	85		
5	L	153	Total	O	0	0
			153	153		
5	C	14	Total	O	0	0
			14	14		


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Fab1437 heavy chain



- Molecule 2: Fab1437 light chain



- Molecule 3: Circumsporozoite protein



4 Data and refinement statistics

Property	Value	Source
Space group	C 2 2 21	Depositor
Cell constants a, b, c, α , β , γ	71.72Å 106.11Å 152.37Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	46.85 – 1.95 46.85 – 1.95	Depositor EDS
% Data completeness (in resolution range)	99.0 (46.85-1.95) 99.4 (46.85-1.95)	Depositor EDS
R_{merge}	0.14	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.63 (at 1.95Å)	Xtriage
Refinement program	PHENIX (1.21rc1_5127: ???)	Depositor
R, R_{free}	0.235 , 0.262 0.235 , 0.262	Depositor DCC
R_{free} test set	2132 reflections (5.02%)	wwPDB-VP
Wilson B-factor (Å ²)	29.0	Xtriage
Anisotropy	0.430	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.38 , 31.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.47$, $\langle L^2 \rangle = 0.30$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	7814	wwPDB-VP
Average B, all atoms (Å ²)	40.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.00% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GOL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	H	0.11	0/1783	0.31	0/2426
2	L	0.12	0/1641	0.34	0/2240
3	C	0.09	0/530	0.26	0/711
All	All	0.11	0/3954	0.31	0/5377

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	H	1701	1672	1618	8	0
2	L	1587	1555	1533	7	0
3	C	517	516	510	1	0
4	L	6	8	8	0	0
5	C	14	0	0	0	0
5	H	85	0	0	0	0
5	L	153	0	0	1	0
All	All	4063	3751	3669	14	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (14) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:H:6:GLU:OE1	1:H:6:GLU:N	2.38	0.56
1:H:33:ALA:HB3	1:H:99:SER:OG	2.09	0.53
1:H:187:LEU:C	1:H:187:LEU:HD12	2.36	0.50
1:H:108:SER:N	1:H:109:PRO:HD3	2.27	0.49
1:H:47:TRP:CZ3	2:L:98:ALA:HB1	2.48	0.48
2:L:4:LEU:HD11	2:L:91:THR:HG22	1.95	0.48
2:L:31:TYR:HH	3:C:350[B]:SER:HB3	1.83	0.42
2:L:33:TYR:HB3	2:L:51:ASP:HA	2.01	0.41
2:L:51:ASP:O	2:L:52:ASN:HB2	2.20	0.41
2:L:136:LEU:N	2:L:136:LEU:HD12	2.35	0.41
2:L:132:ASN:OD1	5:L:401:HOH:O	2.22	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	H	234/225 (104%)	227 (97%)	7 (3%)	0	100	100
2	L	215/216 (100%)	210 (98%)	5 (2%)	0	100	100
3	C	65/73 (89%)	65 (100%)	0	0	100	100
All	All	514/514 (100%)	502 (98%)	12 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	H	197/186 (106%)	197 (100%)	0	100	100
2	L	180/179 (101%)	180 (100%)	0	100	100
3	C	61/68 (90%)	61 (100%)	0	100	100
All	All	438/433 (101%)	438 (100%)	0	100	100

There are no protein residues with a non-rotameric sidechain to report.

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
3	C	340	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry ⓘ

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	GOL	L	301	-	5,5,5	0.32	0	5,5,5	0.50	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	GOL	L	301	-	-	4/4/4/4	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (4) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	L	301	GOL	O1-C1-C2-O2
4	L	301	GOL	O1-C1-C2-C3
4	L	301	GOL	C1-C2-C3-O3
4	L	301	GOL	O2-C2-C3-O3

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	H	225/225 (100%)	1.47	73 (32%) 1 0	12, 44, 61, 75	5 (2%)
2	L	214/216 (99%)	0.22	4 (1%) 66 74	12, 30, 44, 68	1 (0%)
3	C	65/73 (89%)	1.71	26 (40%) 1 0	17, 54, 78, 101	1 (1%)
All	All	504/514 (98%)	0.97	103 (20%) 3 3	12, 38, 62, 101	7 (1%)

All (103) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
3	C	374	CYS	6.3
1	H	120	VAL	5.0
1	H	137	SER	4.7
1	H	225	CYS	4.6
1	H	22	CYS	4.5
2	L	215	CYS	4.5
1	H	119	ILE	4.3
1	H	79	LEU	4.2
3	C	338	CYS	4.1
1	H	82[A]	GLN	4.1
1	H	68	PHE	4.0
1	H	48	VAL	3.9
1	H	55	GLY	3.8
1	H	36	TRP	3.7
3	C	369	CYS	3.6
1	H	24	ALA	3.5
1	H	88	ALA	3.5
3	C	316	ILE	3.5
1	H	18	LEU	3.4
3	C	323	ILE	3.4
3	C	334	CYS	3.4
1	H	80	TYR	3.3
1	H	75	SER	3.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	H	62	VAL	3.3
1	H	64	VAL	3.3
1	H	138	LYS	3.3
1	H	15	GLY	3.3
1	H	94	TYR	3.3
1	H	224	SER	3.3
1	H	95	TYR	3.3
1	H	20	LEU	3.2
1	H	7	SER	3.2
1	H	21	SER	3.2
1	H	140	THR	3.2
1	H	70	ILE	3.1
1	H	89	ALA	3.1
1	H	11	LEU	3.1
1	H	12	VAL	3.1
3	C	337	THR	3.1
1	H	96	CYS	3.0
3	C	346	ILE	3.0
1	H	14	PRO	2.9
1	H	83[A]	MET	2.9
1	H	30	ARG	2.9
2	L	2	SER	2.9
3	C	336	VAL	2.9
1	H	60	TYR	2.9
1	H	29	PHE	2.9
1	H	87	ARG	2.9
1	H	19	ARG	2.8
1	H	17	SER	2.8
3	C	373	LYS	2.8
1	H	77	ASN	2.8
1	H	84	ASN	2.8
3	C	368	ILE	2.8
3	C	371	MET	2.7
1	H	78	THR	2.7
3	C	311	PRO	2.7
1	H	66	GLY	2.7
1	H	61	ALA	2.6
1	H	51	ILE	2.6
3	C	325	ASN	2.6
1	H	13	GLN	2.6
1	H	102	TYR	2.6
1	H	86	LEU	2.6

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	H	69	THR	2.6
1	H	81	LEU	2.5
1	H	5	VAL	2.5
2	L	3	VAL	2.5
1	H	72	ARG	2.5
1	H	73	ASP	2.5
3	C	313	ASP	2.5
1	H	76	GLU	2.5
1	H	118	VAL	2.5
3	C	319	TYR	2.4
1	H	4	LEU	2.4
3	C	370	LYS	2.4
1	H	31	HIS	2.4
2	L	197	CYS	2.4
1	H	28	THR	2.4
1	H	10	GLY	2.4
3	C	320	LEU	2.3
1	H	8	GLY	2.3
1	H	136	SER	2.3
1	H	74	ASN	2.3
1	H	121	SER	2.3
1	H	2	VAL	2.3
3	C	372	GLU	2.3
1	H	71	SER	2.2
1	H	104	GLY	2.2
1	H	139	SER	2.2
1	H	65	LYS	2.2
1	H	117	LEU	2.2
3	C	333	PRO	2.2
3	C	312	SER	2.1
3	C	321	ASN	2.1
3	C	364	ILE	2.1
1	H	85	SER	2.1
1	H	91	THR	2.1
3	C	339	GLY	2.1
3	C	340	ASN	2.1
3	C	322	LYS	2.0
1	H	47	TRP	2.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
4	GOL	L	301	6/6	0.91	0.13	28,33,36,42	14

6.5 Other polymers [i](#)

There are no such residues in this entry.