



# Full wwPDB X-ray Structure Validation Report ⓘ

Apr 6, 2026 – 12:04 PM EDT

PDB ID : 9NZD / pdb\_00009nzd  
Title : Crystal structure of AfNth1-K122A mutant bound to Tg-DNA duplex complex in an intermediate state  
Authors : Hitomi, K.; Arvai, A.S.; Syed, A.; Parikh, S.; Tainer, J.A.  
Deposited on : 2025-03-31  
Resolution : 3.10 Å (reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0  
Mogul : 2022.3.0, CSD as543be (2022)  
Xtriage (Phenix) : 2.0  
EDS : 3.0  
Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)  
CCP4 : 9.0.010 (Gargrove)  
Density-Fitness : 1.0.12  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.48.1


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*



The reported resolution of this entry is 3.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.




Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	164625	1351 (3.10-3.10)
Clashscore	180529	1454 (3.10-3.10)
Ramachandran outliers	177936	1391 (3.10-3.10)
Sidechain outliers	177891	1391 (3.10-3.10)
RSRZ outliers	164620	1351 (3.10-3.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	209	 75% 22% ..
1	D	209	 75% 22% ..
2	B	13	 62% 38%
2	E	13	 54% 46%
3	C	13	 54% 38% 8%

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Length	Quality of chain
3	F	13	 54% 38% 8%

## 2 Entry composition i

There are 5 unique types of molecules in this entry. The entry contains 4342 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Endonuclease III.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	206	Total 1631	C 1046	N 287	O 292	S 6	0	0	0
1	D	206	Total 1622	C 1039	N 283	O 294	S 6	0	0	0

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	122	ALA	LYS	engineered mutation	UNP O28581
D	122	ALA	LYS	engineered mutation	UNP O28581

- Molecule 2 is a DNA chain called DNA (5'-D(\*TP\*TP\*GP\*GP\*CP\*GP\*TP\*AP\*TP\*CP\*GP\*CP\*G)-3').

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	P			
2	B	13	Total 265	C 127	N 47	O 79	P 12	0	0	0
2	E	13	Total 265	C 127	N 47	O 79	P 12	0	0	0

- Molecule 3 is a DNA chain called DNA (5'-D(\*AP\*CP\*GP\*CP\*GP\*AP\*(CTG)P\*AP\*CP\*GP\*CP\*CP\*A)-3').

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	P			
3	C	13	Total 264	C 125	N 52	O 75	P 12	0	0	0
3	F	13	Total 264	C 125	N 52	O 75	P 12	0	0	0

- Molecule 4 is IRON/SULFUR CLUSTER (CCD ID: SF4) (formula: Fe<sub>4</sub>S<sub>4</sub>).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	1	Total Fe S 8 4 4	0	0
4	D	1	Total Fe S 8 4 4	0	0


- Molecule 5 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	5	Total O 5 5	0	0
5	C	1	Total O 1 1	0	0
5	D	5	Total O 5 5	0	0
5	E	2	Total O 2 2	0	0
5	F	2	Total O 2 2	0	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Endonuclease III



- Molecule 1: Endonuclease III



- Molecule 2: DNA (5'-D(\*TP\*TP\*GP\*GP\*CP\*GP\*TP\*AP\*TP\*CP\*GP\*CP\*G)-3')



- Molecule 2: DNA (5'-D(\*TP\*TP\*GP\*GP\*CP\*GP\*TP\*AP\*TP\*CP\*GP\*CP\*G)-3')




- Molecule 3: DNA (5'-D(\*AP\*CP\*GP\*CP\*GP\*AP\*(CTG)P\*AP\*CP\*GP\*CP\*CP\*A)-3')





- Molecule 3: DNA (5'-D(\*AP\*CP\*GP\*CP\*GP\*AP\*(CTG)P\*AP\*CP\*GP\*CP\*CP\*A)-3')

Chain F: 54% 38% 8%



## 4 Data and refinement statistics i

Property	Value	Source
Space group	P 41 21 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	110.25Å 110.25Å 112.87Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	38.98 – 3.10 38.98 – 3.10	Depositor EDS
% Data completeness (in resolution range)	91.6 (38.98-3.10) 91.6 (38.98-3.10)	Depositor EDS
$R_{merge}$	0.07	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	0.88 (at 3.12Å)	Xtrriage
Refinement program	PHENIX (1.21_5207: ???)	Depositor
R, $R_{free}$	0.214 , 0.265 0.214 , 0.268	Depositor DCC
$R_{free}$ test set	671 reflections (5.10%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	100.3	Xtrriage
Anisotropy	0.021	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.27 , 55.7	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.48$ , $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	0.021 for -h,l,k 0.018 for -l,-k,-h	Xtrriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	4342	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	114.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.18% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality i

### 5.1 Standard geometry i

Bond lengths and bond angles in the following residue types are not validated in this section: SF4, CTG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.15	0/1663	0.39	1/2247 (0.0%)
1	D	0.14	0/1654	0.34	0/2236
2	B	0.23	0/296	0.48	0/456
2	E	0.22	0/296	0.50	0/456
3	C	0.22	0/271	0.41	0/413
3	F	0.24	0/271	0.41	0/413
All	All	0.17	0/4451	0.39	1/6221 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed( $^{\circ}$ )	Ideal( $^{\circ}$ )
1	A	200	ILE	N-CA-C	6.01	116.21	110.74

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	199	PRO	Peptide

## 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1631	0	1696	31	0
1	D	1622	0	1669	34	0
2	B	265	0	149	3	0
2	E	265	0	149	4	0
3	C	264	0	148	5	0
3	F	264	0	148	7	0
4	A	8	0	0	0	0
4	D	8	0	0	0	0
5	A	5	0	0	1	0
5	C	1	0	0	0	0
5	D	5	0	0	0	0
5	E	2	0	0	0	0
5	F	2	0	0	0	0
All	All	4342	0	3959	78	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 10.

All (78) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:C:321:CTG:C4'	3:C:321:CTG:O4'	1.68	1.19
3:F:321:CTG:C4'	3:F:321:CTG:O4'	1.68	1.12
1:A:9:VAL:HG11	1:A:199:PRO:HG2	1.62	0.82
1:A:149:LEU:HA	1:A:203:CYS:HB3	1.68	0.75
1:A:204:PRO:O	1:A:205:ARG:NH1	2.23	0.72
1:A:186:THR:HG22	1:A:187:VAL:HG23	1.73	0.71
1:D:186:THR:HG22	1:D:187:VAL:HG23	1.71	0.70
1:D:24:LEU:HB3	1:D:50:VAL:HG21	1.75	0.67
1:A:161:GLU:HG2	1:A:165:LYS:HE2	1.76	0.67
1:D:137:PRO:O	1:D:177:ASN:ND2	2.27	0.67
1:D:148:ARG:NH1	1:D:188:CYS:SG	2.69	0.64
1:D:65:ASP:HA	1:D:68:LYS:HE2	1.79	0.63
1:D:31:PRO:HB2	1:D:106:VAL:HG21	1.83	0.61
1:A:102:TYR:OH	1:A:112:GLU:OE1	2.18	0.60

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:162:GLU:HB3	1:D:166:ARG:HH12	1.68	0.59
1:D:40:LEU:O	1:D:87:LYS:NZ	2.35	0.58
1:A:116:LEU:O	5:A:401:HOH:O	2.17	0.56
1:D:2:ASP:HB3	1:D:5:GLU:HB3	1.88	0.56
2:E:301:DT:H2''	2:E:302:DT:H5'	1.89	0.55
1:D:21:VAL:HG11	1:D:181:VAL:HG23	1.89	0.54
1:D:162:GLU:O	1:D:166:ARG:HG2	2.08	0.54
3:C:315:DA:H4'	3:C:316:DC:H5'	1.89	0.53
3:C:322:DA:H2''	3:C:323:DC:O5'	2.08	0.53
1:D:120:GLY:N	3:F:323:DC:H5''	2.23	0.53
1:D:120:GLY:H	3:F:323:DC:H5''	1.74	0.53
1:D:62:LYS:HE2	1:D:63:PRO:HD2	1.91	0.53
1:A:154:THR:OG1	1:A:160:THR:OG1	2.17	0.53
1:D:94:LEU:HD21	1:D:116:LEU:HD13	1.91	0.53
1:A:20:PRO:HA	1:A:23:HIS:HB3	1.91	0.53
1:D:24:LEU:HD12	1:D:46:ASP:HB3	1.90	0.52
1:D:74:ILE:O	1:D:78:ILE:HG12	2.09	0.52
1:A:132:ASP:HA	1:A:174:GLU:HG3	1.92	0.52
1:D:78:ILE:O	1:D:81:VAL:HG22	2.10	0.51
1:A:142:VAL:HA	1:A:180:MET:HE2	1.92	0.51
3:F:318:DC:H2''	3:F:319:DG:C8	2.46	0.51
1:A:162:GLU:O	1:A:166:ARG:HG3	2.11	0.51
1:D:137:PRO:HG2	1:D:177:ASN:CG	2.37	0.50
2:B:302:DT:H2''	2:B:303:DG:C8	2.47	0.49
1:A:108:LEU:HD12	1:D:29:LYS:HB3	1.95	0.48
2:E:303:DG:H2''	2:E:304:DG:C8	2.47	0.48
1:D:143:HIS:HA	1:D:160:THR:HG21	1.96	0.48
1:A:27:GLU:O	1:A:33:GLN:NE2	2.47	0.47
1:D:13:GLU:OE2	1:D:17:ARG:NE	2.48	0.47
3:C:323:DC:H1'	3:C:324:DG:H5''	1.97	0.46
1:A:69:LEU:HB3	1:A:73:GLU:HG3	1.97	0.46
2:B:310:DC:H2'	2:B:311:DG:C8	2.50	0.46
3:F:322:DA:H2''	3:F:323:DC:O5'	2.15	0.46
1:A:29:LYS:HE2	1:D:108:LEU:HB2	1.97	0.46
1:A:148:ARG:NH1	1:A:188:CYS:SG	2.79	0.46
2:E:305:DC:H2''	2:E:306:DG:C8	2.51	0.46
1:A:91:LEU:HD13	1:A:91:LEU:HA	1.84	0.45
1:A:145:ILE:O	1:A:149:LEU:HG	2.18	0.44
1:A:148:ARG:HG2	1:A:205:ARG:HG3	1.99	0.44
1:A:205:ARG:HD3	1:A:205:ARG:HA	1.63	0.44
3:F:319:DG:H2''	3:F:320:DA:C8	2.52	0.44

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:66:LEU:HD23	1:A:95:ALA:HB2	2.00	0.44
1:A:25:LYS:HD3	1:A:25:LYS:HA	1.72	0.43
1:D:94:LEU:HD23	1:D:116:LEU:HB3	2.00	0.43
3:F:321:CTG:H2'	3:F:321:CTG:H6	1.73	0.43
1:D:143:HIS:HD2	1:D:157:PRO:HA	1.83	0.43
1:D:44:THR:HG21	1:D:81:VAL:HA	2.01	0.42
2:E:302:DT:H2''	2:E:303:DG:C8	2.54	0.42
1:D:66:LEU:HD21	1:D:91:LEU:HD12	2.02	0.42
1:D:145:ILE:O	1:D:149:LEU:HG	2.19	0.42
1:D:141:HIS:O	1:D:145:ILE:HG12	2.20	0.41
1:D:144:ARG:HD3	1:D:148:ARG:NH2	2.35	0.41
1:A:175:LYS:HZ1	1:D:175:LYS:HD2	1.86	0.41
1:A:201:LYS:HE2	1:A:201:LYS:HB3	1.65	0.41
2:B:306:DG:O6	3:C:322:DA:N6	2.53	0.41
1:D:62:LYS:CE	1:D:63:PRO:HD2	2.50	0.41
1:A:73:GLU:H	1:A:73:GLU:HG2	1.73	0.41
1:A:35:LEU:HD11	1:A:94:LEU:HD11	2.02	0.41
1:A:66:LEU:HD21	1:A:91:LEU:HD12	2.02	0.41
1:D:60:VAL:HG11	1:D:66:LEU:HD13	2.02	0.41
1:A:13:GLU:OE2	1:A:17:ARG:NH1	2.54	0.40
1:A:67:LEU:HD11	1:A:96:LYS:HG2	2.03	0.40
1:A:4:ILE:HD11	1:A:169:PRO:HG2	2.03	0.40
1:D:39:LEU:HD22	1:D:91:LEU:HD22	2.03	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	204/209 (98%)	187 (92%)	16 (8%)	1 (0%)	25 58
1	D	204/209 (98%)	189 (93%)	15 (7%)	0	100 100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
All	All	408/418 (98%)	376 (92%)	31 (8%)	1 (0%)	44 74

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	199	PRO

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	176/181 (97%)	171 (97%)	5 (3%)	38 66
1	D	174/181 (96%)	168 (97%)	6 (3%)	32 62
All	All	350/362 (97%)	339 (97%)	11 (3%)	35 63

All (11) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	36	VAL
1	A	67	LEU
1	A	86	VAL
1	A	91	LEU
1	A	203	CYS
1	D	5	GLU
1	D	28	ILE
1	D	36	VAL
1	D	86	VAL
1	D	91	LEU
1	D	94	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	23	HIS

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
3	CTG	C	321	3	18,23,24	4.98	10 (55%)	21,35,38	2.26	6 (28%)
3	CTG	F	321	3	18,23,24	5.00	12 (66%)	21,35,38	2.13	5 (23%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	CTG	C	321	3	-	6/7/45/46	0/2/2/2
3	CTG	F	321	3	-	4/7/45/46	0/2/2/2

All (22) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	C	321	CTG	O4'-C4'	10.51	1.68	1.45
3	F	321	CTG	O4'-C4'	10.40	1.68	1.45
3	F	321	CTG	C4-N3	8.94	1.51	1.37
3	C	321	CTG	C4-N3	8.83	1.51	1.37
3	F	321	CTG	C3'-C4'	-8.82	1.29	1.53
3	C	321	CTG	C3'-C4'	-8.69	1.30	1.53
3	C	321	CTG	C2-N3	7.71	1.51	1.38
3	F	321	CTG	C2-N3	7.62	1.51	1.38
3	F	321	CTG	C2-N1	6.84	1.49	1.36
3	C	321	CTG	C2-N1	6.80	1.49	1.36

*Continued on next page...*

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	C	321	CTG	O4'-C1'	-5.64	1.29	1.42
3	F	321	CTG	O4'-C1'	-5.56	1.30	1.42
3	C	321	CTG	O3'-C3'	3.17	1.50	1.43
3	F	321	CTG	O3'-C3'	3.11	1.49	1.43
3	F	321	CTG	C1'-N1	2.72	1.49	1.45
3	F	321	CTG	O5-C5	2.72	1.47	1.43
3	C	321	CTG	O5-C5	2.58	1.47	1.43
3	C	321	CTG	C1'-N1	2.57	1.49	1.45
3	F	321	CTG	C5-C4	2.44	1.55	1.52
3	F	321	CTG	O6-C6	2.25	1.45	1.40
3	C	321	CTG	O6-C6	2.21	1.45	1.40
3	F	321	CTG	C2'-C1'	2.02	1.57	1.52

All (11) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	F	321	CTG	N3-C2-N1	-5.47	111.06	116.78
3	C	321	CTG	N3-C2-N1	-5.44	111.09	116.78
3	C	321	CTG	C5-C4-N3	-4.49	109.33	115.23
3	F	321	CTG	C5-C4-N3	-4.49	109.34	115.23
3	C	321	CTG	C4-N3-C2	-4.37	120.08	126.67
3	F	321	CTG	C4-N3-C2	-4.34	120.12	126.67
3	F	321	CTG	C2'-C1'-N1	-3.02	111.50	115.59
3	C	321	CTG	C2'-C1'-N1	-2.49	112.22	115.59
3	C	321	CTG	O4-C4-N3	2.48	124.61	120.59
3	C	321	CTG	C3'-C2'-C1'	-2.25	97.09	102.60
3	F	321	CTG	C5M-C5-C4	-2.09	105.74	108.72

There are no chirality outliers.

All (10) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	F	321	CTG	O4'-C1'-N1-C2
3	C	321	CTG	C3'-C4'-C5'-O5'
3	C	321	CTG	O4'-C4'-C5'-O5'
3	F	321	CTG	C2'-C1'-N1-C6
3	F	321	CTG	O4'-C1'-N1-C6
3	C	321	CTG	C2'-C1'-N1-C2
3	C	321	CTG	C2'-C1'-N1-C6
3	F	321	CTG	C2'-C1'-N1-C2
3	C	321	CTG	O4'-C1'-N1-C6
3	C	321	CTG	O4'-C1'-N1-C2

There are no ring outliers.

2 monomers are involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	C	321	CTG	1	0
3	F	321	CTG	2	0

## 5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
4	SF4	D	300	1	0,12,12	-	-	-		
4	SF4	A	300	1	0,12,12	-	-	-		

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	SF4	D	300	1	-	-	0/6/5/5
4	SF4	A	300	1	-	-	0/6/5/5

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	206/209 (98%)	-0.52	0 100 100	74, 104, 150, 178	0
1	D	206/209 (98%)	-0.47	0 100 100	69, 123, 171, 204	0
2	B	13/13 (100%)	-0.88	0 100 100	99, 121, 127, 133	0
2	E	13/13 (100%)	-1.10	0 100 100	91, 103, 128, 130	0
3	C	12/13 (92%)	-0.98	0 100 100	87, 110, 130, 147	0
3	F	12/13 (92%)	-1.03	0 100 100	76, 87, 134, 135	0
All	All	462/470 (98%)	-0.55	0 100 100	69, 112, 160, 204	0

There are no RSRZ outliers to report.

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
3	CTG	F	321	22/23	0.93	0.11	76,111,156,250	0
3	CTG	C	321	22/23	0.94	0.13	76,111,156,250	0

### 6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
4	SF4	D	300	8/8	0.95	0.05	176,192,202,218	0
4	SF4	A	300	8/8	0.97	0.05	119,131,160,169	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.