

Full wwPDB X-ray Structure Validation Report (i)

Dec 1, 2025 – 12:08 PM EST

PDB ID : 9O4X / pdb 00009o4x

Title : Gamma delta T cell receptor bound to CD1d

Authors: Rice, M.T.; Gully, B.S.; Rossjohn, J.

Deposited on : 2025-04-09

Resolution : 2.86 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
https://www.wwpdb.org/validation/2017/XrayValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0

Mogul : 2022.3.0, CSD as543be (2022)

Xtriage (Phenix) : 2.0 EDS : 3.0

buster-report : 1.1.7 (2018)

Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)

CCP4 : 9.0.010 (Gargrove)

Density-Fitness : 1.0.12

Ideal geometry (proteins) : Engh & Huber (2001) Ideal geometry (DNA, RNA) : Parkinson et al. (1996)


Validation Pipeline (wwPDB-VP) : 2.46

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: X-RAY DIFFRACTION

The reported resolution of this entry is 2.86 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive $(\# \mathrm{Entries})$	$\begin{array}{c} {\rm Similar\ resolution} \\ (\#{\rm Entries},{\rm resolution\ range}({\rm \AA})) \end{array}$
R_{free}	164625	1268 (2.88-2.84)
Clashscore	180529	1351 (2.88-2.84)
Ramachandran outliers	177936	1318 (2.88-2.84)
Sidechain outliers	177891	1319 (2.88-2.84)
RSRZ outliers	164620	1269 (2.88-2.84)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain	
1	В	99	70%	29%
2	D	197	78%	22%
3	G	224	85%	15%
4	A	274	78%	22%
5	С	2	100%	



Mol	Chain	Length	Quality of chain
5	E	2	100%



2 Entry composition (i)

There are 8 unique types of molecules in this entry. The entry contains 6542 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called Beta-2-microglobulin.

\mathbf{Mol}	Chain	Residues		At	oms			ZeroOcc	AltConf	Trace
1	В	99	Total 820	C 524	N 140	O 153	S 3	0	0	0

• Molecule 2 is a protein called T cell receptor delta chain.

\mathbf{Mol}	Chain	Residues		\mathbf{At}	oms			ZeroOcc	AltConf	Trace	
2	D	197	Total 1562	C 1001	N 264	O 290	S 7	1	0	0	

• Molecule 3 is a protein called T cell receptor gamma chain.

M	ol	Chain	Residues		At	oms			ZeroOcc	AltConf	Trace
3		G	224	Total 1830	C 1171	N 308	O 345	S 6	2	2	0

• Molecule 4 is a protein called Antigen-presenting glycoprotein CD1d.

Mol	Chain	Residues		Ato	oms			ZeroOcc	AltConf	Trace
4	A	274	Total 2188	C 1397	N 382	O 402	S 7	0	0	0

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	279	SER	GLY	conflict	UNP P15813


• Molecule 5 is an oligosaccharide called 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-a cetamido-2-deoxy-beta-D-glucopyranose.





Mol	Chain	Residues	Atoms	ZeroOcc	AltConf	Trace
5	С	2	Total C N O 28 16 2 10	0	0	0
5	E	2	Total C N O 28 16 2 10	0	0	0


 \bullet Molecule 6 is 2-acetamido-2-deoxy-beta-D-glucopyranose (CCD ID: NAG) (formula: $C_8H_{15}NO_6).$



Mol	Chain	Residues	A	tor	ns		ZeroOcc	AltConf
6	D	1	Total 14	C 8	N 1	O 5	0	0


 \bullet Molecule 7 is sphingomyelin (CCD ID: FO4) (formula: $C_{47}H_{94}N_2O_6P)$ (labeled as "Ligand of Interest" by depositor).





Mol	Chain	Residues		Ato	oms			ZeroOcc	AltConf
7	A	1	Total				P	0	0
			56	47	2	6	1		

 \bullet Molecule 8 is PENTAETHYLENE GLYCOL (CCD ID: 1PE) (formula: $\mathrm{C_{10}H_{22}O_6}).$




Mol	Chain	Residues	At	oms		ZeroOcc	AltConf
8	A	1	Total 16	C 10	O 6	0	0



3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.





• Molecule 5: opyranose	$2\hbox{-}acetamido-2\hbox{-}deoxy-beta-D-glucopyranose-(1-4)-2\hbox{-}acetamido-2\hbox{-}deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido$
Chain C:	100%
NAG2	
• Molecule 5: opyranose	$2\hbox{-}acetamido-2\hbox{-}deoxy-beta-D-glucopyranose-} (1\hbox{-}4)-2\hbox{-}acetamido-2\hbox{-}deoxy-beta-D-glucopyranose-} (1\hbox{-}4)-2\hbox{-}acetamido-2-$
Chain E:	100%



4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 43 21 2	Depositor
Cell constants	178.09Å 178.09Å 75.43Å	Donositor
a, b, c, α , β , γ	90.00° 90.00° 90.00°	Depositor
Resolution (Å)	46.65 - 2.86	Depositor
Resolution (A)	46.65 - 2.86	EDS
% Data completeness	99.8 (46.65-2.86)	Depositor
(in resolution range)	95.6 (46.65-2.86)	EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) > 1$	1.63 (at 2.86Å)	Xtriage
Refinement program	PHENIX 1.21.2_5419	Depositor
D D.	0.215 , 0.255	Depositor
R, R_{free}	0.212 , 0.251	DCC
R_{free} test set	2000 reflections (7.03%)	wwPDB-VP
Wilson B-factor (Å ²)	69.7	Xtriage
Anisotropy	0.122	Xtriage
Bulk solvent $k_{sol}(e/Å^3)$, $B_{sol}(Å^2)$	0.32, 61.1	EDS
L-test for twinning ²	$ < L >=0.48, < L^2>=0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	6542	wwPDB-VP
Average B, all atoms (Å ²)	82.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 2.66% of the height of the origin peak. No significant pseudotranslation is detected.

²Theoretical values of <|L|>, $<L^2>$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



¹Intensities estimated from amplitudes.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: FO4, 1PE, NAG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol Chain		Bond	lengths	Bond angles		
Moi Chair	Chain	RMSZ	# Z > 5	RMSZ	# Z > 5	
1	В	0.32	0/843	0.74	0/1142	
2	D	0.11	0/1598	0.33	0/2161	
3	G	0.11	0/1880	0.31	0/2554	
4	A	0.27	0/2253	0.64	1/3069 (0.0%)	
All	All	0.21	0/6574	0.52	1/8926 (0.0%)	

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	\mathbf{Z}	$Observed(^o)$	$\operatorname{Ideal}({}^{o})$
4	A	246	ARG	CG-CD-NE	-7.36	95.81	112.00

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	В	820	0	786	31	0
2	D	1562	0	1570	30	0
3	G	1830	0	1803	22	0
4	A	2188	0	2107	37	0
5	С	28	0	25	1	0
5	Е	28	0	25	0	0



	Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
	6	D	14	0	13	0	0
	7	A	56	0	0	1	0
	8	A	16	0	22	1	0
ĺ	All	All	6542	0	6351	116	0

The all-atom clash score is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clash score for this structure is 9.

All (116) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

A 4 1	A 4 0	Interatomic	Clash
Atom-1	Atom-2	${\rm distance}\ (\rm \mathring{A})$	overlap (Å)
3:G:134:PHE:HB2	3:G:150:LEU:HB3	1.61	0.80
2:D:39:TRP:HB2	2:D:52:ILE:HG22	1.62	0.80
1:B:23:LEU:HB2	1:B:70:PHE:CE2	2.19	0.78
4:A:187:PRO:HB3	4:A:211:PHE:HB3	1.69	0.74
3:G:22:ALA:HB2	3:G:88:LEU:HD11	1.71	0.73
4:A:130:SER:HB2	4:A:149:ASN:HD21	1.57	0.69
1:B:4:THR:HA	1:B:86:THR:HG21	1.72	0.69
1:B:5:PRO:HB3	1:B:30:PHE:HB3	1.74	0.68
4:A:122:ASP:HB3	4:A:134:THR:HG21	1.78	0.66
4:A:133:PRO:HG3	4:A:145:ILE:HG12	1.78	0.65
4:A:256:ALA:HA	4:A:259:LEU:HD12	1.79	0.64
4:A:217:TRP:HB3	4:A:264:LYS:HB2	1.81	0.63
4:A:221:MET:HA	4:A:227:GLN:HG2	1.80	0.62
1:B:23:LEU:HB2	1:B:70:PHE:HE2	1.64	0.62
1:B:7:ILE:HD11	1:B:93:VAL:CB	2.30	0.61
3:G:28:LEU:HD21	3:G:100:THR:HG22	1.82	0.61
4:A:135:GLN:HG3	4:A:136:GLU:HG3	1.82	0.60
4:A:185:VAL:HG21	8:A:302:1PE:H152	1.82	0.60
2:D:89:SER:HA	2:D:111:VAL:O	2.01	0.60
4:A:140:TRP:HA	4:A:143:LEU:HD23	1.84	0.59
2:D:41:LYS:HB2	2:D:51:LEU:HD11	1.84	0.59
3:G:201:LEU:HD12	3:G:227:ILE:HD13	1.84	0.58
2:D:37:ILE:HG21	2:D:77:VAL:HG11	1.86	0.56
1:B:16:GLU:HG2	1:B:19:LYS:HE3	1.88	0.55
4:A:197:PRO:HG3	4:A:250:ASP:OD1	2.06	0.55
4:A:14:GLN:HB3	4:A:98:VAL:HB	1.89	0.55
1:B:16:GLU:HG2	1:B:19:LYS:HG2	1.88	0.54
3:G:57:ASN:HB3	3:G:59:LYS:HE3	1.89	0.54
1:B:16:GLU:HG2	1:B:19:LYS:CE	2.38	0.54
3:G:37:HIS:HD2	3:G:49:ARG:HH11	1.54	0.54



 $Continued\ from\ previous\ page...$

Continued from prev		Interatomic	Clash
Atom-1	Atom-2	${\rm distance}(\mathring{\rm A})$	overlap (Å)
1:B:7:ILE:HD11	1:B:93:VAL:HB	1.88	0.53
4:A:130:SER:HB2	4:A:149:ASN:ND2	2.23	0.53
1:B:24:ASN:HB3	1:B:65:LEU:HD21	1.91	0.53
2:D:134:VAL:HB	2:D:179:TYR:CE1	2.44	0.53
3:G:207:CYS:HB3	3:G:222:ILE:HB	1.91	0.53
4:A:144:ALA:O	4:A:148:LEU:HD13	2.09	0.53
3:G:21:SER:HA	3:G:85:LEU:O	2.09	0.52
3:G:133:ILE:HD13	5:C:1:NAG:H4	1.91	0.52
1:B:7:ILE:HD11	1:B:93:VAL:HG21	1.92	0.52
3:G:223:ILE:HD12	3:G:223:ILE:O	2.09	0.51
4:A:141:VAL:O	4:A:145:ILE:HD13	2.11	0.51
2:D:71:LYS:HG3	2:D:74:ALA:HB3	1.93	0.51
3:G:163:ILE:HG12	3:G:209:VAL:HG22	1.93	0.51
1:B:33:SER:HB2	1:B:54:LEU:HD21	1.93	0.50
4:A:19:ALA:O	4:A:23:TRP:HB3	2.12	0.50
1:B:99:MET:HB2	4:A:207:HIS:HE1	1.77	0.50
1:B:70:PHE:CE1	1:B:72:PRO:HD3	2.47	0.50
2:D:53:ARG:HD3	3:G:106:TYR:CE2	2.47	0.50
1:B:54:LEU:HA	1:B:64:LEU:HD13	1.93	0.49
2:D:5:GLN:HA	2:D:29:GLU:O	2.12	0.49
2:D:136:CYS:SG	2:D:148:ILE:HD11	2.52	0.49
4:A:221:MET:HB2	4:A:225:GLN:O	2.11	0.49
4:A:138:PRO:O	4:A:141:VAL:HG22	2.13	0.49
1:B:19:LYS:N	1:B:19:LYS:HD3	2.27	0.49
1:B:23:LEU:HB2	1:B:70:PHE:CD2	2.48	0.49
4:A:64:GLU:OE1	4:A:64:GLU:HA	2.13	0.48
1:B:75:LYS:HD3	1:B:75:LYS:H	1.79	0.48
3:G:167:GLU:HB2	3:G:170:SER:HB3	1.96	0.48
4:A:189:ALA:HA	4:A:207:HIS:O	2.14	0.48
2:D:123:PRO:HG3	2:D:141:PHE:HB3	1.96	0.47
1:B:7:ILE:HD11	1:B:93:VAL:CG2	2.44	0.47
3:G:138:ILE:H	3:G:138:ILE:HD12	1.78	0.47
3:G:133:ILE:HG23	3:G:224:PHE:CE2	2.50	0.47
1:B:23:LEU:O	1:B:67:TYR:HA	2.14	0.47
2:D:36:TYR:HA	2:D:55:GLY:HA2	1.97	0.46
1:B:70:PHE:C	1:B:70:PHE:CD1	2.93	0.46
2:D:71:LYS:HD3	2:D:73:ALA:N	2.31	0.46
4:A:197:PRO:HG2	4:A:201:ARG:HB2	1.97	0.45
2:D:37:ILE:HG13	2:D:77:VAL:HG21	1.97	0.45
2:D:71:LYS:HD3	2:D:73:ALA:H	1.81	0.45
2:D:38:PHE:HD2	2:D:50:PHE:HE1	1.64	0.45



 $Continued\ from\ previous\ page...$

Continued from prev		Interatomic	Clash
Atom-1	Atom-2	${\rm distance} \ (\mathring{\rm A})$	overlap (Å)
4:A:32:LEU:HD22	4:A:173:LEU:HB3	1.99	0.45
3:G:167:GLU:HG3	3:G:174:LEU:HD11	1.98	0.45
4:A:20:ASN:C	4:A:22:SER:H	2.25	0.45
2:D:21:ALA:HA	2:D:81:ILE:O	2.17	0.45
3:G:69:GLY:O	3:G:86:ARG:HD2	2.16	0.45
2:D:53:ARG:HD3	3:G:106:TYR:CD2	2.52	0.44
1:B:38:ASP:CG	1:B:45:ARG:HH11	2.26	0.44
1:B:70:PHE:CD1	1:B:72:PRO:HD3	2.53	0.44
1:B:18:GLY:C	1:B:19:LYS:HD3	2.43	0.44
2:D:20:LYS:HD3	2:D:20:LYS:HA	1.73	0.44
2:D:134:VAL:HB	2:D:179:TYR:HE1	1.82	0.44
3:G:149:TYR:HB3	3:G:224:PHE:CE2	2.53	0.44
4:A:90:LEU:HD11	4:A:140:TRP:CE2	2.53	0.44
4:A:118:PHE:CD1	4:A:119:GLN:HG2	2.53	0.44
3:G:130:LYS:O	3:G:153:LEU:HA	2.18	0.43
1:B:17:ASN:O	1:B:19:LYS:HE2	2.18	0.43
1:B:88:SER:N	1:B:89:GLN:OE1	2.52	0.43
2:D:88:ASP:HB2	2:D:113:VAL:HG21	2.00	0.43
4:A:10:LEU:HB2	4:A:170:VAL:HG22	2.00	0.43
4:A:230:THR:O	4:A:232:PRO:HD3	2.19	0.43
1:B:7:ILE:C	1:B:7:ILE:HD12	2.44	0.42
2:D:59:GLN:HG3	2:D:60:ASN:H	1.84	0.42
2:D:144:LYS:HB3	2:D:171:TYR:CZ	2.54	0.42
1:B:33:SER:HB3	1:B:62:PHE:CZ	2.54	0.42
4:A:265:HIS:H	4:A:268:LEU:HD12	1.84	0.42
4:A:251:VAL:HG22	4:A:255:GLU:HB2	2.02	0.42
4:A:204:LEU:O	4:A:248:THR:HA	2.20	0.42
2:D:28:TYR:CE2	2:D:75:LYS:HD2	2.55	0.41
1:B:89:GLN:HA	1:B:90:PRO:HD3	1.80	0.41
2:D:16:MET:HE3	2:D:20:LYS:HB3	2.01	0.41
4:A:103:GLU:HB3	4:A:111:ASN:HD21	1.85	0.41
1:B:60:TRP:HB3	4:A:97:GLN:OE1	2.21	0.41
2:D:52:ILE:HD12	2:D:52:ILE:HA	1.98	0.41
2:D:39:TRP:CE2	2:D:79:LEU:HB2	2.55	0.41
2:D:123:PRO:HB3	2:D:138:VAL:CG1	2.50	0.41
4:A:12:CYS:HB3	7:A:301:FO4:C28	2.51	0.41
2:D:114:GLU:HA	2:D:115:PRO:HD3	1.96	0.41
4:A:82:LYS:HG2	4:A:92:TYR:OH	2.20	0.41
4:A:143:LEU:HD22	4:A:143:LEU:H	1.86	0.41
4:A:251:VAL:CG2	4:A:255:GLU:HB2	2.51	0.41
1:B:81:ARG:HD3	1:B:92:ILE:CG1	2.51	0.40



Atom-1	Atom-2	$egin{aligned} & ext{Interatomic} \ & ext{distance} \ & ext{(Å)} \end{aligned}$	Clash overlap (Å)
3:G:40:LEU:HD13	3:G:96:TYR:CZ	2.55	0.40
2:D:144:LYS:O	2:D:146:ILE:HG22	2.22	0.40
2:D:26:CYS:HB3	2:D:77:VAL:O	2.22	0.40
3:G:123:LEU:HD13	3:G:123:LEU:HA	1.96	0.40

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Perce	entiles
1	В	97/99~(98%)	96 (99%)	1 (1%)	0	100	100
2	D	$195/197\ (99\%)$	188 (96%)	7 (4%)	0	100	100
3	G	$224/224 \ (100\%)$	218 (97%)	6 (3%)	0	100	100
4	A	$272/274\ (99\%)$	263 (97%)	9 (3%)	0	100	100
All	All	788/794 (99%)	765 (97%)	23 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	В	92/94 (98%)	91 (99%)	1 (1%)	70 85
2	D	176/176 (100%)	175 (99%)	1 (1%)	84 92



Mol	Chain	Analysed	Rotameric	Percentiles		
3	G	206/205 (100%)	206 (100%)	0	100	100
4	A	237/239~(99%)	235 (99%)	2 (1%)	79	90
All	All	711/714 (100%)	707 (99%)	4 (1%)	84	92

All (4) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	В	65	LEU
2	D	110	ARG
4	A	88	LEU
4	A	110	SER

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (12) such sidechains are listed below:

Mol	Chain	Res	Type
1	В	31	HIS
2	D	183	ASN
3	G	166	GLN
3	G	171	ASN
3	G	185	ASN
3	G	205	HIS
4	A	105	HIS
4	A	111	ASN
4	A	115	HIS
4	A	146	GLN
4	A	149	ASN
4	A	150	GLN

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.



5.5 Carbohydrates (i)

4 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
MIOI	Type	Chain	nes		Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
5	NAG	С	1	5,2	14,14,15	0.70	0	17,19,21	0.96	0
5	NAG	С	2	5	14,14,15	0.71	0	17,19,21	0.90	1 (5%)
5	NAG	Е	1	5,4	14,14,15	0.81	0	17,19,21	1.33	3 (17%)
5	NAG	Е	2	5	14,14,15	1.42	3 (21%)	17,19,21	2.43	4 (23%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	NAG	С	1	5,2	-	0/6/23/26	0/1/1/1
5	NAG	С	2	5	-	0/6/23/26	0/1/1/1
5	NAG	Е	1	5,4	-	2/6/23/26	0/1/1/1
5	NAG	Е	2	5	-	3/6/23/26	0/1/1/1

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	\mathbf{Z}	$\operatorname{Observed}(\text{\AA})$	Ideal(A)
5	Ε	2	NAG	C1-C2	-3.16	1.48	1.52
5	Е	2	NAG	C2-N2	-2.53	1.42	1.46
5	Е	2	NAG	C4-C5	2.31	1.57	1.53

All (8) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	\mathbf{Z}	$Observed(^o)$	$\operatorname{Ideal}({}^{o})$
5	Ε	2	NAG	C2-N2-C7	7.62	133.11	122.90
5	Е	2	NAG	C8-C7-N2	-3.74	109.91	116.12
5	Е	1	NAG	O5-C1-C2	-2.94	106.74	111.29
5	Ε	2	NAG	O7-C7-N2	2.84	127.00	121.98



Mol	Chain	Res	Type	Atoms	${f Z}$	$\mathbf{Observed}(^o)$	$\operatorname{Ideal}({}^{o})$
5	Е	1	NAG	C1-O5-C5	2.39	115.39	112.19
5	С	2	NAG	C1-O5-C5	2.21	115.15	112.19
5	Е	2	NAG	O3-C3-C2	-2.07	105.11	109.40
5	Е	1	NAG	O4-C4-C3	-2.06	105.51	110.38

There are no chirality outliers.

All (5) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
5	Е	2	NAG	C1-C2-N2-C7
5	Е	1	NAG	C8-C7-N2-C2
5	Е	1	NAG	O7-C7-N2-C2
5	Е	2	NAG	O7-C7-N2-C2
5	Е	2	NAG	O5-C5-C6-O6


There are no ring outliers.

1 monomer is involved in 1 short contact:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	С	1	NAG	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.









5.6 Ligand geometry (i)

3 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Mol Type Cl		Res	Res Link	Bo	Bond lengths			Bond angles		
MIOI	туре	Chain		LIIIK	Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2	
8	1PE	A	302	-	15,15,15	0.30	0	14,14,14	0.20	0	
7	FO4	A	301	-	54,55,55	0.51	1 (1%)	60,63,63	0.78	3 (5%)	
6	NAG	D	301	2	14,14,15	0.73	0	17,19,21	1.03	2 (11%)	



In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
8	1PE	A	302	-	-	3/13/13/13	-
7	FO4	A	301	-	-	21/60/60/60	-
6	NAG	D	301	2	-	3/6/23/26	0/1/1/1

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	\mathbf{Z}	$\operatorname{Observed}(\text{\AA})$	$Ideal(\AA)$
7	A	301	FO4	O-C	-2.18	1.39	1.43

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	\mathbf{Z}	$Observed(^o)$	$Ideal(^{o})$
7	A	301	FO4	C-C16-N	-2.75	105.00	109.66
6	D	301	NAG	C2-N2-C7	2.58	126.35	122.90
7	A	301	FO4	O-C-C16	2.40	114.18	107.85
6	D	301	NAG	O5-C1-C2	-2.35	107.66	111.29
7	A	301	FO4	C16-N-C17	2.29	127.22	123.40

There are no chirality outliers.

All (27) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
7	A	301	FO4	O-C-C16-N
7	A	301	FO4	C1-C-C16-C41
7	A	301	FO4	O-C-C16-C41
7	A	301	FO4	C17-C18-C19-C20
7	A	301	FO4	C33-C34-C35-C36
6	D	301	NAG	C8-C7-N2-C2
6	D	301	NAG	O7-C7-N2-C2
8	A	302	1PE	ОН4-С13-С23-ОН3
7	A	301	FO4	O-C-C1-C2
7	A	301	FO4	C11-C12-C13-C14
7	A	301	FO4	C28-C29-C30-C31
8	A	302	1PE	OH5-C14-C24-OH4
8	A	302	1PE	C15-C25-OH5-C14
7	A	301	FO4	C35-C36-C37-C38
7	A	301	FO4	C42-O4-P-O5



Continued from previous page...

Mol	Chain	Res	Type	Atoms
7	A	301	FO4	C16-C-C1-C2
7	A	301	FO4	C3-C4-C5-C6
7	A	301	FO4	C37-C38-C39-C40
7	A	301	FO4	C10-C11-C12-C13
7	A	301	FO4	C42-C43-N1-C44
6	D	301	NAG	O5-C5-C6-O6
7	A	301	FO4	C36-C37-C38-C39
7	A	301	FO4	C32-C33-C34-C35
7	A	301	FO4	C24-C25-C26-C27
7	A	301	FO4	C11-C10-C9-C8
7	A	301	FO4	C31-C32-C33-C34
7	A	301	FO4	C6-C7-C8-C9


There are no ring outliers.

2 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
8	A	302	1PE	1	0
7	A	301	FO4	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ>2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95^{th} percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<rsrz></rsrz>	# RSRZ > 2	$\mathbf{OWAB}(\mathbf{\mathring{A}}^2)$	Q < 0.9
1	В	99/99 (100%)	0.20	2 (2%) 64 61	59, 90, 123, 169	0
2	D	197/197 (100%)	0.06	1 (0%) 87 86	48, 87, 113, 125	2 (1%)
3	G	224/224 (100%)	-0.23	0 100 100	24, 64, 101, 130	4 (1%)
4	A	274/274 (100%)	-0.00	2 (0%) 84 83	48, 80, 143, 178	0
All	All	794/794 (100%)	-0.02	5 (0%) 85 84	24, 79, 125, 178	6 (0%)

All (5) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	В	1	ILE	2.8
1	В	99	MET	2.6
4	A	102	CYS	2.3
4	A	19	ALA	2.2
2	D	136	CYS	2.0

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)


In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mo	l Type	Chain	Res	Atoms	RSCC	RSR	$\operatorname{B-factors}(\mathrm{\AA}^2)$	Q<0.9
5	NAG	С	1	14/15	-	-	82,87,96,109	0
5	NAG	С	2	14/15	-	_	91,104,115,129	0




Mol	Type	Chain	Res	Atoms	RSCC	RSR	$\mathbf{B} ext{-}\mathbf{factors}(\mathbf{\mathring{A}}^2)$	Q < 0.9
5	NAG	Е	2	14/15	0.80	0.14	129,143,158,162	0
5	NAG	Е	1	14/15	0.88	0.12	80,94,107,115	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.








6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	$\operatorname{B-factors}(\mathrm{A}^2)$	Q<0.9
6	NAG	D	301	14/15	0.53	0.17	125,134,140,143	0
7	FO4	A	301	56/56	0.86	0.20	69,93,156,187	0
8	1PE	A	302	16/16	0.92	0.10	59,73,82,83	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.





6.5 Other polymers (i)

There are no such residues in this entry.

