



Full wwPDB X-ray Structure Validation Report ⓘ

Oct 22, 2026 – 12:06 PM EDT

PDB ID : 9PWI / pdb_00009pwi
Title : EGFR kinase bound to ZNL-1
Authors : Beyett, T.S.; Eck, M.J.
Deposited on : 2025-08-04
Resolution : 2.60 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.2
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.2
EDS : 3.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.015 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.51


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION


The reported resolution of this entry is 2.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	4008 (2.60-2.60)
Clashscore	190562	4347 (2.60-2.60)
Ramachandran outliers	187476	4277 (2.60-2.60)
Sidechain outliers	187428	4277 (2.60-2.60)
RSRZ outliers	180081	4008 (2.60-2.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	331	

2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 2243 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Epidermal growth factor receptor.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	274	2202	1422	373	392	15	0	1	0

There are 3 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	692	GLY	-	expression tag	UNP P00533
A	693	SER	-	expression tag	UNP P00533
A	694	THR	-	expression tag	UNP P00533

- Molecule 2 is N-[2-({1-methyl-7-[4-(4-methylpiperazin-1-yl)anilino]-2-oxo-1,4-dihydropyrimido[4,5-d]pyrimidin-3(2H)-yl}methyl)phenyl]prop-2-enamide (CCD ID: A1CMF) (formula: C₂₈H₃₂N₈O₂) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	N	O		
2	A	1	38	28	8	2	0	0


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	3	Total	O	0	0
			3	3		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Epidermal growth factor receptor



4 Data and refinement statistics

Property	Value	Source
Space group	P 43 21 2	Depositor
Cell constants a, b, c, α , β , γ	70.39Å 70.39Å 161.72Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	64.54 – 2.60 64.54 – 2.60	Depositor EDS
% Data completeness (in resolution range)	99.5 (64.54-2.60) 99.5 (64.54-2.60)	Depositor EDS
R_{merge}	0.09	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.26 (at 2.61Å)	Xtriage
Refinement program	PHENIX (1.21_5207: ???)	Depositor
R, R_{free}	0.224 , 0.254 0.233 , 0.265	Depositor DCC
R_{free} test set	607 reflections (4.61%)	wwPDB-VP
Wilson B-factor (Å ²)	71.2	Xtriage
Anisotropy	0.455	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 55.7	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	2243	wwPDB-VP
Average B, all atoms (Å ²)	84.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.94% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: A1CMF

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.17	0/2253	0.34	0/3050

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2202	0	2268	25	0
2	A	38	0	0	1	0
3	A	3	0	0	1	0
All	All	2243	0	2268	25	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (25) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:825:MET:HE2	1:A:853:ILE:HD13	1.68	0.76

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:825:MET:HE3	1:A:838:LEU:HD22	1.72	0.70
1:A:781:CYS:HB2	1:A:787:GLN:HG3	1.74	0.69
1:A:825:MET:HE1	1:A:835:HIS:CD2	2.34	0.62
1:A:892:THR:H	1:A:895:SER:HB3	1.68	0.58
1:A:884:GLU:OE2	1:A:958:ARG:NH2	2.24	0.58
1:A:777:LEU:HD21	1:A:780:ILE:HG22	1.86	0.57
1:A:705:ARG:NH1	1:A:778:LEU:O	2.36	0.57
1:A:750:ALA:HB1	1:A:755:ALA:HB3	1.89	0.54
1:A:733:PRO:HD2	1:A:738:VAL:HG13	1.90	0.53
1:A:797:CYS:SG	3:A:1203:HOH:O	2.59	0.52
1:A:759:ILE:O	1:A:762:GLU:HG2	2.10	0.52
1:A:736:GLU:HG2	1:A:737:LYS:HG2	1.92	0.52
1:A:773:HIS:HA	1:A:852:LYS:HG2	1.94	0.49
1:A:766:MET:SD	1:A:858:LEU:HD21	2.53	0.48
1:A:698:ALA:HB1	1:A:699:PRO:HD2	1.96	0.47
1:A:782:LEU:HD23	1:A:782:LEU:HA	1.77	0.46
1:A:882:ALA:HA	1:A:898:TRP:CD2	2.53	0.44
1:A:706:ILE:HA	1:A:780:ILE:O	2.18	0.44
1:A:731:TRP:O	1:A:733:PRO:HD3	2.17	0.44
1:A:796:GLY:HA2	2:A:1101:A1CMF:C13	2.49	0.42
1:A:731:TRP:CE2	1:A:733:PRO:HG3	2.55	0.41
1:A:716:LYS:HE3	1:A:716:LYS:HB2	1.84	0.41
1:A:842:ASN:O	1:A:854:THR:HG22	2.20	0.41
1:A:806:LYS:HB3	1:A:806:LYS:HE2	1.87	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	271/331 (82%)	255 (94%)	15 (6%)	1 (0%)	30 51

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	786	VAL

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	244/290 (84%)	233 (96%)	11 (4%)	24	50

All (11) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	738	VAL
1	A	780	ILE
1	A	782	LEU
1	A	783	THR
1	A	787	GLN
1	A	831	ARG
1	A	841	ARG
1	A	861	LEU
1	A	884	GLU
1	A	926	ILE
1	A	927	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	893	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	A1CMF	A	1101	1	42,42,42	2.71	18 (42%)	55,59,59	2.98	23 (41%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	A1CMF	A	1101	1	-	4/18/44/44	0/5/5/5

All (18) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	1101	A1CMF	C22-N20	8.96	1.48	1.38
2	A	1101	A1CMF	C22-N23	4.65	1.40	1.36
2	A	1101	A1CMF	C15-N14	4.37	1.45	1.36
2	A	1101	A1CMF	C19-N38	4.19	1.39	1.34
2	A	1101	A1CMF	C19-N20	4.07	1.43	1.38
2	A	1101	A1CMF	C24-N23	3.84	1.52	1.46
2	A	1101	A1CMF	C21-N20	3.71	1.52	1.46
2	A	1101	A1CMF	C08-N07	3.62	1.48	1.38

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	1101	A1CMF	C13-N14	3.33	1.48	1.40
2	A	1101	A1CMF	C15-N16	3.31	1.39	1.34
2	A	1101	A1CMF	C32-N31	3.18	1.41	1.35
2	A	1101	A1CMF	C06-N07	3.10	1.52	1.46
2	A	1101	A1CMF	C30-C25	3.03	1.44	1.40
2	A	1101	A1CMF	C24-C25	2.74	1.56	1.51
2	A	1101	A1CMF	C30-N31	2.61	1.46	1.41
2	A	1101	A1CMF	C17-C18	2.36	1.42	1.37
2	A	1101	A1CMF	C15-N38	2.17	1.40	1.34
2	A	1101	A1CMF	C12-C13	2.02	1.42	1.39

All (23) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	1101	A1CMF	C33-C32-N31	13.90	123.28	113.84
2	A	1101	A1CMF	N16-C15-N38	-6.44	120.19	126.42
2	A	1101	A1CMF	O37-C22-N20	-4.37	118.39	122.10
2	A	1101	A1CMF	O35-C32-C33	-4.32	116.22	122.70
2	A	1101	A1CMF	C12-C13-C10	-4.00	113.74	119.04
2	A	1101	A1CMF	C04-C03-N02	-3.77	104.80	110.86
2	A	1101	A1CMF	C18-C36-N23	3.77	117.26	111.56
2	A	1101	A1CMF	C24-N23-C22	3.58	123.33	118.85
2	A	1101	A1CMF	N23-C22-N20	3.49	121.78	116.75
2	A	1101	A1CMF	C24-C25-C30	3.42	124.21	121.09
2	A	1101	A1CMF	C34-C33-C32	3.34	125.90	122.22
2	A	1101	A1CMF	C11-C08-N07	3.26	125.92	121.39
2	A	1101	A1CMF	N38-C19-N20	2.92	121.18	117.41
2	A	1101	A1CMF	C11-C08-C09	-2.88	113.69	119.18
2	A	1101	A1CMF	C05-C06-N07	2.53	116.11	110.78
2	A	1101	A1CMF	C09-C10-C13	2.52	123.20	120.30
2	A	1101	A1CMF	C11-C12-C13	2.35	123.00	120.30
2	A	1101	A1CMF	C01-N02-C03	-2.33	106.20	110.63
2	A	1101	A1CMF	C12-C11-C08	2.31	123.23	120.30
2	A	1101	A1CMF	C10-C09-C08	2.24	123.14	120.30
2	A	1101	A1CMF	O37-C22-N23	-2.08	119.83	121.98
2	A	1101	A1CMF	C01-N02-C05	-2.07	106.69	110.63
2	A	1101	A1CMF	C21-N20-C19	2.02	121.95	120.09

There are no chirality outliers.

All (4) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	A	1101	A1CMF	N31-C32-C33-C34
2	A	1101	A1CMF	O35-C32-C33-C34
2	A	1101	A1CMF	C11-C08-N07-C04
2	A	1101	A1CMF	C25-C30-N31-C32

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	1101	A1CMF	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	274/331 (82%)	0.47	6 (2%) 62 57	43, 81, 112, 130	1 (0%)

All (6) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	982	GLN	4.5
1	A	762	GLU	4.3
1	A	777	LEU	2.5
1	A	723	PHE	2.1
1	A	785	THR	2.0
1	A	735	GLY	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)


There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q < 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	A1CMF	A	1101	38/38	0.92	0.16	56,79,110,154	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



6.5 Other polymers [i](#)

There are no such residues in this entry.