



Full wwPDB X-ray Structure Validation Report ⓘ

Jan 5, 2026 – 04:12 pm GMT

PDB ID : 9QCJ / pdb_00009qcj
Title : Crystal structure of PhoC wild type acid phosphatase
Authors : Levy, C.W.; Ortmayer, M.
Deposited on : 2025-03-04
Resolution : 2.24 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : ?? (??), CSD ??CSD?? (???)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.47


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.24 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	164625	3139 (2.26-2.22)
Clashscore	180529	3381 (2.26-2.22)
Ramachandran outliers	177936	3334 (2.26-2.22)
Sidechain outliers	177891	3335 (2.26-2.22)
RSRZ outliers	164620	3138 (2.26-2.22)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	249	4% 84% 6% 10%
1	B	249	5% 82% 6% 12%
1	C	249	11% 74% 14% 12%

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 10081 atoms, of which 4914 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Major phosphate-irrepressible acid phosphatase.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	223	Total	C	H	N	O	S	0	0	0
			3352	1066	1650	294	336	6			
1	B	220	Total	C	H	N	O	S	0	1	0
			3314	1055	1629	291	333	6			
1	C	220	Total	C	H	N	O	S	0	0	0
			3305	1053	1625	290	331	6			

There are 6 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	92	ASP	GLY	conflict	UNP P28581
A	171	THR	ILE	conflict	UNP P28581
B	92	ASP	GLY	conflict	UNP P28581
B	171	THR	ILE	conflict	UNP P28581
C	92	ASP	GLY	conflict	UNP P28581
C	171	THR	ILE	conflict	UNP P28581

- Molecule 2 is PHOSPHATE ION (CCD ID: PO4) (formula: O₄P).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	O	P		0	0
			5	4	1			
2	B	1	Total	O	P		0	0
			5	4	1			
2	C	1	Total	O	P		0	0
			5	4	1			

- Molecule 3 is DI(HYDROXYETHYL)ETHER (CCD ID: PEG) (formula: $C_4H_{10}O_3$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	C	1	Total	C	H	O	0	0
			17	4	10	3		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	46	Total 46	O 46	0	0
4	B	15	Total 15	O 15	0	0
4	C	17	Total 17	O 17	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Major phosphate-irrepressible acid phosphatase



- Molecule 1: Major phosphate-irrepressible acid phosphatase



- Molecule 1: Major phosphate-irrepressible acid phosphatase



4 Data and refinement statistics

Property	Value	Source
Space group	P 32 2 1	Depositor
Cell constants a, b, c, α , β , γ	116.38Å 116.38Å 94.54Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	50.39 – 2.24 50.39 – 2.24	Depositor EDS
% Data completeness (in resolution range)	100.0 (50.39-2.24) 93.3 (50.39-2.24)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.64 (at 2.25Å)	Xtriage
Refinement program	PHENIX 1.21.1_5286	Depositor
R, R_{free}	0.205 , 0.236 0.208 , 0.237	Depositor DCC
R_{free} test set	1927 reflections (5.37%)	wwPDB-VP
Wilson B-factor (Å ²)	45.3	Xtriage
Anisotropy	0.230	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.39 , 43.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.34$	Xtriage
Estimated twinning fraction	0.028 for -h,-k,l	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	10081	wwPDB-VP
Average B, all atoms (Å ²)	75.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.21% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: PO4, PEG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.43	0/1737	0.61	0/2357
1	B	0.41	0/1723	0.56	0/2339
1	C	0.43	0/1715	0.57	0/2328
All	All	0.42	0/5175	0.58	0/7024

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1702	1650	1649	9	0
1	B	1685	1629	1628	12	0
1	C	1680	1625	1624	27	0
2	A	5	0	0	0	0
2	B	5	0	0	0	0
2	C	5	0	0	0	0
3	C	7	10	10	0	0
4	A	46	0	0	1	0
4	B	15	0	0	0	0
4	C	17	0	0	0	0
All	All	5167	4914	4911	47	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (47) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:134:GLU:HA	1:C:137:MET:HE3	1.56	0.87
1:B:34:LEU:HD22	1:B:121:ILE:HD11	1.56	0.86
1:C:22:ILE:HG12	1:C:42:ALA:HB3	1.77	0.67
1:C:139:ILE:HD13	1:C:161:ASN:OD1	1.95	0.66
1:C:101:PHE:CE2	1:C:179:LEU:HD22	2.34	0.62
1:C:110:SER:HA	1:C:242:GLU:OE2	2.02	0.59
1:C:117:LEU:HD11	1:C:179:LEU:HD21	1.84	0.59
1:B:117:LEU:HD11	1:B:179:LEU:HD21	1.84	0.58
1:B:139:ILE:N	1:B:139:ILE:HD12	2.18	0.58
1:B:227:HIS:O	1:B:233:GLN:NE2	2.37	0.57
1:B:139:ILE:HD12	1:B:139:ILE:H	1.70	0.56
1:A:90:ALA:HB1	1:A:121:ILE:HG21	1.86	0.56
1:A:134:GLU:HA	1:A:137:MET:HE2	1.86	0.56
1:C:103:TYR:CD2	1:C:246:LYS:HE2	2.41	0.56
1:C:27:ASP:OD2	1:C:38:LYS:HE2	2.09	0.53
1:C:101:PHE:CD2	1:C:179:LEU:HD22	2.44	0.52
1:A:246:LYS:NZ	4:A:403:HOH:O	2.45	0.50
1:C:101:PHE:CD2	1:C:105:ILE:HD11	2.47	0.50
1:C:38:LYS:HD3	1:C:39:ASN:H	1.76	0.49
1:B:22:ILE:HD12	1:B:42:ALA:CB	2.42	0.49
1:A:44:ASP:OD2	1:A:47:LYS:HG3	2.12	0.49
1:C:38:LYS:HD3	1:C:39:ASN:N	2.27	0.49
1:C:114:TYR:O	1:C:118:THR:OG1	2.23	0.47
1:B:59:GLN:HE22	1:C:59:GLN:HE22	1.62	0.47
1:B:81:GLN:NE2	1:B:85:ASP:OD1	2.49	0.46
1:C:101:PHE:HD1	1:C:101:PHE:O	1.99	0.46
1:C:101:PHE:HE2	1:C:179:LEU:HD22	1.77	0.46
1:C:112:GLU:HB2	1:C:242:GLU:OE1	2.15	0.46
1:C:36:TYR:CE2	1:C:119:ASN:HB3	2.51	0.45
1:C:91:GLY:O	1:C:93:VAL:N	2.42	0.45
1:B:169:THR:HG21	1:B:212:VAL:HG13	1.99	0.44
1:C:38:LYS:HD2	1:C:40:GLU:OE1	2.17	0.43
1:C:36:TYR:CZ	1:C:119:ASN:HB3	2.53	0.43
1:B:83[A]:GLN:HG2	1:B:150:CYS:SG	2.59	0.43
1:C:47:LYS:HD3	1:C:47:LYS:HA	1.70	0.42
1:C:81:GLN:NE2	1:C:85:ASP:OD1	2.52	0.42
1:B:34:LEU:CD2	1:B:121:ILE:HD11	2.40	0.42

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:36:TYR:CE2	1:A:119:ASN:HB3	2.55	0.42
1:A:229:ASP:O	1:A:233:GLN:HG3	2.20	0.41
1:A:36:TYR:CZ	1:A:119:ASN:HB3	2.56	0.41
1:A:126:ASP:O	1:A:127:LEU:C	2.62	0.41
1:C:141:PRO:HG3	1:C:208:TRP:CD2	2.55	0.41
1:C:77:GLU:HA	1:C:80:LYS:HE3	2.02	0.41
1:C:22:ILE:N	1:C:22:ILE:HD12	2.35	0.40
1:A:77:GLU:OE2	1:A:80:LYS:NZ	2.54	0.40
1:B:139:ILE:H	1:B:139:ILE:CD1	2.34	0.40
1:C:26:ASN:OD1	1:C:30:THR:HG21	2.21	0.40

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	219/249 (88%)	216 (99%)	3 (1%)	0	100	100
1	B	217/249 (87%)	215 (99%)	2 (1%)	0	100	100
1	C	216/249 (87%)	211 (98%)	4 (2%)	1 (0%)	25	24
All	All	652/747 (87%)	642 (98%)	9 (1%)	1 (0%)	44	49

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	C	90	ALA

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar

resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	175/197 (89%)	175 (100%)	0	100	100
1	B	173/197 (88%)	173 (100%)	0	100	100
1	C	172/197 (87%)	171 (99%)	1 (1%)	84	88
All	All	520/591 (88%)	519 (100%)	1 (0%)	92	95

All (1) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	119	ASN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (9) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	59	GLN
1	A	64	GLN
1	A	168	HIS
1	B	62	ASN
1	B	235	GLN
1	C	59	GLN
1	C	135	HIS
1	C	168	HIS
1	C	235	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry

4 ligands are modelled in this entry.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	223/249 (89%)	-0.12	11 (4%) 36 36	44, 55, 99, 144	0
1	B	220/249 (88%)	0.38	13 (5%) 29 29	39, 71, 113, 163	1 (0%)
1	C	220/249 (88%)	0.72	27 (12%) 9 9	45, 87, 143, 165	0
All	All	663/747 (88%)	0.32	51 (7%) 21 20	39, 67, 127, 165	1 (0%)

All (51) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	94	ALA	4.8
1	B	159	SER	4.4
1	A	21	ALA	4.3
1	C	243	PHE	4.2
1	B	21	ALA	4.1
1	C	158	LEU	3.8
1	A	244	ALA	3.8
1	C	105	ILE	3.7
1	C	21	ALA	3.6
1	C	88	LEU	3.4
1	C	114	TYR	3.4
1	C	151	ASN	3.4
1	C	117	LEU	3.3
1	A	247	SER	3.2
1	A	243	PHE	3.2
1	C	101	PHE	3.2
1	C	160	THR	3.0
1	C	37	LEU	3.0
1	C	113	LEU	3.0
1	C	93	VAL	3.0
1	B	244	ALA	3.0
1	B	116	LEU	2.9
1	C	90	ALA	2.9

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	C	22	ILE	2.9
1	C	179	LEU	2.8
1	B	246	LYS	2.7
1	B	152	THR	2.7
1	C	96	ALA	2.6
1	C	84	ALA	2.6
1	C	246	LYS	2.6
1	C	89	ALA	2.5
1	A	246	LYS	2.5
1	B	243	PHE	2.5
1	C	244	ALA	2.5
1	A	158	LEU	2.4
1	A	77	GLU	2.4
1	A	245	GLN	2.3
1	A	90	ALA	2.3
1	B	160	THR	2.3
1	B	36	TYR	2.3
1	B	24	ALA	2.3
1	B	161	ASN	2.2
1	C	239	ALA	2.2
1	B	32	PRO	2.2
1	C	41	GLN	2.2
1	A	153	LYS	2.2
1	C	104	PRO	2.1
1	A	161	ASN	2.1
1	C	159	SER	2.0
1	C	34	LEU	2.0
1	B	28	ALA	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum,

median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	PO4	A	301	5/5	0.84	0.12	80,83,83,87	0
2	PO4	C	301	5/5	0.90	0.09	91,93,97,107	0
2	PO4	B	301	5/5	0.91	0.08	76,77,84,88	0
3	PEG	C	302	7/7	0.92	0.11	52,67,78,85	0

6.5 Other polymers [i](#)

There are no such residues in this entry.