



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 16, 2026 – 06:09 pm GMT

PDB ID : 9QHR / pdb_00009qhr
Title : Crystal structure of the second bromodomain of C. albicans Rsc1
Authors : Petosa, C.; Sayou, C.; Govin, J.
Deposited on : 2025-03-16
Resolution : 1.50 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 1.8.4, CSD as541be (2020)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.48.1


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	164625	3717 (1.50-1.50)
Clashscore	180529	4048 (1.50-1.50)
Ramachandran outliers	177936	3970 (1.50-1.50)
Sidechain outliers	177891	3967 (1.50-1.50)
RSRZ outliers	164620	3718 (1.50-1.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	118	4% 92% 8% • •
1	B	118	3% 96% • •
1	C	118	6% 95% • •
1	D	118	12% 92% 8% •

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 4614 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Chromatin structure-remodeling complex subunit RSC1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	117	Total	C	N	O	Se	0	0	0
			974	627	166	180	1			
1	B	116	Total	C	N	O	Se	0	0	0
			970	625	165	179	1			
1	C	117	Total	C	N	O	Se	0	0	0
			974	627	166	180	1			
1	D	117	Total	C	N	O	Se	0	0	0
			974	627	166	180	1			

There are 8 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	223	GLY	-	expression tag	UNP A0A1D8PCV8
A	224	ALA	-	expression tag	UNP A0A1D8PCV8
B	223	GLY	-	expression tag	UNP A0A1D8PCV8
B	224	ALA	-	expression tag	UNP A0A1D8PCV8
C	223	GLY	-	expression tag	UNP A0A1D8PCV8
C	224	ALA	-	expression tag	UNP A0A1D8PCV8
D	223	GLY	-	expression tag	UNP A0A1D8PCV8
D	224	ALA	-	expression tag	UNP A0A1D8PCV8


- Molecule 2 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	183	Total	O	0	0
			183	183		
2	B	212	Total	O	0	0
			212	212		
2	C	167	Total	O	0	0
			167	167		
2	D	160	Total	O	0	0
			160	160		



3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Chromatin structure-remodeling complex subunit RSC1




- Molecule 1: Chromatin structure-remodeling complex subunit RSC1



- Molecule 1: Chromatin structure-remodeling complex subunit RSC1



- Molecule 1: Chromatin structure-remodeling complex subunit RSC1



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	66.18Å 68.65Å 130.27Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	38.50 – 1.50 38.50 – 1.50	Depositor EDS
% Data completeness (in resolution range)	99.0 (38.50-1.50) 99.3 (38.50-1.50)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.73 (at 1.50Å)	Xtriage
Refinement program	PHENIX 1.15.2	Depositor
R, R_{free}	0.202 , 0.226 0.205 , 0.232	Depositor DCC
R_{free} test set	998 reflections (1.05%)	wwPDB-VP
Wilson B-factor (Å ²)	22.4	Xtriage
Anisotropy	0.431	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.31 , 25.2	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	0.025 for k,h,-l	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	4614	wwPDB-VP
Average B, all atoms (Å ²)	29.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.55% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.28	0/994	0.48	0/1334
1	B	0.25	0/990	0.44	0/1329
1	C	0.28	0/994	0.47	0/1334
1	D	0.25	0/994	0.39	0/1334
All	All	0.26	0/3972	0.45	0/5331

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	974	0	980	7	0
1	B	970	0	977	2	0
1	C	974	0	980	4	0
1	D	974	0	980	7	0
2	A	183	0	0	4	0
2	B	212	0	0	0	0
2	C	167	0	0	2	1
2	D	160	0	0	3	1
All	All	4614	0	3917	19	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (19) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:313:ASP:OD1	2:D:401:HOH:O	2.15	0.63
1:D:224:ALA:O	1:D:287:ARG:NH2	2.35	0.59
1:A:312:ASN:ND2	2:A:405:HOH:O	2.37	0.56
1:D:232:ARG:NH2	2:D:407:HOH:O	2.38	0.56
1:D:330:HIS:O	1:D:334:GLN:HG2	2.08	0.53
1:A:238:LYS:HA	1:A:241:LYS:HE2	1.91	0.52
1:A:317:GLU:HG2	2:A:404:HOH:O	2.17	0.44
1:A:285:ARG:NH2	2:A:414:HOH:O	2.51	0.44
1:C:262:LYS:HG3	2:C:401:HOH:O	2.19	0.42
1:D:251:LEU:HD23	1:D:251:LEU:HA	1.88	0.42
1:A:242:LYS:NZ	2:A:402:HOH:O	2.26	0.41
1:A:285:ARG:CZ	1:B:265:GLY:HA3	2.50	0.41
1:D:287:ARG:NH2	2:D:405:HOH:O	2.53	0.41
1:C:307:GLN:O	1:C:311:GLU:HB2	2.20	0.41
1:A:252:THR:HG22	1:A:325:PHE:CZ	2.56	0.40
1:C:260:ASP:OD1	2:C:401:HOH:O	2.21	0.40
1:C:285:ARG:HA	1:C:285:ARG:HD3	1.88	0.40
1:D:256:GLU:OE1	1:D:281:ARG:NH2	2.47	0.40
1:B:229:PHE:HB2	1:B:292:VAL:HG23	2.03	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:C:524:HOH:O	2:D:486:HOH:O[4_556]	2.17	0.03

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	115/118 (98%)	115 (100%)	0	0	100	100
1	B	114/118 (97%)	113 (99%)	1 (1%)	0	100	100
1	C	115/118 (98%)	115 (100%)	0	0	100	100
1	D	115/118 (98%)	115 (100%)	0	0	100	100
All	All	459/472 (97%)	458 (100%)	1 (0%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	107/107 (100%)	106 (99%)	1 (1%)	75	57
1	B	107/107 (100%)	107 (100%)	0	100	100
1	C	107/107 (100%)	107 (100%)	0	100	100
1	D	107/107 (100%)	107 (100%)	0	100	100
All	All	428/428 (100%)	427 (100%)	1 (0%)	92	83

All (1) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	311	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (9) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	249	HIS
1	A	278	ASN
1	B	278	ASN
1	B	323	GLN
1	C	246	ASN
1	C	249	HIS

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	C	278	ASN
1	C	297	ASN
1	D	323	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	116/118 (98%)	0.54	5 (4%) 40 43	16, 25, 39, 50	0
1	B	115/118 (97%)	0.53	3 (2%) 57 60	17, 26, 41, 57	0
1	C	116/118 (98%)	0.64	7 (6%) 29 30	18, 26, 43, 51	0
1	D	116/118 (98%)	0.90	14 (12%) 10 9	18, 31, 58, 78	0
All	All	463/472 (98%)	0.65	29 (6%) 27 28	16, 26, 48, 78	0

All (29) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	331	LEU	4.9
1	B	337	LEU	4.9
1	D	332	VAL	3.6
1	D	223	GLY	3.6
1	B	315	TYR	3.5
1	D	329	ALA	3.5
1	D	333	ILE	3.4
1	D	337	LEU	3.3
1	A	315	TYR	3.1
1	D	315	TYR	2.9
1	C	330	HIS	2.9
1	A	224	ALA	2.8
1	D	263	THR	2.7
1	A	310	TYR	2.7
1	A	223	GLY	2.6
1	D	224	ALA	2.6
1	C	337	LEU	2.5
1	C	338	SER	2.5
1	D	338	SER	2.5
1	D	335	THR	2.5
1	A	337	LEU	2.5

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	D	261	ILE	2.4
1	C	312	ASN	2.4
1	B	338	SER	2.3
1	C	315	TYR	2.3
1	C	310	TYR	2.2
1	D	232	ARG	2.1
1	C	331	LEU	2.0
1	D	339	LYS	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.