



# Full wwPDB X-ray Structure Validation Report ⓘ

Dec 15, 2025 – 03:18 pm GMT

PDB ID : 9RB2 / pdb\_00009rb2  
Title : Crystal Structure of glxR  
Authors : Parkhill, S.; little, O.; Wang, M.; Askenasy, I.; Brear, P.; Cai, W.; Welch, M.  
Deposited on : 2025-05-21  
Resolution : 2.21 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0  
Xtriage (Phenix) : 2.0  
EDS : 3.0  
Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)  
CCP4 : 9.0.010 (Gargrove)  
Density-Fitness : 1.0.12  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.47


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.21 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	164625	7167 (2.24-2.20)
Clashscore	180529	8096 (2.24-2.20)
Ramachandran outliers	177936	8010 (2.24-2.20)
Sidechain outliers	177891	8011 (2.24-2.20)
RSRZ outliers	164620	7166 (2.24-2.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	312	
1	B	312	
1	C	312	
1	D	312	

## 2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 9250 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Probable oxidoreductase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	301	Total	C	N	O	S	0	4	0
			2256	1417	400	426	13			
1	B	296	Total	C	N	O	S	0	3	0
			2195	1377	389	416	13			
1	C	300	Total	C	N	O	S	0	5	0
			2251	1415	400	422	14			
1	D	297	Total	C	N	O	S	0	1	0
			2187	1372	389	413	13			

There are 64 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-15	MET	-	initiating methionine	UNP Q9I3L2
A	-14	GLY	-	expression tag	UNP Q9I3L2
A	-13	HIS	-	expression tag	UNP Q9I3L2
A	-12	HIS	-	expression tag	UNP Q9I3L2
A	-11	HIS	-	expression tag	UNP Q9I3L2
A	-10	HIS	-	expression tag	UNP Q9I3L2
A	-9	HIS	-	expression tag	UNP Q9I3L2
A	-8	HIS	-	expression tag	UNP Q9I3L2
A	-7	ALA	-	expression tag	UNP Q9I3L2
A	-6	GLU	-	expression tag	UNP Q9I3L2
A	-5	ASN	-	expression tag	UNP Q9I3L2
A	-4	LEU	-	expression tag	UNP Q9I3L2
A	-3	TYR	-	expression tag	UNP Q9I3L2
A	-2	PHE	-	expression tag	UNP Q9I3L2
A	-1	GLN	-	expression tag	UNP Q9I3L2
A	0	HIS	-	expression tag	UNP Q9I3L2
B	-15	MET	-	initiating methionine	UNP Q9I3L2
B	-14	GLY	-	expression tag	UNP Q9I3L2
B	-13	HIS	-	expression tag	UNP Q9I3L2
B	-12	HIS	-	expression tag	UNP Q9I3L2
B	-11	HIS	-	expression tag	UNP Q9I3L2

*Continued on next page...*

*Continued from previous page...*

Chain	Residue	Modelled	Actual	Comment	Reference
B	-10	HIS	-	expression tag	UNP Q9I3L2
B	-9	HIS	-	expression tag	UNP Q9I3L2
B	-8	HIS	-	expression tag	UNP Q9I3L2
B	-7	ALA	-	expression tag	UNP Q9I3L2
B	-6	GLU	-	expression tag	UNP Q9I3L2
B	-5	ASN	-	expression tag	UNP Q9I3L2
B	-4	LEU	-	expression tag	UNP Q9I3L2
B	-3	TYR	-	expression tag	UNP Q9I3L2
B	-2	PHE	-	expression tag	UNP Q9I3L2
B	-1	GLN	-	expression tag	UNP Q9I3L2
B	0	HIS	-	expression tag	UNP Q9I3L2
C	-15	MET	-	initiating methionine	UNP Q9I3L2
C	-14	GLY	-	expression tag	UNP Q9I3L2
C	-13	HIS	-	expression tag	UNP Q9I3L2
C	-12	HIS	-	expression tag	UNP Q9I3L2
C	-11	HIS	-	expression tag	UNP Q9I3L2
C	-10	HIS	-	expression tag	UNP Q9I3L2
C	-9	HIS	-	expression tag	UNP Q9I3L2
C	-8	HIS	-	expression tag	UNP Q9I3L2
C	-7	ALA	-	expression tag	UNP Q9I3L2
C	-6	GLU	-	expression tag	UNP Q9I3L2
C	-5	ASN	-	expression tag	UNP Q9I3L2
C	-4	LEU	-	expression tag	UNP Q9I3L2
C	-3	TYR	-	expression tag	UNP Q9I3L2
C	-2	PHE	-	expression tag	UNP Q9I3L2
C	-1	GLN	-	expression tag	UNP Q9I3L2
C	0	HIS	-	expression tag	UNP Q9I3L2
D	-15	MET	-	initiating methionine	UNP Q9I3L2
D	-14	GLY	-	expression tag	UNP Q9I3L2
D	-13	HIS	-	expression tag	UNP Q9I3L2
D	-12	HIS	-	expression tag	UNP Q9I3L2
D	-11	HIS	-	expression tag	UNP Q9I3L2
D	-10	HIS	-	expression tag	UNP Q9I3L2
D	-9	HIS	-	expression tag	UNP Q9I3L2
D	-8	HIS	-	expression tag	UNP Q9I3L2
D	-7	ALA	-	expression tag	UNP Q9I3L2
D	-6	GLU	-	expression tag	UNP Q9I3L2
D	-5	ASN	-	expression tag	UNP Q9I3L2
D	-4	LEU	-	expression tag	UNP Q9I3L2
D	-3	TYR	-	expression tag	UNP Q9I3L2
D	-2	PHE	-	expression tag	UNP Q9I3L2
D	-1	GLN	-	expression tag	UNP Q9I3L2

*Continued on next page...*

*Continued from previous page...*

Chain	Residue	Modelled	Actual	Comment	Reference
D	0	HIS	-	expression tag	UNP Q9I3L2


- Molecule 2 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	98	Total 98	O 98	0	0
2	B	92	Total 92	O 92	0	0
2	C	87	Total 87	O 87	0	0
2	D	84	Total 84	O 84	0	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Probable oxidoreductase




- Molecule 1: Probable oxidoreductase




- Molecule 1: Probable oxidoreductase



- Molecule 1: Probable oxidoreductase





## 4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	79.18Å 109.21Å 146.69Å 90.00° 93.39° 90.00°	Depositor
Resolution (Å)	73.22 – 2.21 73.22 – 2.21	Depositor EDS
% Data completeness (in resolution range)	98.5 (73.22-2.21) 98.5 (73.22-2.21)	Depositor EDS
$R_{merge}$	0.45	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.30 (at 2.20Å)	Xtriage
Refinement program	BUSTER 2.10.4 (10-JUL-2024)	Depositor
R, $R_{free}$	0.235 , 0.273 0.226 , 0.260	Depositor DCC
$R_{free}$ test set	2037 reflections (3.31%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	31.9	Xtriage
Anisotropy	0.496	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.34 , 38.9	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.94	EDS
Total number of atoms	9250	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	39.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.73% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.71	1/2294 (0.0%)	1.12	6/3100 (0.2%)
1	B	0.72	0/2230	1.11	3/3013 (0.1%)
1	C	0.68	0/2289	1.07	2/3094 (0.1%)
1	D	0.69	0/2223	1.09	2/3005 (0.1%)
All	All	0.70	1/9036 (0.0%)	1.10	13/12212 (0.1%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	-1	GLN	CA-C	6.31	1.61	1.52

All (13) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	-2	PHE	CA-CB-CG	-7.99	105.81	113.80
1	B	231	GLY	N-CA-C	-7.04	96.48	113.18
1	C	231	GLY	N-CA-C	-6.51	97.76	113.18
1	A	231	GLY	N-CA-C	-6.28	98.30	113.18
1	A	-2	PHE	CA-C-N	6.21	133.01	122.26
1	A	-2	PHE	C-N-CA	6.21	133.01	122.26
1	B	137	VAL	N-CA-C	5.73	116.73	108.48
1	A	-1	GLN	N-CA-C	-5.35	106.80	113.38
1	B	66	PRO	N-CA-C	5.24	120.81	113.53
1	D	232	PHE	CA-CB-CG	-5.17	108.63	113.80
1	A	50	ASN	N-CA-C	5.15	112.86	108.22
1	D	231	GLY	N-CA-C	-5.14	101.00	113.18
1	C	67	ASP	N-CA-C	5.01	115.93	108.86

There are no chirality outliers.

There are no planarity outliers.

## 5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2256	0	2263	9	0
1	B	2195	0	2214	11	0
1	C	2251	0	2265	15	0
1	D	2187	0	2204	12	0
2	A	98	0	0	0	0
2	B	92	0	0	0	0
2	C	87	0	0	0	0
2	D	84	0	0	0	0
All	All	9250	0	8946	46	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (46) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:151:PHE:HB3	1:D:158:ILE:HG12	1.79	0.64
1:D:207:MET:HE1	1:D:217:GLU:HG2	1.82	0.62
1:C:151:PHE:HB3	1:C:158:ILE:HG12	1.85	0.58
1:C:250[B]:GLU:HG3	1:C:251:LEU:HG	1.88	0.56
1:B:233:ARG:HD2	1:B:275:ASN:ND2	2.21	0.55
1:A:121:VAL:HG12	1:A:135:ILE:HG12	1.88	0.55
1:C:17:ALA:HA	1:C:20:LEU:HD12	1.91	0.53
1:D:71:VAL:HG21	1:D:97:ILE:HG21	1.91	0.52
1:B:7:ILE:HG21	1:B:74:VAL:HG21	1.91	0.51
1:D:219:HIS:O	1:D:223:MET:HG3	2.11	0.50
1:A:32:HIS:CE1	1:A:33:HIS:CE1	2.99	0.50
1:D:186:ALA:HA	1:D:282:ILE:HD12	1.93	0.50
1:A:179:ALA:HB2	1:A:237:HIS:NE2	2.26	0.50
1:C:132:THR:HA	1:C:156:LYS:HB2	1.94	0.49
1:C:215:ILE:CD1	1:C:219:HIS:CE1	2.96	0.49
1:C:207[A]:MET:HE1	1:C:217:GLU:HA	1.94	0.49
1:B:121:VAL:HG12	1:B:135:ILE:HG12	1.95	0.48
1:B:52:LYS:HE3	1:B:79:ASP:HA	1.96	0.48
1:B:215:ILE:HD12	1:B:219:HIS:CE1	2.50	0.47

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:215:ILE:HD13	1:C:219:HIS:ND1	2.30	0.47
1:D:215:ILE:HG23	1:D:219:HIS:HB2	1.97	0.46
1:B:233:ARG:HH11	1:B:275:ASN:ND2	2.13	0.46
1:B:29:LEU:HD13	1:B:45:ALA:HB1	1.96	0.46
1:D:121:VAL:HG12	1:D:135:ILE:HG12	1.97	0.46
1:B:215:ILE:HD12	1:B:219:HIS:ND1	2.31	0.46
1:C:247:GLY:HA2	1:C:250[B]:GLU:HG2	1.98	0.45
1:D:53:GLU:O	1:D:57:GLU:HG2	2.16	0.45
1:A:253:LEU:HD21	1:B:194:ARG:HG2	1.97	0.45
1:B:25:HIS:CE1	1:B:150:LEU:CD2	2.99	0.45
1:C:121:VAL:HG12	1:C:135:ILE:HG12	1.98	0.44
1:D:7:ILE:HG21	1:D:74:VAL:HG21	1.99	0.44
1:C:215:ILE:CD1	1:C:219:HIS:ND1	2.81	0.44
1:D:86:GLY:HA2	1:D:112:THR:O	2.17	0.43
1:C:55:ALA:HA	1:C:61:ILE:HG12	2.01	0.43
1:C:215:ILE:HD13	1:C:219:HIS:CE1	2.53	0.43
1:D:117:LEU:HD23	1:D:139:GLY:HA3	1.99	0.43
1:A:71:VAL:HG21	1:A:97:ILE:HG21	2.01	0.43
1:B:215:ILE:CD1	1:B:219:HIS:CE1	3.02	0.42
1:A:171:LYS:O	1:A:175:GLN:HG2	2.20	0.42
1:D:96:SER:OG	1:D:171:LYS:HE2	2.18	0.42
1:C:215:ILE:HG23	1:C:219:HIS:HB2	2.02	0.42
1:C:202:VAL:O	1:C:206:LEU:HG	2.21	0.41
1:A:192:ALA:HB1	1:A:202:VAL:HG21	2.02	0.41
1:C:29:LEU:HD12	1:C:36:ALA:HB1	2.03	0.41
1:A:202:VAL:O	1:A:206:LEU:HG	2.20	0.41
1:A:140[B]:CYS:HA	1:A:141:PRO:HD3	1.98	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles

### 5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	303/312 (97%)	294 (97%)	9 (3%)	0	100	100
1	B	297/312 (95%)	286 (96%)	10 (3%)	1 (0%)	37	41
1	C	303/312 (97%)	294 (97%)	9 (3%)	0	100	100
1	D	296/312 (95%)	287 (97%)	8 (3%)	1 (0%)	37	41
All	All	1199/1248 (96%)	1161 (97%)	36 (3%)	2 (0%)	44	51

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	2	ALA
1	D	231	GLY

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	229/234 (98%)	223 (97%)	6 (3%)	41	52
1	B	223/234 (95%)	219 (98%)	4 (2%)	54	67
1	C	229/234 (98%)	225 (98%)	4 (2%)	56	69
1	D	222/234 (95%)	216 (97%)	6 (3%)	40	51
All	All	903/936 (96%)	883 (98%)	20 (2%)	48	59

All (20) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	18	GLN
1	A	59[A]	GLU
1	A	59[B]	GLU
1	A	78	LYS
1	A	212	SER
1	A	278	HIS
1	B	74	VAL
1	B	78	LYS
1	B	207	MET

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	B	278	HIS
1	C	1	MET
1	C	212	SER
1	C	217	GLU
1	C	225	LYS
1	D	88	ASN
1	D	212	SER
1	D	237	HIS
1	D	250	GLU
1	D	275	ASN
1	D	278	HIS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (22) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	18	GLN
1	A	32	HIS
1	A	33	HIS
1	A	56	GLN
1	A	263	GLN
1	A	275	ASN
1	B	25	HIS
1	B	33	HIS
1	B	174	ASN
1	B	175	GLN
1	B	195	ASN
1	B	263	GLN
1	B	275	ASN
1	C	-1	GLN
1	C	18	GLN
1	C	152	GLN
1	C	263	GLN
1	D	0	HIS
1	D	174	ASN
1	D	175	GLN
1	D	263	GLN
1	D	275	ASN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

There are no ligands in this entry.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	301/312 (96%)	0.16	7 (2%) 61 58	17, 37, 52, 69	4 (1%)
1	B	296/312 (94%)	0.16	4 (1%) 73 71	18, 36, 60, 70	3 (1%)
1	C	300/312 (96%)	0.32	7 (2%) 61 58	18, 40, 55, 69	5 (1%)
1	D	297/312 (95%)	0.10	3 (1%) 79 77	20, 37, 49, 68	1 (0%)
All	All	1194/1248 (95%)	0.19	21 (1%) 67 64	17, 38, 54, 70	13 (1%)

All (21) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	-4	LEU	5.6
1	C	-3	TYR	4.0
1	C	-2	PHE	3.9
1	B	1	MET	3.8
1	D	295	ASP	3.5
1	A	-4	LEU	3.0
1	D	33	HIS	3.0
1	A	249[A]	ARG	2.7
1	A	296	GLU	2.7
1	A	-1	GLN	2.6
1	B	40	LEU	2.6
1	C	27	LEU	2.6
1	C	-1	GLN	2.4
1	A	59[A]	GLU	2.3
1	C	35	ALA	2.3
1	B	45	ALA	2.2
1	A	1	MET	2.2
1	C	295	ASP	2.0
1	D	0	HIS	2.0
1	B	41	LEU	2.0
1	A	-2	PHE	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

## 6.4 Ligands [i](#)

There are no ligands in this entry.

## 6.5 Other polymers [i](#)

There are no such residues in this entry.