



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 10, 2026 – 03:10 pm GMT

PDB ID : 9RS1 / pdb_00009rs1
Title : Crystal structure of the human METTL3-METTL14 in complex with compound 2 (T148)
Authors : Bedi, R.K.; Cafisch, A.
Deposited on : 2025-06-30
Resolution : 1.73 Å (reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 1.8.4, CSD as541be (2020)
Xtrriage (Phenix) : 2.0
EDS : 3.0
buster-report : 1.1.7 (2018)
Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.48.1


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.73 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	164625	1043 (1.74-1.74)
Clashscore	180529	1119 (1.74-1.74)
Ramachandran outliers	177936	1112 (1.74-1.74)
Sidechain outliers	177891	1112 (1.74-1.74)
RSRZ outliers	164620	1043 (1.74-1.74)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	228	 8% 81% 5% 14%
2	B	307	 13% 76% 5% 20%

2 Entry composition [i](#)

There are 6 unique types of molecules in this entry. The entry contains 3743 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called N6-adenosine-methyltransferase catalytic subunit.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	197	1558	1000	273	276	9	0	0	0

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	353	MET	-	initiating methionine	UNP Q86U44


- Molecule 2 is a protein called N(6)-adenosine-methyltransferase non-catalytic subunit METTL14.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	B	246	1884	1207	321	343	13	0	2	0

There are 18 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
B	89	MET	-	initiating methionine	UNP Q9HCE5
B	90	HIS	-	expression tag	UNP Q9HCE5
B	91	HIS	-	expression tag	UNP Q9HCE5
B	92	HIS	-	expression tag	UNP Q9HCE5
B	93	HIS	-	expression tag	UNP Q9HCE5
B	94	HIS	-	expression tag	UNP Q9HCE5
B	95	HIS	-	expression tag	UNP Q9HCE5
B	96	SER	-	expression tag	UNP Q9HCE5
B	97	SER	-	expression tag	UNP Q9HCE5
B	98	GLY	-	expression tag	UNP Q9HCE5
B	99	ARG	-	expression tag	UNP Q9HCE5
B	100	GLU	-	expression tag	UNP Q9HCE5
B	101	ASN	-	expression tag	UNP Q9HCE5
B	102	LEU	-	expression tag	UNP Q9HCE5

Continued on next page...



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	B	1	Total	C	O	0	0
			4	2	2		

- Molecule 5 is 4-[[[4-[(4,4-dimethylpiperidin-1-yl)methyl]-2-oxidanyl-phenyl]amino]methyl]-1-[6-[(phenylmethyl)amino]pyrimidin-4-yl]piperidin-4-ol (CCD ID: A1JI8) (formula: C₃₁H₄₂N₆O₂) (labeled as "Ligand of Interest" by depositor).




Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	91	Total 91	O 91	0	0
6	B	141	Total 141	O 141	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: N6-adenosine-methyltransferase catalytic subunit



- Molecule 2: N(6)-adenosine-methyltransferase non-catalytic subunit METTL14



4 Data and refinement statistics i

Property	Value	Source
Space group	P 32 2 1	Depositor
Cell constants a, b, c, α , β , γ	63.86Å 63.86Å 225.75Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	44.56 – 1.73 44.56 – 1.73	Depositor EDS
% Data completeness (in resolution range)	99.8 (44.56-1.73) 99.9 (44.56-1.73)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.86 (at 1.73Å)	Xtrriage
Refinement program	PHENIX 1.21.2_5419	Depositor
R, R_{free}	0.200 , 0.220 0.200 , 0.220	Depositor DCC
R_{free} test set	2862 reflections (4.99%)	wwPDB-VP
Wilson B-factor (Å ²)	34.4	Xtrriage
Anisotropy	0.080	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 41.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	0.023 for -h,-k,l	Xtrriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	3743	wwPDB-VP
Average B, all atoms (Å ²)	40.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.97% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: ACT, SAH, A1JI8

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.27	0/1597	0.48	0/2173
2	B	0.30	0/1938	0.49	0/2639
All	All	0.29	0/3535	0.49	0/4812

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1558	0	1502	7	0
2	B	1884	0	1722	10	0
3	A	26	0	18	0	0
4	B	4	0	3	0	0
5	B	39	0	0	2	0
6	A	91	0	0	1	0
6	B	141	0	0	0	0
All	All	3743	0	3245	16	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (16) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:183:THR:HG21	5:B:402:A1JI8:C34	2.34	0.58
1:A:381:LEU:HD13	1:A:574:ILE:HD13	1.88	0.55
2:B:341:ARG:HD2	2:B:385:LEU:HB2	1.89	0.54
2:B:183:THR:HG22	2:B:184:PRO:HD2	1.90	0.52
2:B:198:TYR:HB3	2:B:324:ILE:HG22	1.92	0.51
1:A:373:TRP:HB2	1:A:551:LEU:HD13	1.92	0.50
1:A:391:VAL:HB	1:A:530:LYS:HG2	1.93	0.49
2:B:183:THR:HG21	5:B:402:A1JI8:C33	2.44	0.47
1:A:456:ILE:HD13	2:B:260[B]:CYS:SG	2.57	0.44
2:B:260[A]:CYS:SG	2:B:287:HIS:CE1	3.12	0.43
1:A:539:ASN:O	1:A:541:GLN:HG2	2.19	0.43
2:B:260[B]:CYS:SG	2:B:285:LYS:HD2	2.59	0.43
2:B:311:ILE:HG23	2:B:313:LEU:H	1.84	0.42
2:B:341:ARG:CD	2:B:385:LEU:HB2	2.51	0.41
1:A:436:ALA:HA	1:A:439:LEU:HB3	2.04	0.40
1:A:490:ASN:HB2	6:A:752:HOH:O	2.22	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	191/228 (84%)	188 (98%)	3 (2%)	0	100	100
2	B	240/307 (78%)	230 (96%)	9 (4%)	1 (0%)	30	16
All	All	431/535 (81%)	418 (97%)	12 (3%)	1 (0%)	44	29

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	B	271	LYS

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	163/200 (82%)	163 (100%)	0	100	100
2	B	186/274 (68%)	186 (100%)	0	100	100
All	All	349/474 (74%)	349 (100%)	0	100	100

There are no protein residues with a non-rotameric sidechain to report.

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
2	B	122	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

3 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	SAH	A	601	-	24,28,28	4.68	8 (33%)	25,40,40	3.20	7 (28%)
5	A1JI8	B	402	-	42,43,43	2.34	11 (26%)	51,61,61	1.73	6 (11%)
4	ACT	B	401	-	3,3,3	1.42	0	3,3,3	0.63	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	SAH	A	601	-	-	2/11/31/31	0/3/3/3
5	A1JI8	B	402	-	-	1/19/43/43	0/5/5/5

All (19) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	601	SAH	O4'-C1'	14.91	1.61	1.41
3	A	601	SAH	C2'-C1'	-14.69	1.31	1.53
5	B	402	A1JI8	C07-N06	-7.18	1.33	1.47
5	B	402	A1JI8	C27-N28	6.48	1.45	1.36
3	A	601	SAH	O4'-C4'	-5.82	1.32	1.45
5	B	402	A1JI8	C05-N06	-4.96	1.33	1.46
5	B	402	A1JI8	C38-N06	-4.87	1.33	1.46
5	B	402	A1JI8	C21-N22	-3.85	1.40	1.46
5	B	402	A1JI8	C23-N22	3.61	1.44	1.37
5	B	402	A1JI8	C18-C17	3.49	1.56	1.52
3	A	601	SAH	C6-N6	3.19	1.45	1.34
5	B	402	A1JI8	C14-N15	3.12	1.46	1.37
5	B	402	A1JI8	O37-C17	-3.09	1.38	1.44
3	A	601	SAH	O2'-C2'	2.97	1.50	1.43
3	A	601	SAH	O3'-C3'	-2.79	1.36	1.43
3	A	601	SAH	C2-N3	2.59	1.36	1.32
3	A	601	SAH	C5-C4	-2.40	1.34	1.40
5	B	402	A1JI8	C19-N22	-2.12	1.43	1.46

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	B	402	A1JI8	C36-C23	-2.08	1.36	1.39

All (13) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	601	SAH	C5-C6-N6	10.00	135.56	120.35
3	A	601	SAH	N6-C6-N1	-7.12	103.79	118.57
5	B	402	A1JI8	N26-C25-N24	-5.71	119.67	128.60
5	B	402	A1JI8	C25-N24-C23	5.66	119.89	114.94
3	A	601	SAH	O4'-C1'-C2'	-5.29	99.19	106.93
3	A	601	SAH	C1'-N9-C4	-5.20	117.51	126.64
3	A	601	SAH	N3-C2-N1	-5.16	120.61	128.68
5	B	402	A1JI8	C19-N22-C21	4.35	121.12	111.52
5	B	402	A1JI8	C39-C38-N06	4.27	115.45	111.23
5	B	402	A1JI8	C29-N28-C27	-2.89	117.71	123.40
3	A	601	SAH	OXT-C-CA	2.30	121.23	113.38
3	A	601	SAH	C5'-SD-CG	2.13	108.65	102.27
5	B	402	A1JI8	C36-C23-N22	-2.06	119.84	122.29

There are no chirality outliers.

All (3) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	601	SAH	O-C-CA-N
5	B	402	A1JI8	N15-C16-C17-O37
3	A	601	SAH	C3'-C4'-C5'-SD


There are no ring outliers.

1 monomer is involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	B	402	A1JI8	2	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the

average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	197/228 (86%)	0.64	18 (9%) 16 22	26, 39, 57, 75	0
2	B	246/307 (80%)	0.79	39 (15%) 6 8	20, 37, 72, 86	2 (0%)
All	All	443/535 (82%)	0.72	57 (12%) 9 12	20, 38, 69, 86	2 (0%)

All (57) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	B	179	ILE	5.2
2	B	184	PRO	5.2
1	A	368	LEU	4.9
2	B	177	PHE	4.8
1	A	574	ILE	4.4
2	B	182	LEU	4.2
2	B	272	THR	4.1
2	B	152	LEU	4.1
2	B	153	ILE	4.0
2	B	295	THR	3.9
2	B	202	THR	3.8
2	B	339	LEU	3.8
2	B	274	THR	3.7
2	B	137	VAL	3.7
2	B	322	GLY	3.7
2	B	183	THR	3.5
2	B	321	ILE	3.5
2	B	275	LEU	3.5
1	A	576	LYS	3.4
1	A	467	ILE	3.3
2	B	309	VAL	3.3
2	B	392	ILE	3.3
1	A	573	ILE	3.2
2	B	256	CYS	3.0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
2	B	180	ARG	3.0
2	B	117	ASN	2.9
1	A	400	ILE	2.9
1	A	474	HIS	2.8
1	A	407	GLY	2.8
1	A	490	ASN	2.8
2	B	273	LYS	2.7
2	B	308	ASN	2.7
2	B	155	LEU	2.6
1	A	575	SER	2.6
1	A	369	PHE	2.6
1	A	494	PHE	2.6
2	B	136	ASP	2.6
2	B	311	ILE	2.6
1	A	463	LEU	2.6
2	B	181	GLU	2.6
2	B	390	GLU	2.5
1	A	475	TRP	2.5
2	B	384	TYR	2.5
2	B	385	LEU	2.4
2	B	271	LYS	2.4
1	A	509	SER	2.4
2	B	154	ARG	2.4
1	A	571	ASP	2.2
2	B	268	ASN	2.2
2	B	270	GLY	2.2
1	A	498	LEU	2.2
2	B	151	GLU	2.1
2	B	310	ASP	2.1
2	B	258	ASP	2.1
2	B	387	GLY	2.1
1	A	446	LEU	2.0
2	B	176	ALA	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)


There are no oligosaccharides in this entry.


6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
5	A1JI8	B	402	39/39	0.84	0.14	36,45,55,58	0
3	SAH	A	601	26/26	0.88	0.11	34,43,47,49	0
4	ACT	B	401	4/4	0.93	0.10	32,32,34,34	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.





6.5 Other polymers [i](#)

There are no such residues in this entry.