



Full wwPDB X-ray Structure Validation Report ⓘ

Jul 6, 2026 – 03:01 pm BST

PDB ID : 9RSF / pdb_00009rsf
Title : Macaque CD1d with lipid antigen bound
Authors : Burns, D.; Maly, M.; Tews, I.; Mansour, S.
Deposited on : 2025-07-01
Resolution : 1.83 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	1.8.4, CSD as541be (2020)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.015 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.50


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.83 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1296 (1.84-1.84)
Clashscore	190562	1329 (1.84-1.84)
Ramachandran outliers	187476	1318 (1.84-1.84)
Sidechain outliers	187428	1318 (1.84-1.84)
RSRZ outliers	180081	1296 (1.84-1.84)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	296	3% 83% 9% 8%
1	C	296	5% 86% 5% 7%
2	B	100	% 91% 7% ..
2	D	100	2% 95% ..

2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 13505 atoms, of which 6071 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called CD1d molecule.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	273	Total	C	H	N	O	S	0	14	0
			4395	1439	2137	393	418	8			
1	C	275	Total	C	H	N	O	S	0	7	0
			4360	1430	2116	392	413	9			

There are 42 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	2	MET	-	initiating methionine	UNP F6TR81
A	274	VAL	ILE	engineered mutation	UNP F6TR81
A	279	PRO	-	expression tag	UNP F6TR81
A	280	GLY	-	expression tag	UNP F6TR81
A	281	SER	-	expression tag	UNP F6TR81
A	282	GLY	-	expression tag	UNP F6TR81
A	283	GLY	-	expression tag	UNP F6TR81
A	284	GLY	-	expression tag	UNP F6TR81
A	285	LEU	-	expression tag	UNP F6TR81
A	286	ASN	-	expression tag	UNP F6TR81
A	287	ASP	-	expression tag	UNP F6TR81
A	288	ILE	-	expression tag	UNP F6TR81
A	289	PHE	-	expression tag	UNP F6TR81
A	290	GLU	-	expression tag	UNP F6TR81
A	291	ALA	-	expression tag	UNP F6TR81
A	292	GLN	-	expression tag	UNP F6TR81
A	293	LYS	-	expression tag	UNP F6TR81
A	294	ILE	-	expression tag	UNP F6TR81
A	295	GLU	-	expression tag	UNP F6TR81
A	296	TRP	-	expression tag	UNP F6TR81
A	297	HIS	-	expression tag	UNP F6TR81
C	2	MET	-	initiating methionine	UNP F6TR81
C	274	VAL	ILE	engineered mutation	UNP F6TR81
C	279	PRO	-	expression tag	UNP F6TR81
C	280	GLY	-	expression tag	UNP F6TR81

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
C	281	SER	-	expression tag	UNP F6TR81
C	282	GLY	-	expression tag	UNP F6TR81
C	283	GLY	-	expression tag	UNP F6TR81
C	284	GLY	-	expression tag	UNP F6TR81
C	285	LEU	-	expression tag	UNP F6TR81
C	286	ASN	-	expression tag	UNP F6TR81
C	287	ASP	-	expression tag	UNP F6TR81
C	288	ILE	-	expression tag	UNP F6TR81
C	289	PHE	-	expression tag	UNP F6TR81
C	290	GLU	-	expression tag	UNP F6TR81
C	291	ALA	-	expression tag	UNP F6TR81
C	292	GLN	-	expression tag	UNP F6TR81
C	293	LYS	-	expression tag	UNP F6TR81
C	294	ILE	-	expression tag	UNP F6TR81
C	295	GLU	-	expression tag	UNP F6TR81
C	296	TRP	-	expression tag	UNP F6TR81
C	297	HIS	-	expression tag	UNP F6TR81


- Molecule 2 is a protein called Beta-2-microglobulin.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
2	B	99	Total	C	H	N	O	S	0	3	0
			1631	538	789	141	160	3			
2	D	99	Total	C	H	N	O	S	0	8	0
			1677	551	815	143	165	3			

There are 2 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
B	1	MET	-	initiating methionine	UNP P61769
D	1	MET	-	initiating methionine	UNP P61769

- Molecule 3 is N-{(1S,2R,3S)-1-[(ALPHA-D-GALACTOPYRANOSYLOXY)METHYL]-2,3-DIHYDROXYHEPTADECYL}HEXACOSANAMIDE (CCD ID: AGH) (formula: C₅₀H₉₉NO₉) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total	C	H	N	O	0	0
			153	50	93	1	9		
3	C	1	Total	C	H	N	O	0	0
			153	50	93	1	9		

- Molecule 4 is 1,2-ETHANEDIOL (CCD ID: EDO) (formula: $C_2H_6O_2$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	A	1	Total	C	H	O	0	0
			8	2	4	2		
4	A	1	Total	C	H	O	0	0
			8	2	4	2		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	B	1	Total	C	H	O	0	0
			8	2	4	2		
4	C	1	Total	C	H	O	0	0
			8	2	4	2		
4	C	1	Total	C	H	O	0	0
			8	2	4	2		
4	C	1	Total	C	H	O	0	0
			8	2	4	2		
4	D	1	Total	C	H	O	0	0
			8	2	4	2		

- Molecule 5 is CHLORIDE ION (CCD ID: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	B	1	Total	Cl	0	0
			1	1		
5	C	1	Total	Cl	0	0
			1	1		
5	D	1	Total	Cl	0	0
			1	1		


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	342	Total	O	0	4
			345	345		
6	B	199	Total	O	0	3
			201	201		
6	C	346	Total	O	0	3
			348	348		
6	D	183	Total	O	0	1
			183	183		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: CD1d molecule




- Molecule 1: CD1d molecule



- Molecule 2: Beta-2-microglobulin



- Molecule 2: Beta-2-microglobulin



4 Data and refinement statistics

Property	Value	Source
Space group	P 32 2 1	Depositor
Cell constants a, b, c, α , β , γ	87.06Å 87.06Å 232.81Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	54.08 – 1.83 54.08 – 1.83	Depositor EDS
% Data completeness (in resolution range)	98.1 (54.08-1.83) 98.1 (54.08-1.83)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.12 (at 1.83Å)	Xtriage
Refinement program	Servalcat 0.4.113	Depositor
R, R_{free}	0.171 , 0.217 0.171 , 0.216	Depositor DCC
R_{free} test set	2079 reflections (2.32%)	wwPDB-VP
Wilson B-factor (Å ²)	25.8	Xtriage
Anisotropy	0.340	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 41.1	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.34$	Xtriage
Estimated twinning fraction	0.017 for -h,-k,l	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	13505	wwPDB-VP
Average B, all atoms (Å ²)	33.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.80% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: AGH, CL, EDO

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.86	1/2370 (0.0%)	1.21	8/3223 (0.2%)
1	C	0.86	0/2331	1.21	2/3170 (0.1%)
2	B	0.97	0/874	1.24	4/1184 (0.3%)
2	D	0.91	1/911 (0.1%)	1.20	2/1233 (0.2%)
All	All	0.88	2/6486 (0.0%)	1.21	16/8810 (0.2%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	C	0	1

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	D	52	HIS	CD2-NE2	5.86	1.44	1.37
1	A	79	ARG	NE-CZ	5.13	1.38	1.33

All (16) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	67	GLN	N-CA-CB	6.95	120.33	110.12
1	A	246	ARG	CD-NE-CZ	6.29	133.21	124.40
2	D	46	ARG	CD-NE-CZ	-6.13	115.82	124.40
1	A	114	PHE	CA-CB-CG	6.11	119.91	113.80
2	B	71	PHE	CA-CB-CG	6.01	119.81	113.80
1	A	195	PRO	CB-CA-C	5.96	118.06	111.56
1	C	269	GLU	CB-CG-CD	5.54	122.02	112.60

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	60	ASP	CA-CB-CG	5.54	118.14	112.60
2	B	100	MET	CG-SD-CE	-5.29	89.26	100.90
1	A	183	LYS	N-CA-CB	-5.24	102.34	110.04
1	A	171	SER	CA-C-N	5.21	125.73	120.00
1	A	171	SER	C-N-CA	5.21	125.73	120.00
2	D	49	LYS	CB-CA-C	-5.17	103.96	112.03
2	B	65	LEU	N-CA-CB	-5.13	101.90	110.57
1	A	59	SER	CA-C-O	-5.04	116.25	121.99
1	C	246	ARG	CB-CA-C	-5.04	100.16	109.37

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	C	193	ARG	Sidechain

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2258	2137	2177	10	0
1	C	2244	2116	2169	10	0
2	B	842	789	813	3	0
2	D	862	815	829	1	0
3	A	60	93	99	0	0
3	C	60	93	99	1	0
4	A	8	8	12	0	0
4	B	4	4	6	0	0
4	C	12	12	18	1	0
4	D	4	4	6	0	0
5	B	1	0	0	0	0
5	C	1	0	0	0	0
5	D	1	0	0	0	0
6	A	345	0	0	2	0
6	B	201	0	0	2	0
6	C	348	0	0	2	0
6	D	183	0	0	1	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
All	All	7434	6071	6228	24	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (24) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:135:GLN:HG3	1:A:136[B]:GLU:OE2	2.00	0.60
1:C:197:PRO:HG3	1:C:250:GLU:HG3	1.85	0.57
1:C:246:ARG:HD3	4:C:303:EDO:H21	1.86	0.57
1:A:208:VAL:HG21	1:A:218[A]:VAL:HG21	1.89	0.53
1:C:115[A]:HIS:HE1	6:C:680:HOH:O	1.93	0.52
1:A:68:ARG:NH2	6:A:407:HOH:O	2.43	0.51
2:D:84[B]:ASN:ND2	6:D:311:HOH:O	2.47	0.48
1:A:61:GLN:H	1:A:61:GLN:CD	2.21	0.47
1:A:186:LYS:HA	1:A:267:SER:OG	2.15	0.45
1:C:82:LYS:NZ	6:C:409:HOH:O	2.49	0.45
1:C:197:PRO:CG	1:C:250:GLU:HG3	2.46	0.45
2:B:84[B]:ASN:OD1	2:B:91:PRO:HG3	2.15	0.45
1:A:201:ARG:HD2	1:A:252:ALA:HA	1.98	0.45
1:C:183:LYS:HG2	1:C:184:GLN:N	2.32	0.45
1:C:217:TRP:HB3	1:C:264:LYS:HB2	2.00	0.43
2:B:49:LYS:HB2	6:B:533:HOH:O	2.18	0.42
1:C:66:LEU:HD21	3:C:301:AGH:HAV2	2.00	0.42
1:A:189:ALA:HA	1:A:207:HIS:O	2.19	0.41
1:A:200:GLY:O	1:A:253:ALA:HB2	2.19	0.41
1:A:271:GLN:HG3	6:A:591:HOH:O	2.20	0.41
1:C:177:GLY:O	1:C:178:LYS:C	2.63	0.41
2:B:5[A]:THR:HG22	6:B:598:HOH:O	2.19	0.41
1:C:201:ARG:NH1	1:C:250:GLU:OE2	2.45	0.41
1:A:105:HIS:C	1:A:106:PRO:O	2.64	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	284/296 (96%)	276 (97%)	7 (2%)	1 (0%)	30	18
1	C	280/296 (95%)	274 (98%)	5 (2%)	1 (0%)	30	18
2	B	100/100 (100%)	100 (100%)	0	0	100	100
2	D	105/100 (105%)	104 (99%)	1 (1%)	0	100	100
All	All	769/792 (97%)	754 (98%)	13 (2%)	2 (0%)	36	25

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	106	PRO
1	C	266	SER

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	249/253 (98%)	247 (99%)	2 (1%)	73	65
1	C	244/253 (96%)	239 (98%)	5 (2%)	48	32
2	B	97/95 (102%)	96 (99%)	1 (1%)	68	58
2	D	102/95 (107%)	102 (100%)	0	100	100
All	All	692/696 (99%)	684 (99%)	8 (1%)	63	51

All (8) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	217	TRP
1	A	231	GLN
2	B	71	PHE
1	C	29	LEU
1	C	67	GLN
1	C	143	LEU
1	C	217	TRP
1	C	269	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	152	ASN
1	C	127	GLN
1	C	146	GLN
1	C	184	GLN
1	C	228	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 12 ligands modelled in this entry, 3 are monoatomic - leaving 9 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the

expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	EDO	D	201	-	3,3,3	0.13	0	2,2,2	0.44	0
4	EDO	C	303	-	3,3,3	0.57	0	2,2,2	0.47	0
4	EDO	A	302	-	3,3,3	0.53	0	2,2,2	0.33	0
4	EDO	C	302	-	3,3,3	0.39	0	2,2,2	0.46	0
4	EDO	A	303	-	3,3,3	1.46	1 (33%)	2,2,2	0.41	0
4	EDO	C	304	-	3,3,3	0.48	0	2,2,2	0.19	0
3	AGH	C	301	-	60,60,60	0.83	1 (1%)	65,69,69	0.92	4 (6%)
3	AGH	A	301	-	60,60,60	0.51	0	65,69,69	0.70	1 (1%)
4	EDO	B	402	-	3,3,3	0.30	0	2,2,2	0.29	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	EDO	D	201	-	-	0/1/1/1	-
4	EDO	C	303	-	-	0/1/1/1	-
4	EDO	A	302	-	-	0/1/1/1	-
4	EDO	C	302	-	-	0/1/1/1	-
4	EDO	A	303	-	-	0/1/1/1	-
4	EDO	C	304	-	-	0/1/1/1	-
3	AGH	C	301	-	-	6/58/78/78	0/1/1/1
3	AGH	A	301	-	-	7/58/78/78	0/1/1/1
4	EDO	B	402	-	-	0/1/1/1	-

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	C	301	AGH	O4-C4	2.85	1.49	1.43
4	A	303	EDO	O1-C1	2.27	1.53	1.42

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	301	AGH	C1-C2-N2	3.08	114.15	109.61
3	C	301	AGH	C1-C2-C3	-2.98	106.85	112.71
3	C	301	AGH	C1-C2-N2	2.82	113.77	109.61
3	C	301	AGH	OAA-CAA-N2	-2.70	118.39	122.95
3	C	301	AGH	C6-C5-C4	-2.05	110.81	114.18

There are no chirality outliers.

All (13) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
3	A	301	AGH	CAR-CAS-CAT-CAU
3	A	301	AGH	CAL-CAM-CAN-CAO
3	A	301	AGH	CAI-CAJ-CAK-CAL
3	C	301	AGH	CAL-CAM-CAN-CAO
3	A	301	AGH	CAE-CAF-CAG-CAH
3	C	301	AGH	CAE-CAF-CAG-CAH
3	C	301	AGH	CAI-CAJ-CAK-CAL
3	C	301	AGH	CAN-CAO-CAP-CAQ
3	A	301	AGH	CAU-CAV-CAW-CAX
3	C	301	AGH	CAP-CAQ-CAR-CAS
3	A	301	AGH	CAJ-CAK-CAL-CAM
3	A	301	AGH	CAP-CAQ-CAR-CAS
3	C	301	AGH	C4-C5-C6-C7

There are no ring outliers.

2 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	C	303	EDO	1	0
3	C	301	AGH	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	273/296 (92%)	-0.13	10 (3%) 45 48	12, 28, 68, 99	13 (4%)
1	C	275/296 (92%)	0.01	15 (5%) 30 33	11, 29, 69, 102	7 (2%)
2	B	99/100 (99%)	-0.50	1 (1%) 79 86	12, 24, 43, 58	3 (3%)
2	D	99/100 (99%)	-0.54	2 (2%) 65 72	13, 25, 44, 66	7 (7%)
All	All	746/792 (94%)	-0.18	28 (3%) 44 48	11, 27, 65, 102	30 (4%)

All (28) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	278	GLY	4.8
1	C	278	GLY	4.7
1	C	106	PRO	3.7
1	A	106	PRO	3.7
1	C	109	ALA	3.5
2	D	2	ILE	3.4
1	C	202	LEU	3.4
1	C	199	PRO	3.1
1	C	200	GLY	3.1
1	A	107	GLY	2.8
1	A	199	PRO	2.8
1	A	200	GLY	2.8
1	A	253	ALA	2.5
1	C	198	GLY	2.5
1	C	252	ALA	2.5
1	C	253	ALA	2.5
2	B	2	ILE	2.4
1	A	109	ALA	2.3
1	C	256	ALA	2.3
1	C	195	PRO	2.3
1	A	277	TRP	2.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	198	GLY	2.3
1	C	107	GLY	2.2
1	C	273	ILE	2.2
1	A	257	ALA	2.2
1	C	197	PRO	2.1
1	C	196	SER	2.0
2	D	100	MET	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	AGH	C	301	60/60	0.89	0.11	32,42,56,61	0
4	EDO	B	402	4/4	0.92	0.11	37,38,41,44	0
4	EDO	D	201	4/4	0.93	0.09	35,35,37,42	0
3	AGH	A	301	60/60	0.94	0.08	24,33,43,48	0
5	CL	B	401	1/1	0.95	0.17	44,44,44,44	0
4	EDO	C	303	4/4	0.96	0.07	24,29,33,33	0
4	EDO	A	303	4/4	0.96	0.07	22,29,33,33	0
4	EDO	A	302	4/4	0.96	0.08	28,28,29,34	0
4	EDO	C	302	4/4	0.97	0.07	27,28,28,40	0
4	EDO	C	304	4/4	0.97	0.06	30,33,34,34	0
5	CL	D	202	1/1	0.98	0.15	42,42,42,42	0
5	CL	C	305	1/1	1.00	0.03	26,26,26,26	1

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around AGH C 301:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.5 Other polymers ⓘ

There are no such residues in this entry.