



Full wwPDB X-ray Structure Validation Report ⓘ

Aug 3, 2026 – 01:16 pm BST

PDB ID : 9S6K / pdb_00009s6k
Title : Structure of C. tepidum Rel in an open state in complex with Nb898
Authors : Garcia-Pino, A.; Van Nerom, K.; Talavera Perez, A.
Deposited on : 2025-08-01
Resolution : 3.15 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.015 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.50


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.15 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	2361 (3.20-3.12)
Clashscore	190562	2486 (3.20-3.12)
Ramachandran outliers	187476	2405 (3.20-3.12)
Sidechain outliers	187428	2404 (3.20-3.12)
RSRZ outliers	180081	2361 (3.20-3.12)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	125	6% 72% 18% 10%
1	C	125	6% 81% 10% 9%
2	B	346	3% 73% 17% 7%
2	D	346	4% 75% 15% 8%

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 6261 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Nanobody Nb898.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	113	Total	C	N	O	S	0	0	0
			801	503	137	158	3			
1	C	114	Total	C	N	O	S	0	0	0
			807	507	140	156	4			

- Molecule 2 is a protein called GTP pyrophosphokinase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	322	Total	C	N	O	S	0	0	0
			2367	1512	406	438	11			
2	D	318	Total	C	N	O	S	0	0	0
			2284	1466	395	411	12			

- Molecule 3 is MANGANESE (II) ION (CCD ID: MN) (formula: Mn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	B	1	Total	Mn	1	0
			1	1		
3	D	1	Total	Mn	1	0
			1	1		



4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	197.43Å 88.30Å 62.52Å 90.00° 101.88° 90.00°	Depositor
Resolution (Å)	96.07 – 3.15 96.07 – 3.15	Depositor EDS
% Data completeness (in resolution range)	60.1 (96.07-3.15) 61.3 (96.07-3.15)	Depositor EDS
R_{merge}	0.62	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.75 (at 3.13Å)	Xtriage
Refinement program	PHENIX 2.0_5885	Depositor
R, R_{free}	0.279 , 0.304 0.267 , 0.292	Depositor DCC
R_{free} test set	535 reflections (2.91%)	wwPDB-VP
Wilson B-factor (Å ²)	40.3	Xtriage
Anisotropy	0.303	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 64.2	EDS
L-test for twinning ²	$\langle L \rangle = 0.40$, $\langle L^2 \rangle = 0.23$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.79	EDS
Total number of atoms	6261	wwPDB-VP
Average B, all atoms (Å ²)	37.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.72% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: MN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.62	0/816	0.89	0/1117
1	C	0.64	0/822	0.80	0/1123
2	B	0.65	0/2414	0.99	0/3291
2	D	0.64	0/2327	0.98	0/3172
All	All	0.64	0/6379	0.95	0/8703

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	801	0	743	16	0
1	C	807	0	760	5	0
2	B	2367	0	2154	49	0
2	D	2284	0	2091	32	0
3	B	1	0	0	0	0
3	D	1	0	0	0	0
All	All	6261	0	5748	92	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

All (92) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:16:ILE:HD11	2:B:59:VAL:HG13	1.62	0.82
2:B:223:ASN:HA	2:B:226:ILE:HG13	1.73	0.69
2:B:223:ASN:HA	2:B:226:ILE:CG1	2.22	0.69
2:B:69:LEU:HD13	2:B:143:LEU:HD22	1.75	0.67
2:B:338:MET:O	2:B:339:HIS:C	2.37	0.67
2:D:82:HIS:NE2	2:D:83:ASP:OD1	2.29	0.66
2:B:298:ILE:HG22	2:B:301:HIS:HD2	1.61	0.66
1:C:28:ILE:HG12	1:C:77:ASN:HA	1.77	0.65
1:A:99:ILE:HG21	2:B:285:PHE:HB3	1.78	0.65
1:A:39:GLN:HB2	1:A:45:ARG:HG3	1.79	0.65
2:B:236:GLN:HG3	2:B:290:TYR:CE2	2.34	0.62
1:A:62:VAL:CG2	2:B:236:GLN:HA	2.29	0.62
2:D:84:VAL:O	2:D:88:SER:HB3	2.00	0.62
2:D:223:ASN:HA	2:D:226:ILE:HG12	1.83	0.61
2:D:298:ILE:HG22	2:D:300:GLN:HG2	1.82	0.61
2:B:227:ALA:HB3	2:B:228:PRO:HD3	1.83	0.59
1:A:4:LEU:HD22	1:A:22:CYS:SG	2.42	0.59
1:A:17:SER:HA	1:A:83:MET:O	2.03	0.59
1:C:83:MET:HE1	1:C:111:VAL:HG21	1.83	0.58
2:D:106:ILE:HG23	2:D:141:VAL:HG22	1.85	0.58
2:D:113:ILE:HG23	2:D:132:LEU:HD21	1.85	0.58
1:A:36:TRP:HE1	1:A:79:VAL:HG12	1.68	0.58
2:B:298:ILE:HG22	2:B:301:HIS:CD2	2.38	0.58
2:B:295:PHE:HE1	2:D:181:ARG:HH22	1.47	0.57
1:A:57:SER:HB2	2:B:280:LYS:HG2	1.86	0.57
2:B:236:GLN:HG3	2:B:290:TYR:HE2	1.69	0.57
2:B:335:THR:O	2:B:336:ARG:C	2.45	0.56
2:B:133:LEU:HD21	2:B:182:PHE:HZ	1.72	0.55
2:B:295:PHE:HE1	2:D:181:ARG:NH2	2.04	0.55
2:D:136:VAL:HG13	2:D:137:LYS:N	2.23	0.54
2:B:146:PHE:HB3	2:B:187:MET:HE3	1.90	0.54
2:B:106:ILE:HG23	2:B:141:VAL:HG22	1.89	0.54
2:B:156:LEU:O	2:B:159:LEU:HG	2.09	0.53
2:B:84:VAL:O	2:B:88:SER:HB3	2.09	0.52
2:B:57:HIS:HD1	2:B:151:HIS:CE1	2.27	0.52
2:B:113:ILE:HD11	2:B:136:VAL:HG21	1.92	0.52
1:A:60:TYR:HE1	1:A:70:ILE:HG22	1.74	0.51
2:B:138:ASP:HB3	2:B:141:VAL:HG23	1.92	0.51
2:B:287:VAL:HG12	2:B:333:ILE:HD13	1.93	0.51
1:A:6:GLU:HG2	1:A:95:TYR:HA	1.92	0.50

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:D:74:VAL:HG13	2:D:103:VAL:HG22	1.93	0.49
2:B:296:PRO:HG2	2:B:322:ILE:HB	1.95	0.49
2:D:69:LEU:HD13	2:D:143:LEU:HD22	1.94	0.49
2:D:136:VAL:HG13	2:D:137:LYS:H	1.76	0.49
2:B:43:HIS:HB3	2:B:54:PHE:HB3	1.95	0.49
2:B:221:TYR:OH	2:B:327:HIS:HB2	2.13	0.49
2:B:225:MET:O	2:B:228:PRO:HD2	2.13	0.49
2:D:9:HIS:HA	2:D:55:PHE:CE1	2.48	0.48
2:B:331:LEU:HD21	2:B:333:ILE:HD11	1.96	0.48
1:A:39:GLN:NE2	2:D:135:MET:SD	2.87	0.48
1:A:62:VAL:HG23	2:B:236:GLN:HA	1.96	0.48
1:A:62:VAL:HG23	2:B:236:GLN:OE1	2.14	0.48
2:B:298:ILE:CG2	2:B:301:HIS:CD2	2.96	0.47
1:A:47:LEU:HD21	1:A:50:VAL:HG23	1.96	0.47
2:B:284:CYS:SG	2:B:335:THR:HG23	2.55	0.47
2:B:295:PHE:CE1	2:D:181:ARG:NH2	2.75	0.47
2:B:142:ILE:HG21	2:B:182:PHE:CE2	2.50	0.46
1:C:100:PRO:O	1:C:101:ASN:HB2	2.15	0.46
2:D:12:LYS:HD2	2:D:59:VAL:HG11	1.97	0.46
2:D:227:ALA:HB3	2:D:228:PRO:HD3	1.98	0.45
2:D:246:ALA:O	2:D:247:LYS:C	2.59	0.45
2:B:233:LEU:HD22	2:B:287:VAL:HG22	1.98	0.45
2:D:110:LEU:HG	2:D:141:VAL:HG13	1.99	0.45
2:B:223:ASN:HA	2:B:226:ILE:HG12	1.97	0.45
1:A:40:ALA:HB3	1:A:43:LYS:HB2	1.98	0.44
2:D:216:ASN:HA	2:D:219:VAL:HG22	2.00	0.44
1:C:36:TRP:HE1	1:C:79:VAL:HG12	1.83	0.43
2:D:180:HIS:HB2	2:D:188:LYS:NZ	2.34	0.43
1:A:28:ILE:O	1:A:72:ARG:NH2	2.52	0.43
2:B:74:VAL:HG13	2:B:103:VAL:HG22	2.01	0.42
2:D:9:HIS:HA	2:D:55:PHE:HE1	1.84	0.42
2:D:314:TYR:OH	2:D:332:GLN:OE1	2.34	0.42
2:D:160:PRO:HD2	2:D:163:ARG:HD2	2.01	0.42
2:B:4:GLN:HB2	2:B:41:ARG:HH12	1.83	0.42
2:D:65:LEU:HD12	2:D:69:LEU:HB2	2.01	0.42
2:B:13:LEU:HB2	2:B:36:PHE:CE2	2.55	0.42
2:B:142:ILE:HG21	2:B:182:PHE:HE2	1.84	0.42
2:B:161:GLU:O	2:B:162:HIS:C	2.62	0.42
2:D:289:GLY:O	2:D:293:GLN:HG3	2.19	0.42
2:D:233:LEU:HD22	2:D:287:VAL:HG22	2.01	0.42
2:D:142:ILE:HG22	2:D:146:PHE:CE2	2.55	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:57:HIS:CE1	2:B:61:VAL:HG11	2.55	0.41
2:B:226:ILE:HG22	2:B:242:LEU:HD12	2.02	0.41
1:C:40:ALA:HB3	1:C:43:LYS:HB2	2.02	0.41
2:B:302:PHE:CZ	2:B:317:LEU:HD22	2.56	0.41
2:D:136:VAL:HG21	2:D:141:VAL:HB	2.02	0.41
2:B:275:ILE:HG12	2:B:334:ARG:HG3	2.02	0.41
2:D:114:SER:C	2:D:116:ILE:N	2.77	0.41
2:D:231:ASP:O	2:D:234:GLU:HG3	2.21	0.41
2:D:303:LYS:HB2	2:D:318:HIS:HB3	2.02	0.41
1:A:62:VAL:HG21	2:B:236:GLN:HA	2.04	0.40
2:B:43:HIS:HB2	2:B:55:PHE:HB3	2.04	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	111/125 (89%)	109 (98%)	2 (2%)	0	100	100
1	C	112/125 (90%)	110 (98%)	2 (2%)	0	100	100
2	B	316/346 (91%)	308 (98%)	8 (2%)	0	100	100
2	D	312/346 (90%)	299 (96%)	13 (4%)	0	100	100
All	All	851/942 (90%)	826 (97%)	25 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	77/102 (76%)	75 (97%)	2 (3%)	40	64
1	C	78/102 (76%)	75 (96%)	3 (4%)	29	58
2	B	220/306 (72%)	206 (94%)	14 (6%)	16	43
2	D	206/306 (67%)	192 (93%)	14 (7%)	14	41
All	All	581/816 (71%)	548 (94%)	33 (6%)	18	47

All (33) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	48	VAL
1	A	103	THR
2	B	41	ARG
2	B	61	VAL
2	B	84	VAL
2	B	159	LEU
2	B	162	HIS
2	B	202	GLU
2	B	248	HIS
2	B	249	LEU
2	B	268	ASP
2	B	276	ILE
2	B	280	LYS
2	B	332	GLN
2	B	335	THR
2	B	339	HIS
1	C	7	SER
1	C	57	SER
1	C	103	THR
2	D	10	TYR
2	D	30	SER
2	D	83	ASP
2	D	130	LYS
2	D	181	ARG
2	D	215	ARG
2	D	231	ASP
2	D	234	GLU
2	D	268	ASP
2	D	276	ILE
2	D	281	ILE

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
2	D	300	GLN
2	D	312	ASN
2	D	314	TYR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (7) such sidechains are listed below:

Mol	Chain	Res	Type
2	B	223	ASN
2	B	301	HIS
1	C	32	ASN
2	D	14	HIS
2	D	243	GLN
2	D	300	GLN
2	D	315	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry ⓘ

Of 2 ligands modelled in this entry, 2 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2			OWAB(Å²)	Q<0.9
1	A	113/125 (90%)	0.69	8 (7%)	22	12	27, 41, 48, 52	0
1	C	114/125 (91%)	0.66	8 (7%)	22	13	21, 34, 44, 50	0
2	B	322/346 (93%)	0.54	10 (3%)	51	31	18, 35, 48, 60	0
2	D	318/346 (91%)	0.77	15 (4%)	36	21	20, 39, 51, 59	0
All	All	867/942 (92%)	0.66	41 (4%)	36	21	18, 37, 49, 60	0

All (41) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	102	PHE	5.1
2	D	299	PRO	4.7
2	B	268	ASP	4.1
1	C	74	ASP	4.1
2	D	248	HIS	3.9
1	A	17	SER	3.9
1	A	99	ILE	3.7
2	D	297	PRO	3.6
2	D	207	LEU	3.5
2	B	249	LEU	3.1
1	A	102	PHE	3.0
2	D	298	ILE	2.9
2	B	315	GLN	2.8
2	B	183	GLY	2.8
2	B	162	HIS	2.8
1	A	114	SER	2.7
2	D	22	ALA	2.6
2	D	247	LYS	2.6
1	C	99	ILE	2.6
1	A	86	LEU	2.6
1	C	100	PRO	2.5

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
2	B	54	PHE	2.5
1	C	56	GLY	2.4
2	D	139	ILE	2.4
1	C	71	SER	2.4
1	C	85	SER	2.4
2	D	253	TYR	2.3
2	B	312	ASN	2.3
2	B	275	ILE	2.3
2	B	267	HIS	2.3
2	B	314	TYR	2.3
2	D	234	GLU	2.2
2	D	45	GLY	2.2
2	D	109	GLY	2.2
1	A	57	SER	2.2
1	C	21	SER	2.1
2	D	336	ARG	2.1
1	A	94	TYR	2.1
2	D	8	GLU	2.1
2	D	254	ASN	2.0
1	A	100	PRO	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	MN	B	401	1/1	-	-	43,43,43,43	1
3	MN	D	401	1/1	-	-	31,31,31,31	1

6.5 Other polymers [i](#)

There are no such residues in this entry.