



Full wwPDB X-ray Structure Validation Report ⓘ

Aug 3, 2026 – 01:32 pm BST

PDB ID : 9S6O / pdb_00009s6o
Title : HIV-1 capsid (M-group) - native in complex with JW3-076
Authors : Govasli, M.A.L.; Pinotsis, N.; Towers, G.; Selwood, D.; Jacques, D.A.
Deposited on : 2025-08-01
Resolution : 2.09 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	1.8.4, CSD as541be (2020)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.015 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.50


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.09 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	8172 (2.10-2.06)
Clashscore	190562	8714 (2.10-2.06)
Ramachandran outliers	187476	8641 (2.10-2.06)
Sidechain outliers	187428	8642 (2.10-2.06)
RSRZ outliers	180081	8177 (2.10-2.06)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	231	19% 71% 22% ...

2 Entry composition [i](#)


There are 6 unique types of molecules in this entry. The entry contains 1822 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Gag polypeptide.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	221	Total	C	N	O	S	0	1	0
			1728	1090	303	322	13			

- Molecule 2 is (2 {S})-3-[3,5-difluorophenyl]ethyl- {N}-(4-methoxyphenyl)- {N}-methyl-2-[2-(2-methyl-1 {H}-indol-3-yl)ethanoylamino]propanamide (CCD ID: A1JL3) (formula: C₂₈H₂₇F₂N₃O₃) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	F	N	O	0	0
			36	28	2	3	3		


- Molecule 3 is IODIDE ION (CCD ID: IOD) (formula: I).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	2	Total	I	0	0
			2	2		

- Molecule 4 is CHLORIDE ION (CCD ID: CL) (formula: Cl).

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	4	Total	Cl		0	0
			4	4			

- Molecule 5 is BETA-MERCAPTOETHANOL (CCD ID: BME) (formula: C₂H₆OS).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
5	A	1	Total	C	O	S	0	0
			4	2	1	1		


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	48	Total	O	0	0
			48	48		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Gag polypeptide



4 Data and refinement statistics

Property	Value	Source
Space group	P 6	Depositor
Cell constants a, b, c, α , β , γ	92.71Å 92.71Å 57.36Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	57.36 – 2.09 57.36 – 2.09	Depositor EDS
% Data completeness (in resolution range)	76.2 (57.36-2.09) 76.2 (57.36-2.09)	Depositor EDS
R_{merge}	0.09	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.68 (at 2.08Å)	Xtriage
Refinement program	BUSTER 2.10.3	Depositor
R, R_{free}	0.224 , 0.249 0.225 , 0.270	Depositor DCC
R_{free} test set	643 reflections (3.82%)	wwPDB-VP
Wilson B-factor (Å ²)	57.7	Xtriage
Anisotropy	0.081	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 54.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	0.052 for h,-h-k,-l	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	1822	wwPDB-VP
Average B, all atoms (Å ²)	70.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.12% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality ⓘ

5.1 Standard geometry ⓘ

Bond lengths and bond angles in the following residue types are not validated in this section: CL, A1JL3, IOD, BME

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.27	8/1767 (0.5%)	1.36	18/2402 (0.7%)

All (8) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	68	MET	SD-CE	7.23	1.97	1.79
1	A	96	MET	SD-CE	6.39	1.95	1.79
1	A	31	ALA	CA-C	6.20	1.61	1.52
1	A	135	ILE	CG1-CD1	-5.88	1.28	1.51
1	A	134	ILE	CG1-CD1	5.80	1.74	1.51
1	A	122	PRO	CA-C	5.79	1.57	1.52
1	A	214	MET	SD-CE	-5.61	1.65	1.79
1	A	186	THR	CA-C	5.18	1.59	1.52

All (18) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	121	ASN	CA-CB-CG	6.63	119.23	112.60
1	A	36	VAL	CA-C-N	6.34	125.40	120.33
1	A	36	VAL	C-N-CA	6.34	125.40	120.33
1	A	185	MET	CA-C-N	6.32	128.66	120.44
1	A	185	MET	C-N-CA	6.32	128.66	120.44
1	A	195	ASN	CA-CB-CG	6.00	118.60	112.60
1	A	197	ASP	CA-CB-CG	5.83	118.43	112.60
1	A	148	THR	CA-C-N	5.77	132.08	122.73
1	A	148	THR	C-N-CA	5.77	132.08	122.73
1	A	5	ASN	CA-CB-CG	5.66	118.26	112.60
1	A	26	VAL	N-CA-CB	5.57	118.93	110.58
1	A	144	MET	CB-CG-SD	-5.48	96.26	112.70
1	A	198	CYS	CA-C-N	5.30	128.12	120.38

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	198	CYS	C-N-CA	5.30	128.12	120.38
1	A	122	PRO	N-CA-C	-5.24	104.31	110.70
1	A	218	CYS	CA-C-N	5.08	127.54	120.63
1	A	218	CYS	C-N-CA	5.08	127.54	120.63
1	A	33	SER	N-CA-C	-5.00	102.89	110.39

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1728	0	1727	31	0
2	A	36	0	0	1	0
3	A	2	0	0	1	0
4	A	4	0	0	0	0
5	A	4	0	6	0	0
6	A	48	0	0	0	0
All	All	1822	0	1733	31	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 9.

All (31) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:117:TRP:CD1	1:A:124:ILE:HD13	2.12	0.83
1:A:86:VAL:HG23	1:A:98:GLU:HB2	1.60	0.81
1:A:117:TRP:HD1	1:A:124:ILE:HD13	1.44	0.81
1:A:176:GLN:HG2	1:A:176:GLN:O	1.89	0.70
1:A:143:ARG:HD2	1:A:176:GLN:HB3	1.75	0.66
1:A:48:THR:HG22	1:A:118:MET:SD	2.36	0.66
1:A:143:ARG:HD3	1:A:176:GLN:HA	1.80	0.64
1:A:54:THR:O	1:A:58:THR:HG23	1.98	0.63
1:A:199:LYS:O	1:A:203:LYS:HG3	2.00	0.62

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:121:ASN:CG	1:A:122:PRO:HD3	2.28	0.58
1:A:143:ARG:HD2	1:A:176:GLN:CB	2.34	0.57
1:A:108:THR:O	1:A:108:THR:HG22	2.05	0.56
1:A:170:LYS:HG3	3:A:602:IOD:I	2.79	0.53
1:A:155:GLN:HG3	1:A:164:TYR:HB2	1.90	0.53
1:A:143:ARG:CD	1:A:176:GLN:HB3	2.39	0.52
1:A:96:MET:SD	1:A:116:GLY:O	2.68	0.52
1:A:151:LEU:HD23	1:A:189:LEU:HD11	1.93	0.51
1:A:97[A]:ARG:HD3	1:A:98:GLU:O	2.11	0.50
1:A:33:SER:HB2	1:A:35:GLU:OE1	2.13	0.49
1:A:117:TRP:HD1	1:A:124:ILE:CD1	2.19	0.48
1:A:35:GLU:C	1:A:38:PRO:HD2	2.38	0.48
1:A:143:ARG:HD3	1:A:176:GLN:CA	2.43	0.48
1:A:56:LEU:HD22	1:A:66:MET:HE1	1.97	0.46
1:A:77:ALA:HB1	1:A:101:GLY:HA3	1.97	0.46
1:A:143:ARG:CD	1:A:176:GLN:CB	2.94	0.45
1:A:66:MET:HE3	2:A:601:A1JL3:C19	2.47	0.44
1:A:92:ALA:HB3	1:A:95:GLN:HG3	2.01	0.42
1:A:1:PRO:N	1:A:13:GLN:O	2.52	0.41
1:A:108:THR:O	1:A:108:THR:CG2	2.67	0.41
1:A:144:MET:HE3	1:A:144:MET:HB2	1.85	0.40
1:A:202:LEU:HD22	1:A:214:MET:HG2	2.03	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	220/231 (95%)	217 (99%)	2 (1%)	1 (0%)	24 21

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	122	PRO

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	187/194 (96%)	176 (94%)	11 (6%)	18 15

All (11) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	6	LEU
1	A	7	GLN
1	A	33	SER
1	A	50	GLN
1	A	86	VAL
1	A	112	GLN
1	A	119	THR
1	A	121	ASN
1	A	124	ILE
1	A	180	GLU
1	A	221	VAL

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	7	GLN
1	A	21	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 8 ligands modelled in this entry, 6 are monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
5	BME	A	608	-	3,3,3	0.58	0	1,2,2	2.78	1 (100%)
2	A1JL3	A	601	-	38,39,39	1.82	9 (23%)	51,55,55	1.91	12 (23%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	BME	A	608	-	-	0/1/1/1	-
2	A1JL3	A	601	-	-	0/26/26/26	0/4/4/4

All (9) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	601	A1JL3	C24-N23	5.99	1.46	1.34
2	A	601	A1JL3	C11-N09	4.27	1.42	1.35
2	A	601	A1JL3	C28-N30	-4.21	1.31	1.37
2	A	601	A1JL3	O12-C11	-3.27	1.16	1.22
2	A	601	A1JL3	C36-C31	-3.01	1.37	1.41
2	A	601	A1JL3	O02-C03	2.20	1.42	1.37

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	601	A1JL3	C35-C36	2.19	1.43	1.39
2	A	601	A1JL3	O25-C24	-2.19	1.18	1.23
2	A	601	A1JL3	C32-C31	2.15	1.43	1.39

All (13) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	601	A1JL3	C06-N09-C11	-5.35	116.42	124.43
2	A	601	A1JL3	C07-C06-N09	-4.96	113.63	120.10
2	A	601	A1JL3	C10-N09-C06	4.51	124.60	117.02
2	A	601	A1JL3	O12-C11-N09	-3.60	116.71	121.44
2	A	601	A1JL3	C29-C28-N30	3.36	123.62	120.03
2	A	601	A1JL3	C32-C31-C36	-3.36	118.82	122.19
5	A	608	BME	O1-C1-C2	-2.78	99.86	110.83
2	A	601	A1JL3	C19-C20-C22	-2.76	120.03	123.52
2	A	601	A1JL3	C13-N23-C24	2.44	127.93	121.65
2	A	601	A1JL3	C05-C06-N09	2.30	123.10	120.10
2	A	601	A1JL3	O25-C24-C26	-2.22	117.71	121.33
2	A	601	A1JL3	C20-C19-C17	2.14	119.45	116.13
2	A	601	A1JL3	C07-C06-C05	2.01	123.26	119.16

There are no chirality outliers.


There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	601	A1JL3	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	221/231 (95%)	1.19	43 (19%) 3 3	37, 67, 113, 143	1 (0%)

All (43) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	2	ILE	6.1
1	A	97[A]	ARG	5.5
1	A	6	LEU	5.3
1	A	3	VAL	4.5
1	A	1	PRO	4.1
1	A	221	VAL	4.0
1	A	32	PHE	4.0
1	A	4	GLN	3.7
1	A	8	GLY	3.6
1	A	191	VAL	3.3
1	A	37	ILE	3.2
1	A	145	TYR	3.1
1	A	141	ILE	3.1
1	A	108	THR	2.9
1	A	10	MET	2.8
1	A	119	THR	2.8
1	A	220	GLY	2.8
1	A	5	ASN	2.8
1	A	39	MET	2.8
1	A	69	LEU	2.7
1	A	12	HIS	2.7
1	A	11	VAL	2.7
1	A	136	LEU	2.6
1	A	138	LEU	2.6
1	A	207	PRO	2.4
1	A	217	ALA	2.4
1	A	202	LEU	2.4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	184	TRP	2.4
1	A	161	PHE	2.3
1	A	94	GLY	2.2
1	A	190	LEU	2.2
1	A	13	GLN	2.2
1	A	62	HIS	2.2
1	A	126	VAL	2.2
1	A	29	GLU	2.2
1	A	68	MET	2.2
1	A	54	THR	2.1
1	A	205	LEU	2.1
1	A	18	ARG	2.1
1	A	15	ILE	2.1
1	A	150	ILE	2.1
1	A	26	VAL	2.1
1	A	168	PHE	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
4	CL	A	606	1/1	0.77	0.18	88,88,88,88	0
4	CL	A	605	1/1	0.87	0.17	69,69,69,69	0
4	CL	A	607	1/1	0.93	0.17	83,83,83,83	0
2	A1JL3	A	601	36/36	0.94	0.08	41,50,56,58	0
5	BME	A	608	4/4	0.94	0.12	54,57,58,60	0
3	IOD	A	603	1/1	0.98	0.04	75,75,75,75	0
3	IOD	A	602	1/1	0.99	0.04	63,63,63,63	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
4	CL	A	604	1/1	0.99	0.09	49,49,49,49	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



6.5 Other polymers ⓘ

There are no such residues in this entry.