



Full wwPDB X-ray Structure Validation Report ⓘ

Aug 29, 2026 – 07:21 pm BST

PDB ID : 9SP3 / pdb_00009sp3
Title : SARS-CoV-2 nsp16-nsp10 in complex with SAM derivative inhibitor 1196
Authors : Kalnins, G.
Deposited on : 2025-09-16
Resolution : 1.70 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 1.8.4, CSD as541be (2020)
Xtriage (Phenix) : 2.0
EDS : 3.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.015 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.50


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	5551 (1.70-1.70)
Clashscore	190562	5924 (1.70-1.70)
Ramachandran outliers	187476	5846 (1.70-1.70)
Sidechain outliers	187428	5846 (1.70-1.70)
RSRZ outliers	180081	5554 (1.70-1.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	302	11% 91% 7%
2	B	142	20% 78% 20%

2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 3549 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called 2'-O-methyltransferase nsp16.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	297	2345	1498	391	438	18	0	4	0

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-3	GLY	-	expression tag	UNP P0DTD1
A	-2	SER	-	expression tag	UNP P0DTD1
A	-1	MET	-	expression tag	UNP P0DTD1
A	0	ALA	-	expression tag	UNP P0DTD1


- Molecule 2 is a protein called Non-structural protein 10.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	B	113	842	526	142	158	16	0	2	0

There are 4 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
B	-2	GLY	-	expression tag	UNP P0DTC1
B	-1	SER	-	expression tag	UNP P0DTC1
B	0	MET	-	expression tag	UNP P0DTC1
B	113	ARG	LYS	conflict	UNP P0DTC1

- Molecule 3 is 4-[4-[(2 {R},3 {S},4 {R},5 {R})-5-(6-aminopurin-9-yl)-3,4-bis(oxidanyl)oxolan-2-yl]-1,2,3-triazol-1-yl]butanoic acid (CCD ID: A1JPE) (formula: C₁₅H₁₈N₈O₅) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C	N	O	0	0
			28	15	8	5		

- Molecule 4 is 2-(N-MORPHOLINO)-ETHANESULFONIC ACID (CCD ID: MES) (formula: $C_6H_{13}NO_4S$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
4	A	1	Total	C	N	O	S	0	0
			12	6	1	4	1		

- Molecule 5 is ZINC ION (CCD ID: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	B	2	Total 2	Zn 2	0	0


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	246	Total 246	O 246	0	0
6	B	74	Total 74	O 74	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: 2'-O-methyltransferase nsp16



• Molecule 2: Non-structural protein 10



4 Data and refinement statistics

Property	Value	Source
Space group	P 31 2 1	Depositor
Cell constants a, b, c, α , β , γ	168.22Å 168.22Å 52.03Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	42.34 – 1.70 42.34 – 1.70	Depositor EDS
% Data completeness (in resolution range)	99.8 (42.34-1.70) 99.8 (42.34-1.70)	Depositor EDS
R_{merge}	0.20	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.62 (at 1.70Å)	Xtriage
Refinement program	REFMAC 5.8.0267, PHENIX 1.17.1_3660	Depositor
R, R_{free}	0.209 , 0.222 0.213 , 0.226	Depositor DCC
R_{free} test set	4599 reflections (4.96%)	wwPDB-VP
Wilson B-factor (Å ²)	19.4	Xtriage
Anisotropy	0.056	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 36.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.34$	Xtriage
Estimated twinning fraction	0.027 for -h,-k,l	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	3549	wwPDB-VP
Average B, all atoms (Å ²)	28.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.02% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: A1JPE, ZN, MES

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.32	0/2408	0.54	0/3269
2	B	0.26	0/867	0.49	0/1176
All	All	0.31	0/3275	0.53	0/4445

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2345	0	2333	17	0
2	B	842	0	811	1	0
3	A	28	0	0	1	0
4	A	12	0	12	0	0
5	B	2	0	0	0	0
6	A	246	0	0	5	0
6	B	74	0	0	1	0
All	All	3549	0	3156	18	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (18) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:141:LYS:HG2	1:A:142:GLU:H	1.59	0.66
1:A:268:ASN:HB2	1:A:271:ILE:H	1.62	0.65
1:A:260:MET:HE2	1:A:262:LEU:HD21	1.83	0.61
2:B:75:LEU:HD22	2:B:112:LEU:HD11	1.84	0.59
1:A:141:LYS:HG2	1:A:142:GLU:N	2.18	0.59
1:A:214:LYS:NZ	6:A:402:HOH:O	2.32	0.53
1:A:41[A]:MET:HG3	6:B:312:HOH:O	2.12	0.50
1:A:163:LEU:HB2	1:A:282:ILE:HB	1.93	0.49
1:A:254:LEU:HD13	6:A:535:HOH:O	2.13	0.48
1:A:147[A]:GLU:OE2	6:A:401:HOH:O	2.20	0.45
1:A:148:GLY:HA3	6:A:567:HOH:O	2.17	0.44
1:A:122:ASN:OD1	1:A:288:VAL:HG13	2.17	0.44
1:A:182:LYS:NZ	6:A:414:HOH:O	2.52	0.43
1:A:66:ARG:NH1	1:A:122:ASN:HB2	2.34	0.42
1:A:99:ASP:OD2	3:A:301:A1JPE:O3	2.38	0.41
1:A:76:LYS:HD3	1:A:76:LYS:HA	1.70	0.41
1:A:41[A]:MET:HE1	1:A:80:PRO:HG2	2.02	0.41
1:A:130:ASP:OD1	1:A:170:LYS:HD2	2.20	0.41

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	299/302 (99%)	290 (97%)	9 (3%)	0	100	100
2	B	113/142 (80%)	111 (98%)	2 (2%)	0	100	100
All	All	412/444 (93%)	401 (97%)	11 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	259/260 (100%)	259 (100%)	0	100	100
2	B	93/115 (81%)	93 (100%)	0	100	100
All	All	352/375 (94%)	352 (100%)	0	100	100

There are no protein residues with a non-rotameric sidechain to report.

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	18	GLN
1	A	28	GLN
1	A	29	ASN
1	A	174	HIS
1	A	235	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 4 ligands modelled in this entry, 2 are monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
3	A1JPE	A	301	-	31,31,31	1.68	4 (12%)	41,45,45	2.20	13 (31%)
4	MES	A	302	-	12,12,12	1.86	1 (8%)	14,16,16	2.03	3 (21%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	A1JPE	A	301	-	-	4/12/30/30	0/4/4/4
4	MES	A	302	-	-	1/6/14/14	0/1/1/1

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	302	MES	C8-S	-6.06	1.68	1.77
3	A	301	A1JPE	N3-N2	6.05	1.42	1.32
3	A	301	A1JPE	N1-N2	4.18	1.41	1.34
3	A	301	A1JPE	C11-N5	2.80	1.36	1.31
3	A	301	A1JPE	C6-C5	2.39	1.39	1.36

All (16) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	301	A1JPE	C12-C13-N6	-6.11	118.78	126.75
4	A	302	MES	C5-N4-C3	4.49	118.94	108.83
3	A	301	A1JPE	C5-N3-N2	-4.41	105.58	109.24
3	A	301	A1JPE	N7-C14-N6	-3.63	122.93	128.60
3	A	301	A1JPE	O5-C10-N4	3.58	115.11	108.06
3	A	301	A1JPE	N6-C13-N4	3.56	132.94	127.08
3	A	301	A1JPE	C6-C5-N3	3.55	112.09	107.97
4	A	302	MES	O2S-S-C8	3.46	111.08	106.92
3	A	301	A1JPE	C14-N6-C13	3.45	119.90	111.75
3	A	301	A1JPE	N4-C11-N5	-3.35	109.34	113.91

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	301	A1JPE	C13-C12-N5	-3.22	106.70	110.62
4	A	302	MES	C7-N4-C5	3.18	119.36	111.23
3	A	301	A1JPE	C12-N5-C11	3.17	108.01	103.51
3	A	301	A1JPE	C9-C8-C7	2.71	104.79	101.64
3	A	301	A1JPE	C12-C13-N4	2.14	108.27	105.78
3	A	301	A1JPE	C15-C12-C13	2.02	119.89	117.18

There are no chirality outliers.

All (5) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
3	A	301	A1JPE	C6-C5-C7-O5
4	A	302	MES	C8-C7-N4-C5
3	A	301	A1JPE	O1-C1-C2-C3
3	A	301	A1JPE	O2-C1-C2-C3
3	A	301	A1JPE	C6-C5-C7-C8

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	301	A1JPE	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å ²)	Q<0.9
1	A	297/302 (98%)	0.40	33 (11%)	10 10	11, 22, 50, 93	4 (1%)
2	B	113/142 (79%)	1.11	28 (24%)	2 1	15, 31, 53, 66	2 (1%)
All	All	410/444 (92%)	0.59	61 (14%)	5 5	11, 24, 51, 93	6 (1%)

All (61) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	1	SER	7.7
2	B	18	ALA	5.5
1	A	297	ASN	5.0
1	A	268	ASN	4.6
1	A	254	LEU	4.6
1	A	139	VAL	4.5
1	A	30	TYR	4.5
2	B	86	PRO	4.5
2	B	21	VAL	4.3
1	A	264	GLU	4.2
1	A	140	THR	4.2
2	B	119	VAL	4.0
2	B	89	PHE	4.0
1	A	262	LEU	3.8
1	A	2	SER	3.7
1	A	265	GLY	3.7
2	B	130	CYS	3.7
1	A	263	LYS	3.6
1	A	141	LYS	3.5
1	A	138	ASN	3.2
1	A	137	LYS	3.1
2	B	33	SER	3.1
1	A	31	GLY	3.1
2	B	84	PRO	3.0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
2	B	87	LYS	2.9
1	A	97	ASP	2.9
1	A	261	SER	2.9
2	B	36	GLN	2.8
1	A	287	ARG	2.8
2	B	88	GLY	2.7
2	B	63[A]	MET	2.7
2	B	85	ASN	2.7
2	B	91	ASP	2.7
2	B	22	ASP	2.7
2	B	113	ARG	2.6
1	A	266	GLN	2.6
2	B	20	ALA	2.6
1	A	267	ILE	2.6
2	B	32	ALA	2.6
2	B	115	THR	2.5
2	B	30	TYR	2.4
1	A	269	ASP	2.4
1	A	144	ASP	2.4
2	B	64	ASP	2.4
2	B	118	THR	2.4
1	A	271	ILE	2.4
2	B	19	PHE	2.3
2	B	110	PHE	2.3
2	B	25	LYS	2.3
2	B	24	ALA	2.2
1	A	104	VAL	2.2
1	A	63	TYR	2.2
1	A	217	GLU	2.1
1	A	295	LEU	2.1
2	B	117	CYS	2.1
2	B	120	CYS	2.1
1	A	64	ASN	2.1
1	A	136	THR	2.1
1	A	135	LYS	2.1
1	A	285	ASN	2.0
1	A	32	ASP	2.0

6.2 Non-standard residues in protein, DNA, RNA chains

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)


There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	A1JPE	A	301	28/28	0.68	0.21	31,44,63,65	0
4	MES	A	302	12/12	0.87	0.14	22,27,47,52	0
5	ZN	B	202	1/1	0.93	0.08	46,46,46,46	0
5	ZN	B	201	1/1	0.97	0.05	26,26,26,26	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



6.5 Other polymers [i](#)

There are no such residues in this entry.