



Full wwPDB X-ray Structure Validation Report ⓘ

Aug 17, 2026 – 01:57 pm BST

PDB ID : 9SWH / pdb_00009swh
Title : Crystal structure of aldo-keto reductase 1C3 (AKR1C3) in complex with NADP+ and Myxoglucamide A
Authors : Siemon, T.; Broenstrup, M.; Blankenfeldt, W.
Deposited on : 2025-10-06
Resolution : 1.99 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 1.8.4, CSD as541be (2020)
Xtriage (Phenix) : 2.0
EDS : 3.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.015 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.50


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.99 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	10052 (2.00-2.00)
Clashscore	190562	11152 (2.00-2.00)
Ramachandran outliers	187476	11031 (2.00-2.00)
Sidechain outliers	187428	11029 (2.00-2.00)
RSRZ outliers	180081	10067 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	344	7% 85% 7% 8%
1	B	344	8% 85% 8% 8%

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 10762 atoms, of which 5281 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Aldo-keto reductase family 1 member C3.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	318	Total	C	H	N	O	S	0	0	0
			5104	1627	2551	443	471	12			
1	B	318	Total	C	H	N	O	S	0	0	0
			5103	1627	2550	443	471	12			

There are 42 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	-20	MET	-	initiating methionine	UNP P42330
A	-19	GLY	-	expression tag	UNP P42330
A	-18	SER	-	expression tag	UNP P42330
A	-17	SER	-	expression tag	UNP P42330
A	-16	HIS	-	expression tag	UNP P42330
A	-15	HIS	-	expression tag	UNP P42330
A	-14	HIS	-	expression tag	UNP P42330
A	-13	HIS	-	expression tag	UNP P42330
A	-12	HIS	-	expression tag	UNP P42330
A	-11	HIS	-	expression tag	UNP P42330
A	-10	SER	-	expression tag	UNP P42330
A	-9	SER	-	expression tag	UNP P42330
A	-8	GLY	-	expression tag	UNP P42330
A	-7	GLU	-	expression tag	UNP P42330
A	-6	ASN	-	expression tag	UNP P42330
A	-5	LEU	-	expression tag	UNP P42330
A	-4	TYR	-	expression tag	UNP P42330
A	-3	PHE	-	expression tag	UNP P42330
A	-2	GLN	-	expression tag	UNP P42330
A	-1	GLY	-	expression tag	UNP P42330
A	0	HIS	-	expression tag	UNP P42330
B	-20	MET	-	initiating methionine	UNP P42330
B	-19	GLY	-	expression tag	UNP P42330
B	-18	SER	-	expression tag	UNP P42330
B	-17	SER	-	expression tag	UNP P42330

Continued on next page...

Continued from previous page...


Chain	Residue	Modelled	Actual	Comment	Reference
B	-16	HIS	-	expression tag	UNP P42330
B	-15	HIS	-	expression tag	UNP P42330
B	-14	HIS	-	expression tag	UNP P42330
B	-13	HIS	-	expression tag	UNP P42330
B	-12	HIS	-	expression tag	UNP P42330
B	-11	HIS	-	expression tag	UNP P42330
B	-10	SER	-	expression tag	UNP P42330
B	-9	SER	-	expression tag	UNP P42330
B	-8	GLY	-	expression tag	UNP P42330
B	-7	GLU	-	expression tag	UNP P42330
B	-6	ASN	-	expression tag	UNP P42330
B	-5	LEU	-	expression tag	UNP P42330
B	-4	TYR	-	expression tag	UNP P42330
B	-3	PHE	-	expression tag	UNP P42330
B	-2	GLN	-	expression tag	UNP P42330
B	-1	GLY	-	expression tag	UNP P42330
B	0	HIS	-	expression tag	UNP P42330

- Molecule 2 is Myxoglucamide A (CCD ID: A1JRA) (formula: $C_{32}H_{58}N_2O_{11}$) (labeled as "Ligand of Interest" by depositor).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	H	N	O	0	0
			100	32	55	2	11		
2	B	1	Total	C	H	N	O	0	0
			100	32	55	2	11		

- Molecule 3 is NADP NICOTINAMIDE-ADENINE-DINUCLEOTIDE PHOSPHATE (CCD ID: NAP) (formula: C₂₁H₂₈N₇O₁₇P₃).



Mol	Chain	Residues	Atoms						ZeroOcc	AltConf
3	A	1	Total	C	H	N	O	P	0	0
			73	21	25	7	17	3		
3	B	1	Total	C	H	N	O	P	0	0
			73	21	25	7	17	3		

- Molecule 4 is (R,R)-2,3-BUTANEDIOL (CCD ID: BU3) (formula: C₄H₁₀O₂).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	B	1	Total	C	H	O	0	0
			16	4	10	2		
4	B	1	Total	C	H	O	0	0
			16	4	10	2		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	83	Total	O	0	0
			83	83		
5	B	94	Total	O	0	0
			94	94		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Aldo-keto reductase family 1 member C3



- Molecule 1: Aldo-keto reductase family 1 member C3



4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, α , β , γ	41.59Å 52.93Å 76.09Å 77.23° 85.03° 73.60°	Depositor
Resolution (Å)	49.69 – 1.99 49.69 – 1.99	Depositor EDS
% Data completeness (in resolution range)	69.1 (49.69-1.99) 69.1 (49.69-1.99)	Depositor EDS
R_{merge}	0.21	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.53 (at 2.00Å)	Xtriage
Refinement program	PHENIX V1.20.1-4487	Depositor
R, R_{free}	0.236 , 0.286 0.236 , 0.286	Depositor DCC
R_{free} test set	1379 reflections (3.32%)	wwPDB-VP
Wilson B-factor (Å ²)	16.7	Xtriage
Anisotropy	0.547	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 21.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.89	EDS
Total number of atoms	10762	wwPDB-VP
Average B, all atoms (Å ²)	20.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 8.27% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: NAP, A1JRA, BU3

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.28	0/2612	0.43	1/3535 (0.0%)
1	B	0.33	1/2612 (0.0%)	0.45	0/3535
All	All	0.30	1/5224 (0.0%)	0.44	1/7070 (0.0%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	121	SER	C-O	-6.17	1.16	1.23

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	309	ASP	N-CA-C	-7.12	104.60	113.28

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2553	2551	2550	15	0
1	B	2553	2550	2550	18	0
2	A	45	55	0	0	0
2	B	45	55	0	1	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	A	48	25	24	1	0
3	B	48	25	24	2	0
4	B	12	20	20	1	0
5	A	83	0	0	0	0
5	B	94	0	0	0	0
All	All	5481	5281	5168	35	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (35) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:A:402:NAP:C1D	3:A:402:NAP:O4D	1.68	1.16
3:B:402:NAP:C1D	3:B:402:NAP:O4D	1.68	1.10
1:B:137:VAL:HG11	1:B:311:PHE:CE2	2.19	0.77
1:B:77:GLU:O	4:B:403:BU3:H3	2.00	0.61
1:B:234:VAL:HG13	1:B:237:GLU:HB2	1.84	0.59
1:B:270:LYS:NZ	3:B:402:NAP:O2X	2.34	0.59
1:B:304:HIS:O	1:B:304:HIS:ND1	2.41	0.54
1:B:137:VAL:HG11	1:B:311:PHE:HE2	1.71	0.54
1:B:74:VAL:HG22	1:B:78:ASP:HB2	1.90	0.53
1:A:37:VAL:HG11	1:A:272:TYR:CD1	2.44	0.53
1:B:244:LEU:HD12	1:B:296:ILE:HD11	1.90	0.53
1:A:199:ARG:HA	1:B:323:TYR:C	2.35	0.52
2:B:401:A1JRA:O42	2:B:401:A1JRA:N29	2.45	0.50
1:B:130:PRO:CB	1:B:138:ILE:HD12	2.43	0.49
1:A:54:LEU:HD13	1:A:86:TRP:CE2	2.48	0.48
1:A:174:GLU:OE2	1:A:174:GLU:HA	2.13	0.47
1:A:240:VAL:HG21	1:A:299:LEU:HD21	1.97	0.47
1:A:191:VAL:HG22	1:A:213:LEU:HD21	1.96	0.46
1:A:71:ASP:C	1:A:71:ASP:OD1	2.58	0.46
1:A:103:LEU:HD21	1:A:111:VAL:HG13	1.99	0.45
1:A:39:LYS:O	1:A:43:GLU:HG3	2.17	0.44
1:B:131:THR:HG22	1:B:137:VAL:HG12	1.98	0.44
1:B:130:PRO:HB2	1:B:138:ILE:HD12	2.00	0.44
1:B:39:LYS:O	1:B:43:GLU:HG3	2.18	0.44
1:B:105:LYS:HE3	1:B:105:LYS:HB3	1.89	0.43
1:A:137:VAL:HG11	1:A:311:PHE:CE1	2.54	0.43
1:B:71:ASP:OD1	1:B:71:ASP:C	2.62	0.43
1:B:114:TYR:CD2	1:B:151:MET:HE1	2.55	0.42

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:140:ASP:C	1:A:141:ILE:HD13	2.45	0.42
1:B:30:PRO:HD2	1:B:33:LYS:HE3	2.01	0.41
1:A:254:LEU:HD21	1:A:279:GLN:O	2.20	0.41
1:A:20:GLY:HA2	1:A:48:HIS:HB3	2.03	0.41
1:A:244:LEU:HD13	1:A:244:LEU:HA	1.96	0.41
1:A:48:HIS:CD2	1:A:48:HIS:C	2.98	0.41
1:B:308:SER:HB3	1:B:311:PHE:CE1	2.56	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	316/344 (92%)	303 (96%)	13 (4%)	0	100	100
1	B	316/344 (92%)	305 (96%)	10 (3%)	1 (0%)	36	35
All	All	632/688 (92%)	608 (96%)	23 (4%)	1 (0%)	43	42

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	134	ASN

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	282/305 (92%)	282 (100%)	0	100	100
1	B	282/305 (92%)	281 (100%)	1 (0%)	84	89
All	All	564/610 (92%)	563 (100%)	1 (0%)	87	92

All (1) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	123	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry ⓘ

6 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	BU3	B	404	-	4,5,5	0.71	0	6,6,6	0.37	0
4	BU3	B	403	-	4,5,5	0.29	0	6,6,6	0.36	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	A1JRA	A	401	-	44,45,45	1.94	7 (15%)	51,58,58	1.67	11 (21%)
3	NAP	B	402	-	49,52,52	4.90	18 (36%)	69,80,80	1.51	11 (15%)
3	NAP	A	402	-	49,52,52	4.90	18 (36%)	69,80,80	1.63	12 (17%)
2	A1JRA	B	401	-	44,45,45	1.94	10 (22%)	51,58,58	1.36	8 (15%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	BU3	B	404	-	-	3/4/4/4	-
4	BU3	B	403	-	-	0/4/4/4	-
2	A1JRA	A	401	-	-	18/48/68/68	0/1/1/1
3	NAP	B	402	-	-	4/35/67/67	0/5/5/5
3	NAP	A	402	-	-	9/35/67/67	0/5/5/5
2	A1JRA	B	401	-	-	25/48/68/68	0/1/1/1

All (53) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	B	402	NAP	O4D-C1D	19.85	1.68	1.41
3	A	402	NAP	O4D-C1D	19.71	1.68	1.41
3	A	402	NAP	C2D-C1D	-15.88	1.29	1.53
3	B	402	NAP	C2D-C1D	-15.69	1.30	1.53
3	A	402	NAP	C3B-C2B	-12.64	1.24	1.52
3	B	402	NAP	C3B-C2B	-12.51	1.25	1.52
3	B	402	NAP	C7N-N7N	8.99	1.50	1.33
3	A	402	NAP	C7N-N7N	8.95	1.50	1.33
2	A	401	A1JRA	C33-N34	6.91	1.48	1.34
2	B	401	A1JRA	C33-N34	6.87	1.47	1.34
2	A	401	A1JRA	C28-N29	6.58	1.48	1.34
3	B	402	NAP	O4D-C4D	-6.58	1.30	1.45
3	A	402	NAP	O4D-C4D	-6.57	1.30	1.45
2	B	401	A1JRA	C28-N29	6.50	1.47	1.34
3	A	402	NAP	O4B-C1B	-6.12	1.27	1.42
3	B	402	NAP	O4B-C1B	-6.09	1.27	1.42
3	B	402	NAP	C6A-N6A	5.60	1.48	1.34
3	A	402	NAP	C6A-N6A	5.58	1.48	1.34
3	B	402	NAP	C2B-C1B	5.19	1.66	1.53

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	402	NAP	C2B-C1B	5.19	1.66	1.53
3	A	402	NAP	P2B-O2B	5.08	1.68	1.59
3	B	402	NAP	P2B-O2B	5.06	1.68	1.59
3	B	402	NAP	O3D-C3D	-4.85	1.31	1.43
3	A	402	NAP	O3D-C3D	-4.71	1.31	1.43
3	B	402	NAP	O3B-C3B	4.33	1.53	1.43
3	A	402	NAP	O3B-C3B	4.28	1.53	1.43
3	B	402	NAP	C3N-C7N	4.13	1.56	1.50
3	A	402	NAP	C3N-C7N	4.11	1.56	1.50
3	A	402	NAP	C5B-C4B	-4.02	1.39	1.51
3	B	402	NAP	C5B-C4B	-3.85	1.39	1.51
3	B	402	NAP	O4B-C4B	3.21	1.52	1.45
3	A	402	NAP	O4B-C4B	3.15	1.52	1.45
2	A	401	A1JRA	C44-C43	-3.01	1.32	1.51
2	B	401	A1JRA	C44-C43	-2.98	1.33	1.51
2	A	401	A1JRA	O09-C06	2.74	1.48	1.41
3	B	402	NAP	PN-O5D	2.51	1.69	1.59
3	B	402	NAP	O2D-C2D	2.47	1.48	1.43
3	A	402	NAP	O2D-C2D	2.39	1.48	1.43
3	A	402	NAP	PN-O5D	2.38	1.68	1.59
2	A	401	A1JRA	C31-C32	2.37	1.55	1.51
2	B	401	A1JRA	O05-C04	-2.34	1.40	1.44
2	B	401	A1JRA	C31-C32	2.32	1.55	1.51
2	B	401	A1JRA	O09-C06	2.28	1.47	1.41
2	A	401	A1JRA	O41-C33	-2.26	1.19	1.23
3	B	402	NAP	PA-O5B	2.22	1.68	1.59
2	B	401	A1JRA	C27-C28	2.22	1.55	1.51
2	A	401	A1JRA	O16-C15	2.15	1.48	1.43
2	B	401	A1JRA	O41-C33	-2.10	1.19	1.23
3	A	402	NAP	PA-O5B	2.05	1.67	1.59
2	B	401	A1JRA	O45-C28	-2.04	1.19	1.23
2	B	401	A1JRA	O16-C15	2.03	1.47	1.43
3	A	402	NAP	C8A-N9A	-2.01	1.33	1.37
3	B	402	NAP	C8A-N9A	-2.01	1.33	1.37

All (42) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	401	A1JRA	C31-C32-C33	5.49	122.81	115.98
3	A	402	NAP	C5A-C4A-N3A	-5.01	120.21	126.75
3	B	402	NAP	C5A-C4A-N3A	-4.98	120.25	126.75
3	A	402	NAP	N3A-C2A-N1A	-4.04	122.28	128.60

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	401	A1JRA	C06-C07-C15	4.00	118.32	110.00
3	B	402	NAP	N3A-C2A-N1A	-3.99	122.37	128.60
2	B	401	A1JRA	C44-C43-C30	3.93	130.47	113.86
2	A	401	A1JRA	C44-C43-C30	3.70	129.49	113.86
3	A	402	NAP	C5A-N7A-C8A	3.60	108.62	103.51
3	A	402	NAP	C3D-C2D-C1D	3.54	106.31	100.98
3	A	402	NAP	C2A-N3A-C4A	3.49	119.99	111.75
3	B	402	NAP	C5A-N7A-C8A	3.45	108.41	103.51
2	B	401	A1JRA	C31-C32-C33	3.44	120.27	115.98
3	B	402	NAP	C2A-N3A-C4A	3.41	119.82	111.75
3	A	402	NAP	C4A-C5A-N7A	-3.36	106.53	110.62
3	B	402	NAP	N3A-C4A-N9A	3.26	132.46	127.08
3	A	402	NAP	N3A-C4A-N9A	3.20	132.36	127.08
3	A	402	NAP	PN-O3-PA	-3.18	121.92	132.83
3	B	402	NAP	C4A-C5A-N7A	-3.16	106.77	110.62
3	A	402	NAP	N9A-C8A-N7A	-3.08	109.71	113.91
3	B	402	NAP	C3D-C2D-C1D	3.00	105.50	100.98
2	A	401	A1JRA	C31-C30-N29	2.98	113.56	110.11
3	A	402	NAP	C4B-O4B-C1B	-2.95	102.96	109.47
3	B	402	NAP	N9A-C8A-N7A	-2.90	109.95	113.91
2	B	401	A1JRA	O09-C10-C13	2.66	114.53	109.69
2	B	401	A1JRA	C43-C30-N29	2.64	113.53	110.32
3	B	402	NAP	C4B-O4B-C1B	-2.64	103.64	109.47
3	B	402	NAP	C3B-C2B-C1B	2.54	107.66	102.89
2	A	401	A1JRA	O09-C06-C07	2.53	115.70	110.35
3	A	402	NAP	C3B-C2B-C1B	2.49	107.56	102.89
2	A	401	A1JRA	C30-N29-C28	-2.41	119.36	122.92
2	B	401	A1JRA	C31-C30-N29	-2.39	107.33	110.11
2	A	401	A1JRA	O09-C10-C13	2.25	113.77	109.69
2	B	401	A1JRA	C15-C13-C10	2.18	114.13	110.24
3	A	402	NAP	C6A-C5A-N7A	2.18	136.08	132.02
2	A	401	A1JRA	C18-C17-C04	-2.17	108.16	113.65
2	B	401	A1JRA	C27-C28-N29	2.11	119.49	115.83
2	A	401	A1JRA	C06-O09-C10	2.08	117.77	113.69
2	A	401	A1JRA	O38-C36-C35	2.07	120.29	113.40
2	B	401	A1JRA	C26-C27-C28	-2.04	107.55	113.26
2	A	401	A1JRA	C03-C02-C04	2.03	116.17	112.03
3	B	402	NAP	C6A-C5A-N7A	2.01	135.76	132.02

There are no chirality outliers.

All (59) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	401	A1JRA	C01-C02-C04-C17
2	A	401	A1JRA	C01-C02-C04-O05
2	A	401	A1JRA	C03-C02-C04-C17
2	A	401	A1JRA	C03-C02-C04-O05
2	A	401	A1JRA	C31-C30-C43-C44
2	A	401	A1JRA	N29-C30-C43-C44
2	B	401	A1JRA	C02-C04-C17-C18
2	B	401	A1JRA	C03-C02-C04-C17
2	B	401	A1JRA	C03-C02-C04-O05
2	B	401	A1JRA	C31-C30-C43-C44
2	B	401	A1JRA	N29-C30-C43-C44
3	A	402	NAP	C5B-O5B-PA-O1A
3	A	402	NAP	C5D-O5D-PN-O3
3	A	402	NAP	C5D-O5D-PN-O1N
3	A	402	NAP	O4D-C1D-N1N-C2N
3	A	402	NAP	O4D-C1D-N1N-C6N
3	B	402	NAP	PA-O3-PN-O5D
2	B	401	A1JRA	C13-C10-C11-O12
2	A	401	A1JRA	C18-C19-C20-C21
2	B	401	A1JRA	O09-C10-C11-O12
2	B	401	A1JRA	C25-C26-C27-C28
2	B	401	A1JRA	C39-C35-N34-C33
2	B	401	A1JRA	C20-C21-C22-C23
2	A	401	A1JRA	O09-C06-O05-C04
2	A	401	A1JRA	C39-C35-N34-C33
2	A	401	A1JRA	C24-C25-C26-C27
2	A	401	A1JRA	C19-C20-C21-C22
2	A	401	A1JRA	C17-C18-C19-C20
2	B	401	A1JRA	C24-C25-C26-C27
2	B	401	A1JRA	C01-C02-C04-O05
2	B	401	A1JRA	C23-C24-C25-C26
2	B	401	A1JRA	C01-C02-C04-C17
2	B	401	A1JRA	C31-C30-N29-C28
2	A	401	A1JRA	O09-C10-C11-O12
2	A	401	A1JRA	C43-C30-N29-C28
2	B	401	A1JRA	C43-C30-N29-C28
2	B	401	A1JRA	C18-C19-C20-C21
3	A	402	NAP	PA-O3-PN-O5D
3	B	402	NAP	C2B-O2B-P2B-O2X
4	B	404	BU3	C1-C2-C3-O6
4	B	404	BU3	O5-C2-C3-C4
2	A	401	A1JRA	C31-C30-N29-C28
2	A	401	A1JRA	C39-C35-C36-O38

Continued on next page...

Continued from previous page...



Mol	Chain	Res	Type	Atoms
2	B	401	A1JRA	C19-C20-C21-C22
2	A	401	A1JRA	C07-C06-O05-C04
2	A	401	A1JRA	C39-C35-C36-O37
2	B	401	A1JRA	O09-C06-O05-C04
2	B	401	A1JRA	C07-C06-O05-C04
2	B	401	A1JRA	C22-C23-C24-C25
2	B	401	A1JRA	C39-C35-C36-O38
2	B	401	A1JRA	C39-C35-C36-O37
2	B	401	A1JRA	C17-C18-C19-C20
4	B	404	BU3	O5-C2-C3-O6
3	A	402	NAP	C5B-O5B-PA-O3
3	B	402	NAP	C2B-O2B-P2B-O3X
3	B	402	NAP	C4D-C5D-O5D-PN
3	A	402	NAP	O4D-C4D-C5D-O5D
2	B	401	A1JRA	O05-C04-C17-C18
3	A	402	NAP	C4D-C5D-O5D-PN


There are no ring outliers.

4 monomers are involved in 5 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	B	403	BU3	1	0
3	B	402	NAP	2	0
3	A	402	NAP	1	0
2	B	401	A1JRA	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	318/344 (92%)	0.88	24 (7%) 20 19	11, 20, 31, 39	0
1	B	318/344 (92%)	0.89	28 (8%) 15 14	11, 20, 32, 38	0
All	All	636/688 (92%)	0.89	52 (8%) 17 16	11, 20, 32, 39	0

All (52) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	309	ASP	7.0
1	B	309	ASP	4.8
1	B	134	ASN	4.2
1	A	227	TRP	4.0
1	B	227	TRP	3.8
1	B	311	PHE	3.7
1	B	283	VAL	3.6
1	A	125	GLY	3.4
1	B	133	GLU	3.3
1	B	321	ASP	3.3
1	B	203	LEU	3.2
1	A	310	SER	3.2
1	A	176	ILE	3.2
1	B	307	ASN	3.2
1	B	306	PHE	3.2
1	B	81	TYR	3.1
1	B	323	TYR	3.1
1	B	219	LEU	3.0
1	B	37	VAL	3.0
1	B	322	GLU	2.9
1	B	135	GLY	2.8
1	A	133	GLU	2.8
1	A	244	LEU	2.7
1	A	308	SER	2.7

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	283	VAL	2.7
1	B	139	PHE	2.7
1	A	210	ASP	2.6
1	B	226	ARG	2.5
1	A	75	LYS	2.5
1	B	126	GLU	2.5
1	B	65	ILE	2.5
1	A	74	VAL	2.5
1	B	138	ILE	2.5
1	A	307	ASN	2.4
1	B	141	ILE	2.4
1	A	131	THR	2.4
1	B	253	ALA	2.4
1	A	320	SER	2.3
1	B	137	VAL	2.3
1	A	137	VAL	2.3
1	B	315	PRO	2.3
1	B	261	LEU	2.2
1	A	205	PHE	2.1
1	A	7	CYS	2.1
1	A	225	LYS	2.1
1	A	294	LYS	2.1
1	A	141	ILE	2.1
1	B	67	SER	2.1
1	A	261	LEU	2.1
1	B	122	LEU	2.1
1	A	37	VAL	2.0
1	A	116	ILE	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum,




median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
4	BU3	B	404	6/6	0.75	0.15	20,20,20,20	0
2	A1JRA	A	401	45/45	0.77	0.14	14,27,40,44	0
2	A1JRA	B	401	45/45	0.80	0.13	13,23,30,30	0
4	BU3	B	403	6/6	0.81	0.11	20,24,29,31	0
3	NAP	A	402	48/48	0.93	0.09	10,15,18,21	0
3	NAP	B	402	48/48	0.93	0.08	10,15,21,23	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



Electron density around A1JRA A 401:

2mF_o-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)







Electron density around A1JRA B 401:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around NAP A 402:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.5 Other polymers [i](#)

There are no such residues in this entry.