



# Full wwPDB X-ray Structure Validation Report ⓘ

Jun 30, 2026 – 05:54 pm BST

PDB ID : 9T07 / pdb\_00009t07  
Title : HUMAN PI3KDELTA IN COMPLEX WITH Roginolisib  
Authors : Graedler, U.; Augustin, M.; Goesser, C.; Kiefersauer, R.  
Deposited on : 2025-10-16  
Resolution : 2.75 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0  
Mogul : 1.8.4, CSD as541be (2020)  
Xtriage (Phenix) : 2.0  
EDS : 3.0  
Buster-report : wwPDB partial adaption of 1.1.7 (2018)  
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)  
CCP4 : 9.0.015 (Gargrove)  
Density-Fitness : 1.0.12  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.50


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.75 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	180053	1009 (2.76-2.76)
Clashscore	190562	1044 (2.76-2.76)
Ramachandran outliers	187476	1024 (2.76-2.76)
Sidechain outliers	187428	1024 (2.76-2.76)
RSRZ outliers	180081	1009 (2.76-2.76)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	1044	 6% 75% 12% • 12%
2	B	169	 2% 79% 14% • 7%

## 2 Entry composition i

There are 4 unique types of molecules in this entry. The entry contains 8814 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Phosphatidylinositol 4,5-bisphosphate 3-kinase catalytic subunit delta isoform.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	914	7379	4724	1254	1348	53	22	0	0

- Molecule 2 is a protein called Phosphatidylinositol 3-kinase regulatory subunit alpha.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	B	158	1370	857	247	261	5	0	0	0

- Molecule 3 is [6-fluoranyl-1-[4-(morpholin-4-ylmethyl)phenyl]-5,5-bis(oxidanylidene)-4H-thiochromeno[4,3-c]pyrazol-3-yl]-morpholin-4-yl-methanone (CCD ID: QGU) (formula: C<sub>26</sub>H<sub>27</sub>FN<sub>4</sub>O<sub>5</sub>S) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms						ZeroOcc	AltConf
			Total	C	F	N	O	S		
3	A	1	37	26	1	4	5	1	0	0


- Molecule 4 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	21	Total O 21 21	0	1
4	B	7	Total O 7 7	0	0


### 3 Residue-property plots


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Phosphatidylinositol 4,5-bisphosphate 3-kinase catalytic subunit delta isoform



- Molecule 2: Phosphatidylinositol 3-kinase regulatory subunit alpha





## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	93.03Å 113.39Å 144.51Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	46.52 – 2.75 46.52 – 2.75	Depositor EDS
% Data completeness (in resolution range)	96.8 (46.52-2.75) 96.7 (46.52-2.75)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	0.04	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.28 (at 2.77Å)	Xtrriage
Refinement program	BUSTER 2.11.8 (17-JUL-2025)	Depositor
R, $R_{free}$	0.231 , 0.273 0.233 , 0.266	Depositor DCC
$R_{free}$ test set	746 reflections (1.91%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	85.9	Xtrriage
Anisotropy	0.247	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.31 , 57.8	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.94	EDS
Total number of atoms	8814	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	101.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.67% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: QGU

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.66	0/7542	1.03	0/10196
2	B	0.65	0/1388	1.16	0/1849
All	All	0.66	0/8930	1.05	0/12045

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	7379	0	7346	64	0
2	B	1370	0	1377	19	0
3	A	37	0	0	0	0
4	A	21	0	0	0	0
4	B	7	0	0	0	0
All	All	8814	0	8723	82	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (82) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:439:GLU:HB2	2:B:445:VAL:HG22	1.41	0.99
2:B:442:ILE:HD13	2:B:589:VAL:HG22	1.62	0.81
1:A:427:ASP:HB3	1:A:433:LYS:HD2	1.67	0.77
1:A:386:ARG:HG3	1:A:387:MET:HE2	1.70	0.74
1:A:424:MET:O	1:A:433:LYS:NZ	2.24	0.71
1:A:755:LYS:HZ2	1:A:756:MET:HG2	1.55	0.69
2:B:439:GLU:HB2	2:B:445:VAL:CG2	2.19	0.69
2:B:442:ILE:HD13	2:B:589:VAL:CG2	2.23	0.67
1:A:641:ARG:HH22	1:A:732:GLN:HE22	1.44	0.65
1:A:336:ASP:O	1:A:365:VAL:HG11	1.97	0.64
2:B:439:GLU:CB	2:B:445:VAL:HG22	2.22	0.63
1:A:589:ASP:OD2	1:A:591:HIS:ND1	2.29	0.62
1:A:491:ILE:HG21	1:A:565:LEU:CD1	2.30	0.62
1:A:396:ALA:HB2	1:A:418:ILE:HD11	1.83	0.60
1:A:549:LEU:HD23	1:A:564:MET:HE1	1.84	0.60
2:B:439:GLU:CB	2:B:445:VAL:CG2	2.79	0.60
1:A:512:ARG:HG2	1:A:542:PHE:HZ	1.66	0.59
1:A:641:ARG:NH2	1:A:732:GLN:HE22	2.00	0.59
1:A:73:ILE:HD12	2:B:489:GLU:HB3	1.83	0.59
1:A:321:GLN:HG3	1:A:381:ILE:HD12	1.84	0.58
1:A:159:TRP:HB3	1:A:283:LEU:HG	1.85	0.58
1:A:902:ARG:HG3	1:A:906:GLN:HB2	1.86	0.56
1:A:387:MET:HE3	1:A:590:CYS:HB2	1.88	0.56
1:A:789:LEU:HD13	1:A:981:MET:HE2	1.88	0.56
1:A:322:PRO:HA	1:A:380:ASN:HA	1.88	0.55
1:A:387:MET:HG3	1:A:589:ASP:HA	1.88	0.55
1:A:755:LYS:NZ	1:A:756:MET:HG2	2.20	0.55
1:A:895:HIS:ND1	1:A:897:ASP:OD1	2.40	0.54
1:A:491:ILE:HG21	1:A:565:LEU:HD12	1.91	0.52
1:A:113:ILE:HG23	1:A:679:MET:HE1	1.91	0.52
2:B:526:HIS:CE1	2:B:530:LYS:NZ	2.78	0.52
1:A:76:THR:HG21	1:A:118:SER:HB3	1.92	0.51
1:A:241:LEU:HB2	1:A:250:LEU:HB2	1.92	0.51
1:A:394:LEU:HB2	1:A:441:MET:HE1	1.93	0.51
1:A:656:HIS:CE1	1:A:820:ASP:HA	2.45	0.51
1:A:716:HIS:O	1:A:720:ARG:HG2	2.11	0.51
1:A:608:LEU:HD23	1:A:643:ILE:HD13	1.93	0.50
1:A:213:PRO:HD3	1:A:254:TYR:O	2.11	0.50
1:A:246:ARG:NH2	1:A:739:THR:OG1	2.44	0.50
1:A:172:GLU:HG3	1:A:263:CYS:SG	2.52	0.50
1:A:517:ARG:O	1:A:518:ARG:O	2.29	0.50
1:A:169:LEU:HD22	1:A:259:PHE:HE1	1.77	0.49

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:610:GLN:HE22	1:A:795:GLN:NE2	2.10	0.49
2:B:439:GLU:HB3	2:B:445:VAL:HG23	1.94	0.49
1:A:324:ARG:HD2	1:A:374:ARG:HD2	1.94	0.49
2:B:526:HIS:CE1	2:B:530:LYS:HZ2	2.31	0.49
1:A:18:GLN:NE2	1:A:37:SER:OG	2.46	0.48
1:A:207:VAL:HG11	1:A:216:LEU:HD12	1.95	0.48
1:A:436:GLU:HA	1:A:474:CYS:HA	1.96	0.48
1:A:383:ASP:HA	1:A:558:HIS:HB3	1.95	0.48
1:A:69:VAL:HG23	1:A:104:ARG:HB2	1.95	0.47
1:A:549:LEU:CD2	1:A:564:MET:HE1	2.44	0.47
1:A:587:PHE:HB3	1:A:592:VAL:HG11	1.97	0.47
1:A:763:TYR:HB2	1:A:774:VAL:HG23	1.96	0.47
1:A:491:ILE:HG21	1:A:565:LEU:HD13	1.94	0.47
1:A:142:LYS:HG2	1:A:143:MET:HE2	1.97	0.47
1:A:350:GLY:HA3	1:A:588:PRO:HG3	1.95	0.46
1:A:74:ASN:OD1	1:A:76:THR:HG22	2.15	0.46
2:B:526:HIS:HD2	2:B:529:GLU:OE1	2.00	0.45
2:B:463:TYR:HB2	2:B:566:ILE:HG21	1.98	0.45
1:A:860:LYS:HA	1:A:868:LEU:HD13	1.98	0.45
1:A:337:GLU:H	1:A:337:GLU:HG3	1.47	0.45
1:A:31:TYR:CD1	1:A:31:TYR:O	2.71	0.44
1:A:715:MET:HE2	1:A:759:LEU:HD22	2.00	0.44
1:A:742:ALA:HB3	1:A:765:ASN:HA	1.99	0.43
2:B:593:LYS:HA	2:B:596:GLU:HG2	2.01	0.43
1:A:558:HIS:CE1	1:A:559:GLU:HG3	2.53	0.43
2:B:526:HIS:HE1	2:B:530:LYS:NZ	2.17	0.43
1:A:491:ILE:CG2	1:A:565:LEU:HD12	2.48	0.43
2:B:542:ARG:O	2:B:546:GLU:HG3	2.19	0.43
1:A:535:ARG:NH2	1:A:563:GLN:NE2	2.67	0.43
1:A:419:ALA:HB3	1:A:441:MET:HE3	1.99	0.42
2:B:463:TYR:CD1	2:B:463:TYR:C	2.97	0.42
2:B:484:ILE:HG13	2:B:545:LEU:HD23	2.00	0.42
1:A:34:PHE:HB2	1:A:35:PRO:HD2	2.00	0.42
1:A:394:LEU:CB	1:A:441:MET:HE1	2.50	0.41
1:A:433:LYS:HB3	1:A:475:LEU:HD13	2.03	0.41
1:A:917:GLY:HA2	1:A:930:VAL:HG23	2.03	0.40
2:B:480:LYS:HG2	2:B:545:LEU:HD11	2.03	0.40
1:A:336:ASP:O	1:A:365:VAL:HG21	2.21	0.40
1:A:329:GLN:HG2	1:A:370:VAL:HG22	2.03	0.40
2:B:583:TRP:CZ2	2:B:587:LYS:HE3	2.57	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	898/1044 (86%)	863 (96%)	32 (4%)	3 (0%)	36	56
2	B	154/169 (91%)	154 (100%)	0	0	100	100
All	All	1052/1213 (87%)	1017 (97%)	32 (3%)	3 (0%)	36	56

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	518	ARG
1	A	495	GLY
1	A	766	GLU

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	815/927 (88%)	771 (95%)	44 (5%)	20	38
2	B	150/160 (94%)	143 (95%)	7 (5%)	23	45
All	All	965/1087 (89%)	914 (95%)	51 (5%)	20	39

All (51) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	18	GLN
1	A	32	LEU
1	A	33	ASN

*Continued on next page...*

*Continued from previous page...*

<b>Mol</b>	<b>Chain</b>	<b>Res</b>	<b>Type</b>
1	A	56	GLU
1	A	66	GLU
1	A	71	THR
1	A	93	GLN
1	A	100	ARG
1	A	104	ARG
1	A	105	GLU
1	A	109	VAL
1	A	110	LYS
1	A	112	LEU
1	A	207	VAL
1	A	240	THR
1	A	263	CYS
1	A	337	GLU
1	A	344	GLN
1	A	368	GLU
1	A	431	GLN
1	A	433	LYS
1	A	441	MET
1	A	445	VAL
1	A	459	VAL
1	A	473	ILE
1	A	475	LEU
1	A	507	GLU
1	A	518	ARG
1	A	602	LYS
1	A	731	LEU
1	A	740	LEU
1	A	755	LYS
1	A	757	LYS
1	A	767	GLU
1	A	782	ASP
1	A	795	GLN
1	A	804	GLU
1	A	902	ARG
1	A	915	PHE
1	A	933	ILE
1	A	937	ASP
1	A	940	HIS
1	A	948	ASN
1	A	968	ARG
2	B	458	GLU

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
2	B	460	SER
2	B	482	THR
2	B	520	GLU
2	B	530	LYS
2	B	543	ARG
2	B	558	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (21) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	18	GLN
1	A	51	HIS
1	A	80	GLN
1	A	86	GLN
1	A	258	GLN
1	A	321	GLN
1	A	558	HIS
1	A	563	GLN
1	A	610	GLN
1	A	656	HIS
1	A	696	ASN
1	A	730	HIS
1	A	732	GLN
1	A	748	GLN
1	A	765	ASN
1	A	795	GLN
1	A	918	ASN
1	A	943	GLN
1	A	948	ASN
2	B	441	ASN
2	B	526	HIS

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
3	QGU	A	1101	-	38,42,42	1.16	2 (5%)	52,62,62	2.08	12 (23%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	QGU	A	1101	-	-	1/16/47/47	0/5/6/6

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	1101	QGU	N14-N13	-4.99	1.28	1.39
3	A	1101	QGU	C2-C10	2.18	1.49	1.45

All (12) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	1101	QGU	C12-N13-N14	7.62	112.52	104.75
3	A	1101	QGU	C11-C12-N13	-7.16	105.40	111.61
3	A	1101	QGU	O31-S3-C1	-3.45	105.22	108.60
3	A	1101	QGU	C10-N14-N13	-3.40	109.19	112.29
3	A	1101	QGU	C26-N25-C30	3.09	115.79	108.83
3	A	1101	QGU	C6-C11-C12	-2.84	129.90	132.81
3	A	1101	QGU	O32-S3-O31	2.82	120.14	117.75

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	1101	QGU	C10-C11-C12	2.55	106.94	104.93
3	A	1101	QGU	C37-N17-C33	2.46	117.35	112.62
3	A	1101	QGU	C29-O28-C27	2.45	118.07	109.89
3	A	1101	QGU	C36-O35-C34	2.20	117.23	109.89
3	A	1101	QGU	O16-C15-N17	-2.16	119.22	122.67

There are no chirality outliers.


All (1) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	1101	QGU	C21-C24-N25-C30

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	914/1044 (87%)	0.52	58 (6%) 26 26	27, 99, 152, 182	7 (0%)
2	B	158/169 (93%)	0.11	3 (1%) 66 64	68, 87, 105, 120	0
All	All	1072/1213 (88%)	0.46	61 (5%) 29 29	27, 96, 149, 182	7 (0%)

All (61) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	194	VAL	4.1
1	A	549	LEU	3.8
1	A	32	LEU	3.8
1	A	471	LEU	3.8
1	A	216	LEU	3.7
1	A	856	LEU	3.7
1	A	171	LEU	3.6
1	A	442	TRP	3.4
1	A	236	PRO	3.3
1	A	458	THR	3.3
1	A	659	SER	3.2
1	A	683	MET	3.1
1	A	877	LEU	3.0
1	A	317	TRP	3.0
1	A	534	LEU	3.0
1	A	266	LEU	2.9
1	A	97	PRO	2.9
1	A	865	GLY	2.9
1	A	209	THR	2.9
1	A	257	CYS	2.9
1	A	192	VAL	2.9
1	A	272	PRO	2.8
1	A	395	TYR	2.8
1	A	459	VAL	2.8

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	A	947	THR	2.8
1	A	439	LEU	2.7
1	A	441	MET	2.7
1	A	892	GLY	2.7
2	B	524	ILE	2.7
1	A	288	GLU	2.7
1	A	101	LEU	2.6
1	A	523	LEU	2.5
1	A	590	CYS	2.4
1	A	497	HIS	2.4
1	A	942	ILE	2.4
2	B	520	GLU	2.4
1	A	222	ARG	2.4
2	B	508	TYR	2.4
1	A	274	LEU	2.3
1	A	864	PRO	2.3
1	A	258	GLN	2.3
1	A	207	VAL	2.3
1	A	470	ALA	2.3
1	A	1027	TRP	2.3
1	A	425	LEU	2.3
1	A	714	LEU	2.3
1	A	419	ALA	2.2
1	A	259	PHE	2.2
1	A	191	LEU	2.2
1	A	468	ALA	2.2
1	A	316	LEU	2.1
1	A	561	VAL	2.1
1	A	893	ASP	2.1
1	A	465	THR	2.1
1	A	936	TYR	2.1
1	A	950	SER	2.0
1	A	162	TRP	2.0
1	A	546	LEU	2.0
1	A	160	GLU	2.0
1	A	262	ILE	2.0
1	A	450	GLY	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates


There are no oligosaccharides in this entry.

### 6.4 Ligands

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
3	QGU	A	1101	37/37	0.86	0.12	124,126,129,129	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



## 6.5 Other polymers [i](#)

There are no such residues in this entry.