



Full wwPDB X-ray Structure Validation Report ⓘ

Aug 18, 2026 – 10:20 am BST

PDB ID : 9T0W / pdb_00009t0w
Title : Crystal structure of the Mycobacterium tuberculosis histidinol-phosphate aminotransferase HisC in complex with the adduct PLP-BVL3572S
Authors : Wintjens, R.; Grosse, C.
Deposited on : 2025-10-20
Resolution : 1.74 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	1.8.4, CSD as541be (2020)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.015 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.50


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.74 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1187 (1.74-1.74)
Clashscore	190562	1207 (1.74-1.74)
Ramachandran outliers	187476	1200 (1.74-1.74)
Sidechain outliers	187428	1200 (1.74-1.74)
RSRZ outliers	180081	1188 (1.74-1.74)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	393	7% 83% 9% 7%
1	B	393	10% 83% 7% 7%

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 5831 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Histidinol-phosphate aminotransferase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	364	Total	C	N	O	S	0	0	0
			2748	1737	483	523	5			
1	B	364	Total	C	N	O	S	0	1	0
			2754	1740	484	525	5			

There are 26 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	381	LYS	-	expression tag	UNP P9WML7
A	382	LEU	-	expression tag	UNP P9WML7
A	383	ALA	-	expression tag	UNP P9WML7
A	384	ALA	-	expression tag	UNP P9WML7
A	385	ALA	-	expression tag	UNP P9WML7
A	386	LEU	-	expression tag	UNP P9WML7
A	387	GLU	-	expression tag	UNP P9WML7
A	388	HIS	-	expression tag	UNP P9WML7
A	389	HIS	-	expression tag	UNP P9WML7
A	390	HIS	-	expression tag	UNP P9WML7
A	391	HIS	-	expression tag	UNP P9WML7
A	392	HIS	-	expression tag	UNP P9WML7
A	393	HIS	-	expression tag	UNP P9WML7
B	381	LYS	-	expression tag	UNP P9WML7
B	382	LEU	-	expression tag	UNP P9WML7
B	383	ALA	-	expression tag	UNP P9WML7
B	384	ALA	-	expression tag	UNP P9WML7
B	385	ALA	-	expression tag	UNP P9WML7
B	386	LEU	-	expression tag	UNP P9WML7
B	387	GLU	-	expression tag	UNP P9WML7
B	388	HIS	-	expression tag	UNP P9WML7
B	389	HIS	-	expression tag	UNP P9WML7
B	390	HIS	-	expression tag	UNP P9WML7
B	391	HIS	-	expression tag	UNP P9WML7
B	392	HIS	-	expression tag	UNP P9WML7

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	393	HIS	-	expression tag	UNP P9WML7

- Molecule 2 is [4-[({E})-[(3 {S})-7-bromanyl-1-oxidanyl-2-oxidanylidene-3,4-dihydroquinolin-3-yl]iminomethyl]-6-methyl-5-oxidanyl-pyridin-3-yl]methyl dihydrogen phosphate (CCD ID: A1JSV) (formula: C₁₇H₁₇BrN₃O₇P) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms						ZeroOcc	AltConf
2	A	1	Total	Br	C	N	O	P	0	0
			29	1	17	3	7	1		
2	B	1	Total	Br	C	N	O	P	0	0
			29	1	17	3	7	1		

- Molecule 3 is DI(HYDROXYETHYL)ETHER (CCD ID: PEG) (formula: C₄H₁₀O₃).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			7	4	3		

- Molecule 4 is GLYCEROL (CCD ID: GOL) (formula: $C_3H_8O_3$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			6	3	3		
4	B	1	Total	C	O	0	0
			6	3	3		
4	B	1	Total	C	O	0	0
			6	3	3		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	B	1	Total	C	O	0	0
			6	3	3		
4	B	1	Total	C	O	0	0
			6	3	3		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	119	Total	O	0	0
			119	119		
5	B	115	Total	O	0	0
			115	115		


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Histidinol-phosphate aminotransferase



- Molecule 1: Histidinol-phosphate aminotransferase



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	67.60Å 101.59Å 115.53Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	58.42 – 1.74 58.42 – 1.74	Depositor EDS
% Data completeness (in resolution range)	99.6 (58.42-1.74) 99.7 (58.42-1.74)	Depositor EDS
R_{merge}	0.17	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.52 (at 1.74Å)	Xtriage
Refinement program	REFMAC 5.8.0425	Depositor
R, R_{free}	0.194 , 0.231 0.203 , 0.237	Depositor DCC
R_{free} test set	4098 reflections (4.95%)	wwPDB-VP
Wilson B-factor (Å ²)	21.8	Xtriage
Anisotropy	0.152	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 26.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	5831	wwPDB-VP
Average B, all atoms (Å ²)	29.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.57% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: PEG, GOL, A1JSV

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.95	2/2805 (0.1%)	1.34	17/3833 (0.4%)
1	B	0.94	1/2811 (0.0%)	1.34	21/3841 (0.5%)
All	All	0.94	3/5616 (0.1%)	1.34	38/7674 (0.5%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	3
1	B	0	6
All	All	0	9

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	208	SER	CA-CB	-6.88	1.43	1.53
1	B	305	ARG	NE-CZ	5.80	1.39	1.33
1	A	305	ARG	NE-CZ	5.04	1.38	1.33

All (38) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	280	ASP	CB-CA-C	-9.76	95.93	110.96
1	A	297	THR	CA-CB-OG1	-8.72	96.52	109.60
1	B	297	THR	CA-CB-OG1	-8.26	97.21	109.60
1	A	305	ARG	CD-NE-CZ	8.16	135.82	124.40
1	B	305	ARG	CD-NE-CZ	7.99	135.59	124.40
1	B	281	THR	CA-CB-OG1	-7.57	98.24	109.60

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	234	PHE	CA-CB-CG	6.96	120.75	113.80
1	B	20	ARG	NE-CZ-NH2	6.80	125.32	119.20
1	B	305	ARG	CG-CD-NE	6.68	126.69	112.00
1	B	59	GLU	CB-CG-CD	6.47	123.61	112.60
1	B	280	ASP	CB-CA-C	-6.43	100.74	110.90
1	A	25	TYR	N-CA-CB	6.36	120.97	110.42
1	B	202	GLU	CG-CD-OE2	-6.13	104.31	118.40
1	B	258	ARG	NE-CZ-NH2	-6.12	113.69	119.20
1	B	121	ILE	CA-C-N	6.11	126.31	121.61
1	B	121	ILE	C-N-CA	6.11	126.31	121.61
1	B	165	LYS	CG-CD-CE	6.04	125.20	111.30
1	A	305	ARG	CG-CD-NE	5.97	125.14	112.00
1	B	20	ARG	NE-CZ-NH1	-5.83	115.67	121.50
1	B	59	GLU	CB-CA-C	5.64	120.15	110.79
1	A	280	ASP	CA-CB-CG	5.62	118.22	112.60
1	B	368	ASP	CA-CB-CG	5.60	118.20	112.60
1	B	94	GLU	CB-CG-CD	5.57	122.07	112.60
1	A	277	HIS	CA-CB-CG	-5.56	108.24	113.80
1	B	202	GLU	CG-CD-OE1	5.49	131.03	118.40
1	A	367	THR	CA-CB-OG1	-5.39	101.51	109.60
1	A	104	GLU	CB-CG-CD	5.36	121.70	112.60
1	A	356	ASP	CA-CB-CG	5.27	117.87	112.60
1	B	127	TYR	CA-C-N	5.27	128.07	120.38
1	B	127	TYR	C-N-CA	5.27	128.07	120.38
1	A	284	SER	CA-CB-OG	5.26	121.62	111.10
1	B	338	ASP	CA-CB-CG	5.24	117.84	112.60
1	A	8	VAL	N-CA-CB	-5.23	103.88	110.31
1	A	86	GLN	CB-CA-C	-5.23	101.97	110.85
1	A	55	ARG	CB-CA-C	-5.21	102.71	110.88
1	A	331	GLU	CB-CA-C	-5.12	99.60	110.31
1	A	94	GLU	CB-CG-CD	5.06	121.21	112.60
1	B	55	ARG	NE-CZ-NH1	-5.01	116.49	121.50

There are no chirality outliers.

All (9) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	305	ARG	Sidechain
1	A	337	ARG	Sidechain
1	A	58	ARG	Sidechain
1	B	16	ARG	Sidechain
1	B	258	ARG	Sidechain

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Group
1	B	305	ARG	Sidechain
1	B	337	ARG	Sidechain
1	B	35	ARG	Sidechain
1	B	55	ARG	Sidechain

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2748	0	2741	20	0
1	B	2754	0	2745	13	0
2	A	29	0	0	0	0
2	B	29	0	0	0	0
3	A	7	0	10	2	0
4	A	6	0	8	1	0
4	B	24	0	32	1	0
5	A	119	0	0	0	0
5	B	115	0	0	0	0
All	All	5831	0	5536	27	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (27) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:54:VAL:HG13	1:B:54:VAL:HG13	1.84	0.58
1:A:233:ALA:HB1	1:A:285:VAL:HG21	1.86	0.57
1:A:125:PRO:HD3	3:A:402:PEG:H22	1.87	0.57
1:B:47:ARG:HA	4:B:404:GOL:H32	1.91	0.53
1:A:322:ALA:HB3	1:A:323:PRO:HD3	1.91	0.52
1:A:266:VAL:HG11	1:B:266:VAL:HG12	1.93	0.50
1:A:266:VAL:CG1	1:B:266:VAL:HG12	2.41	0.50
1:B:127:TYR:CE2	1:B:129:MET:HB2	2.47	0.50
1:B:322:ALA:HB3	1:B:323:PRO:HD3	1.94	0.50
1:B:278:SER:HA	1:B:281:THR:HG22	1.94	0.49

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:127:TYR:CE2	1:A:129:MET:HB2	2.51	0.46
1:A:233:ALA:HB1	1:A:285:VAL:CG2	2.47	0.45
1:A:148:ASN:HA	3:A:402:PEG:H41	1.99	0.44
1:A:321:ASP:OD1	1:A:323:PRO:HD2	2.18	0.44
1:B:83:LEU:HD11	1:B:243:TYR:HE2	1.82	0.44
1:A:32:VAL:HG21	1:A:335:LEU:HB2	1.99	0.44
1:A:266:VAL:HG12	1:B:266:VAL:CG1	2.48	0.42
1:A:110:LEU:HD12	1:A:133:ILE:HG22	2.01	0.42
1:B:61:ALA:HA	1:B:64:LEU:HG	2.01	0.42
1:A:176:ASN:HA	1:A:177:PRO:HA	1.86	0.41
1:A:25:TYR:CE2	4:A:403:GOL:O1	2.69	0.41
1:A:266:VAL:CG1	1:B:266:VAL:CG1	2.98	0.41
1:B:43:HIS:CE1	1:B:351:LEU:HD23	2.56	0.41
1:A:41:ASN:HB2	1:A:232:LYS:O	2.21	0.41
1:A:232:LYS:O	1:A:233:ALA:C	2.62	0.40
1:A:266:VAL:HG12	1:B:266:VAL:HG11	2.03	0.40
1:A:322:ALA:N	1:A:323:PRO:CD	2.85	0.40

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	362/393 (92%)	350 (97%)	11 (3%)	1 (0%)	36	23
1	B	363/393 (92%)	348 (96%)	14 (4%)	1 (0%)	36	23
All	All	725/786 (92%)	698 (96%)	25 (3%)	2 (0%)	36	23

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	28	PRO

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	27	ALA

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	288/311 (93%)	286 (99%)	2 (1%)	76	67
1	B	289/311 (93%)	282 (98%)	7 (2%)	43	19
All	All	577/622 (93%)	568 (98%)	9 (2%)	55	35

All (9) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	305	ARG
1	A	370	VAL
1	B	32	VAL
1	B	35	ARG
1	B	177	PRO
1	B	305	ARG
1	B	310	ASP
1	B	368	ASP
1	B	370	VAL

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

8 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	GOL	A	403	-	5,5,5	0.20	0	5,5,5	0.57	0
4	GOL	B	405	-	5,5,5	0.20	0	5,5,5	0.49	0
2	A1JSV	A	401	-	28,31,31	1.25	2 (7%)	37,46,46	2.33	8 (21%)
4	GOL	B	404	-	5,5,5	0.14	0	5,5,5	0.49	0
3	PEG	A	402	-	6,6,6	0.31	0	5,5,5	0.61	0
4	GOL	B	402	-	5,5,5	0.34	0	5,5,5	0.90	0
4	GOL	B	403	-	5,5,5	0.35	0	5,5,5	0.95	0
2	A1JSV	B	401	-	28,31,31	1.21	4 (14%)	37,46,46	2.37	9 (24%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	GOL	A	403	-	-	4/4/4/4	-
4	GOL	B	405	-	-	2/4/4/4	-
2	A1JSV	A	401	-	-	4/11/27/27	0/3/3/3
4	GOL	B	404	-	-	2/4/4/4	-
3	PEG	A	402	-	-	1/4/4/4	-
4	GOL	B	402	-	-	3/4/4/4	-
4	GOL	B	403	-	-	4/4/4/4	-
2	A1JSV	B	401	-	-	4/11/27/27	0/3/3/3

All (6) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	401	A1JSV	P1-O5	-3.09	1.42	1.54
2	B	401	A1JSV	C10-N1	2.88	1.32	1.27
2	B	401	A1JSV	C4-N2	2.67	1.43	1.39
2	B	401	A1JSV	P1-O5	-2.55	1.45	1.54
2	A	401	A1JSV	C10-N1	2.46	1.31	1.27
2	B	401	A1JSV	C11-C10	2.32	1.51	1.46

All (17) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	401	A1JSV	C1-N1-C10	11.30	133.62	117.31
2	B	401	A1JSV	C1-N1-C10	9.82	131.48	117.31
2	B	401	A1JSV	O7-P1-O5	4.59	125.16	107.64
2	B	401	A1JSV	O5-P1-O4	-4.01	96.06	106.73
2	A	401	A1JSV	O5-P1-O4	-3.53	97.35	106.73
2	B	401	A1JSV	C11-C15-C14	3.02	122.06	120.19
2	A	401	A1JSV	O7-P1-O5	2.73	118.08	107.64
2	B	401	A1JSV	C11-C10-N1	-2.72	116.97	123.01
2	A	401	A1JSV	C4-N2-C9	-2.54	122.90	126.20
2	B	401	A1JSV	C15-C11-C12	-2.50	116.34	118.26
2	B	401	A1JSV	C12-C11-C10	2.41	125.52	121.56
2	A	401	A1JSV	O4-P1-O6	2.35	113.08	106.47
2	A	401	A1JSV	C15-C11-C12	-2.27	116.52	118.26
2	B	401	A1JSV	O2-C9-C1	2.26	125.33	120.30
2	A	401	A1JSV	C16-C14-C15	2.16	123.56	120.89
2	B	401	A1JSV	BR1-C6-C5	-2.10	116.35	119.27
2	A	401	A1JSV	C11-C15-C14	2.09	121.48	120.19

There are no chirality outliers.

All (24) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	401	A1JSV	C17-O4-P1-O5
2	A	401	A1JSV	C17-O4-P1-O7
2	B	401	A1JSV	C17-O4-P1-O5
2	B	401	A1JSV	C17-O4-P1-O7
4	A	403	GOL	C1-C2-C3-O3
4	B	402	GOL	C1-C2-C3-O3
4	B	403	GOL	O1-C1-C2-C3
4	B	404	GOL	C1-C2-C3-O3
4	B	404	GOL	O2-C2-C3-O3
2	A	401	A1JSV	C2-C1-N1-C10

Continued on next page...

Continued from previous page...


Mol	Chain	Res	Type	Atoms
2	B	401	A1JSV	C2-C1-N1-C10
4	B	403	GOL	C1-C2-C3-O3
4	B	405	GOL	O1-C1-C2-C3
4	A	403	GOL	O2-C2-C3-O3
4	B	402	GOL	O2-C2-C3-O3
4	B	403	GOL	O1-C1-C2-O2
4	B	405	GOL	O1-C1-C2-O2
3	A	402	PEG	O2-C3-C4-O4
4	B	403	GOL	O2-C2-C3-O3
2	A	401	A1JSV	C17-O4-P1-O6
2	B	401	A1JSV	C17-O4-P1-O6
4	A	403	GOL	O1-C1-C2-O2
4	A	403	GOL	O1-C1-C2-C3
4	B	402	GOL	O1-C1-C2-O2


There are no ring outliers.

3 monomers are involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	403	GOL	1	0
4	B	404	GOL	1	0
3	A	402	PEG	2	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å ²)	Q<0.9
1	A	364/393 (92%)	0.25	28 (7%)	19 24	15, 24, 53, 108	0
1	B	364/393 (92%)	0.41	38 (10%)	11 15	14, 25, 63, 126	1 (0%)
All	All	728/786 (92%)	0.33	66 (9%)	15 19	14, 24, 58, 126	1 (0%)

All (66) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	25	TYR	7.4
1	B	233	ALA	7.2
1	A	25	TYR	6.8
1	B	370	VAL	6.8
1	A	27	ALA	6.1
1	B	28	PRO	6.1
1	B	24	PRO	5.7
1	B	27	ALA	5.5
1	B	31	ALA	5.4
1	B	7	PRO	5.0
1	B	26	GLY	4.8
1	B	30	LEU	4.8
1	A	26	GLY	4.6
1	A	28	PRO	4.4
1	A	285	VAL	4.4
1	B	369	LEU	4.4
1	A	24	PRO	4.2
1	A	370	VAL	4.2
1	A	31	ALA	4.0
1	A	32	VAL	3.7
1	A	23	ALA	3.7
1	A	7	PRO	3.6
1	B	32	VAL	3.6
1	B	363	ALA	3.5

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	29	GLN	3.5
1	A	30	LEU	3.4
1	A	367	THR	3.4
1	B	184	LEU	3.3
1	B	367	THR	3.3
1	A	286	ALA	3.3
1	B	326	TRP	3.2
1	A	184	LEU	3.0
1	A	29	GLN	3.0
1	A	332	ALA	3.0
1	B	334	ILE	2.9
1	B	330	LEU	2.9
1	B	332	ALA	2.9
1	B	34	VAL	2.8
1	B	335	LEU	2.7
1	B	279	ASP	2.7
1	B	23	ALA	2.7
1	B	305	ARG	2.7
1	A	35	ARG	2.6
1	A	364	ARG	2.6
1	A	369	LEU	2.5
1	B	320	ALA	2.5
1	A	319	PHE	2.4
1	B	35	ARG	2.4
1	A	320	ALA	2.4
1	B	301	ASP	2.4
1	B	327	ARG	2.4
1	B	321	ASP	2.4
1	B	365	ILE	2.3
1	A	305	ARG	2.3
1	A	287	ALA	2.3
1	B	333	GLY	2.2
1	A	365	ILE	2.2
1	B	285	VAL	2.2
1	A	148	ASN	2.2
1	B	328	ARG	2.2
1	B	324	ALA	2.2
1	B	329	TYR	2.2
1	B	352	ALA	2.1
1	A	284	SER	2.1
1	B	22	LYS	2.1
1	A	303	GLY	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
4	GOL	B	404	6/6	0.71	0.18	52,60,65,66	0
4	GOL	B	402	6/6	0.85	0.17	44,47,52,52	0
4	GOL	B	405	6/6	0.85	0.20	50,55,59,73	0
4	GOL	B	403	6/6	0.87	0.15	38,47,51,54	0
3	PEG	A	402	7/7	0.88	0.13	40,43,46,49	0
2	A1JSV	B	401	29/29	0.94	0.08	19,26,35,46	0
4	GOL	A	403	6/6	0.94	0.10	31,47,48,58	0
2	A1JSV	A	401	29/29	0.96	0.07	17,25,34,52	0




The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around A1JSV B 401:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around A1JSV A 401:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.