



Full wwPDB X-ray Structure Validation Report ⓘ

Jul 30, 2026 – 01:12 pm BST

PDB ID : 9T7I / pdb_00009t7i
Title : Plasmodium falciparum Myosin A full-length, post-rigor state, complexed to the inhibitor KNX-115 and ATP-gamma-S
Authors : Moussaoui, D.; Spudich, J.A.; Trybus, K.M.; Trivedi, D.; Robert-Paganin, J.; Houdusse, A.
Deposited on : 2025-11-10
Resolution : 2.35 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	1.8.4, CSD as541be (2020)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.015 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.50


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.35 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1596 (2.36-2.36)
Clashscore	190562	1663 (2.36-2.36)
Ramachandran outliers	187476	1646 (2.36-2.36)
Sidechain outliers	187428	1646 (2.36-2.36)
RSRZ outliers	180081	1598 (2.36-2.36)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	818	5% 86% 12% .
2	B	204	3% 53% 10% 36%
3	E	134	14% 73% 17% . 9%

2 Entry composition [i](#)

There are 8 unique types of molecules in this entry. The entry contains 8898 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Myosin-A.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	807	Total	C	N	O	P	S	0	5	0
			6454	4105	1097	1218	1	33			


- Molecule 2 is a protein called Myosin A tail domain interacting protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	130	Total	C	N	O	S	0	0	0
			1042	656	166	215	5			

- Molecule 3 is a protein called Myosin essential light chain ELC.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	E	122	Total	C	N	O	S	0	2	0
			1020	651	164	199	6			

- Molecule 4 is PHOSPHOTHIOPHOSPHORIC ACID-ADENYLATE ESTER (CCD ID: AGS) (formula: $C_{10}H_{16}N_5O_{12}P_3S$) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms						ZeroOcc	AltConf
4	A	1	Total	C	N	O	P	S	0	0
			31	10	5	12	3	1		

- Molecule 5 is GLYCEROL (CCD ID: GOL) (formula: $C_3H_8O_3$).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	C	O	0	0
			6	3	3		

- Molecule 6 is MAGNESIUM ION (CCD ID: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	1	Total	Mg	0	0
			1	1		

- Molecule 7 is 4-methoxy- {N}-[(3-thiophen-2-yl-1 {H}-pyrazol-4-yl)methyl]benzenesul fonamide (CCD ID: A1JT8) (formula: $C_{15}H_{15}N_3O_3S_2$) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
7	A	1	Total	C	N	O	S	0	0
			23	15	3	3	2		


- Molecule 8 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
8	A	282	Total	O	0	0
			282	282		
8	B	11	Total	O	0	0
			11	11		
8	E	28	Total	O	0	0
			28	28		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: Myosin-A



• Molecule 2: Myosin A tail domain interacting protein



• Molecule 3: Myosin essential light chain ELC



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	90.55Å 114.86Å 171.69Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	71.11 – 2.35 71.11 – 2.35	Depositor EDS
% Data completeness (in resolution range)	76.5 (71.11-2.35) 76.9 (71.11-2.35)	Depositor EDS
R_{merge}	0.11	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.82 (at 2.34Å)	Xtriage
Refinement program	PHENIX 2.0_5867	Depositor
R, R_{free}	0.187 , 0.236 0.188 , 0.236	Depositor DCC
R_{free} test set	2811 reflections (3.72%)	wwPDB-VP
Wilson B-factor (Å ²)	37.9	Xtriage
Anisotropy	0.059	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 47.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	8898	wwPDB-VP
Average B, all atoms (Å ²)	49.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.55% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, MG, SEP, A1JT8, AGS

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.40	1/6555 (0.0%)	0.56	2/8835 (0.0%)
2	B	0.29	0/1059	0.48	0/1428
3	E	0.32	0/1037	0.49	0/1394
All	All	0.38	1/8651 (0.0%)	0.54	2/11657 (0.0%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	462	LYS	CD-CE	6.57	1.72	1.52

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	462	LYS	CG-CD-CE	5.80	124.65	111.30
1	A	462	LYS	CB-CG-CD	-5.60	98.42	111.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	6454	0	6548	56	0
2	B	1042	0	999	17	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	E	1020	0	992	17	0
4	A	31	0	12	0	0
5	A	6	0	8	1	0
6	A	1	0	0	0	0
7	A	23	0	0	0	0
8	A	282	0	0	2	0
8	B	11	0	0	0	0
8	E	28	0	0	2	0
All	All	8898	0	8559	85	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (85) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:552:THR:HG23	1:A:555:LYS:H	1.45	0.81
1:A:731:ASN:HA	1:A:734:LYS:HD2	1.65	0.79
2:B:114:THR:HG23	2:B:117:GLN:H	1.48	0.78
2:B:131:ASP:O	2:B:135:ILE:HG13	1.90	0.71
1:A:745:SER:H	1:A:748:MET:HE2	1.59	0.68
1:A:508:LEU:HD11	1:A:755:MET:HE1	1.82	0.62
1:A:554:GLU:HG3	1:A:579:LYS:HE3	1.81	0.61
1:A:618:ASN:HD22	1:A:621:VAL:H	1.48	0.61
2:B:90:HIS:HA	2:B:92:LYS:NZ	2.17	0.59
1:A:584:GLN:NE2	1:A:589:PRO:HB3	2.18	0.58
3:E:14:ILE:HB	3:E:19:CYS:HA	1.85	0.58
3:E:62:LYS:HE2	3:E:63:HIS:CE1	2.38	0.58
1:A:558:SER:O	1:A:562:THR:HG23	2.03	0.58
1:A:226:ALA:O	1:A:229:PRO:HD2	2.04	0.57
1:A:232:GLU:HG2	1:A:236:ASN:ND2	2.19	0.57
1:A:792:LYS:HD2	3:E:105:ASP:OD1	2.05	0.56
1:A:14:ILE:HD12	1:A:688:HIS:HB2	1.85	0.56
1:A:576:ALA:O	1:A:580:ASN:HB2	2.05	0.56
1:A:367:LEU:HD12	1:A:429:GLU:HG3	1.88	0.55
2:B:78:TYR:CE2	2:B:82:LEU:HD11	2.41	0.55
3:E:113:THR:HG22	3:E:129:LYS:HB2	1.87	0.54
1:A:14:ILE:CD1	1:A:688:HIS:HB2	2.37	0.54
1:A:578:ASN:ND2	1:A:578:ASN:H	2.06	0.54
1:A:238:LYS:HG2	1:A:243:ASN:HA	1.90	0.53
3:E:87:ILE:HG23	3:E:120:THR:HA	1.90	0.53

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:90:HIS:HA	2:B:92:LYS:HZ1	1.74	0.53
1:A:167:VAL:HG12	1:A:664:ILE:HD13	1.91	0.53
3:E:91:LEU:HD22	3:E:106:LEU:HD22	1.90	0.52
1:A:610:VAL:HG12	1:A:614:LYS:HD2	1.91	0.52
1:A:516:THR:HB	1:A:519:LEU:HD12	1.91	0.52
1:A:12:SER:HA	1:A:15:VAL:HG23	1.92	0.52
2:B:58:LEU:HD22	2:B:89:VAL:HA	1.92	0.52
3:E:113:THR:O	3:E:114:ASN:HB2	2.09	0.51
1:A:201:THR:HG22	1:A:205:MET:HE2	1.92	0.50
2:B:91:ASP:OD2	2:B:91:ASP:N	2.44	0.50
2:B:114:THR:HG22	2:B:117:GLN:OE1	2.13	0.49
1:A:619:PRO:O	1:A:623:GLN:HG3	2.12	0.49
2:B:35:PHE:HD2	2:B:81:TYR:CD2	2.30	0.49
3:E:89:ASN:O	3:E:93:ILE:HG13	2.14	0.48
1:A:369[A]:GLU:HG2	1:A:425:LYS:HB2	1.96	0.48
1:A:507:LYS:HE2	1:A:511:ASP:OD2	2.13	0.47
1:A:304:LYS:HE2	1:A:350:GLU:OE2	2.14	0.47
1:A:815:MET:HE1	2:B:133:GLU:OE2	2.15	0.47
1:A:634:LYS:HD3	1:A:634:LYS:HA	1.74	0.47
1:A:813:LYS:HE3	2:B:57:GLY:O	2.15	0.47
1:A:274:LYS:O	1:A:277:ILE:HD12	2.15	0.46
2:B:52:ASN:O	2:B:56:LEU:HG	2.16	0.46
2:B:65:GLU:O	2:B:69:LYS:HB2	2.16	0.46
1:A:609:LEU:HD23	1:A:609:LEU:HA	1.81	0.46
2:B:35:PHE:CE2	2:B:78:TYR:HA	2.51	0.46
1:A:585:HIS:CD2	1:A:590:ILE:HD12	2.50	0.46
3:E:49[A]:ASN:ND2	8:E:201:HOH:O	2.49	0.46
1:A:462:LYS:HE2	1:A:462:LYS:HB2	1.88	0.45
1:A:729:VAL:O	1:A:734:LYS:HE2	2.17	0.45
1:A:652:LEU:O	1:A:656:ILE:HG13	2.17	0.45
1:A:17:ARG:HG3	1:A:88:TRP:CZ2	2.53	0.44
3:E:113:THR:HG21	3:E:126:VAL:HA	1.99	0.44
2:B:50:SER:O	2:B:54:ARG:HG3	2.17	0.43
1:A:813:LYS:HD2	2:B:58:LEU:C	2.42	0.43
1:A:14:ILE:HD11	1:A:693:LEU:HD12	1.99	0.43
1:A:655:LEU:HD23	1:A:655:LEU:HA	1.82	0.43
1:A:632:LYS:HA	1:A:632:LYS:HD3	1.78	0.43
1:A:705:TYR:HB3	1:A:758:LEU:HB2	2.00	0.43
3:E:90:PHE:O	3:E:94:MET:HG2	2.18	0.43
1:A:657:ASN:O	5:A:902:GOL:H31	2.19	0.43
1:A:246:SER:HB2	8:A:1068:HOH:O	2.18	0.42

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:282:ASP:O	1:A:283:ASN:HB2	2.17	0.42
1:A:813:LYS:O	1:A:816:VAL:HG22	2.18	0.42
3:E:24[B]:GLU:HG2	3:E:27:LYS:HB2	2.00	0.42
1:A:96:PRO:HG2	1:A:119:HIS:CG	2.54	0.42
1:A:699[B]:ARG:HD3	8:A:1056:HOH:O	2.20	0.42
2:B:35:PHE:CD2	2:B:78:TYR:HD1	2.38	0.42
1:A:605:LEU:HD12	1:A:605:LEU:HA	1.91	0.42
3:E:114:ASN:ND2	3:E:121:LEU:HD13	2.34	0.42
1:A:794:LYS:HE2	1:A:798:ASN:OD1	2.18	0.42
1:A:665:ARG:HB3	1:A:690:LEU:HD13	2.02	0.42
3:E:87:ILE:O	3:E:90:PHE:HB3	2.19	0.42
1:A:16:ARG:HD2	1:A:16:ARG:HA	1.81	0.41
1:A:618:ASN:ND2	1:A:620:ILE:H	2.18	0.41
1:A:587:ILE:O	1:A:688:HIS:HE1	2.03	0.41
3:E:121:LEU:HD12	3:E:121:LEU:HA	1.91	0.41
3:E:76:VAL:HG23	8:E:219:HOH:O	2.20	0.41
1:A:785:GLU:OE2	3:E:100:ARG:HG3	2.20	0.41
1:A:795:GLN:O	1:A:799:LYS:HD3	2.20	0.41
1:A:14:ILE:HD12	1:A:688:HIS:CD2	2.57	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	807/818 (99%)	788 (98%)	19 (2%)	0	100	100
2	B	128/204 (63%)	125 (98%)	3 (2%)	0	100	100
3	E	118/134 (88%)	112 (95%)	6 (5%)	0	100	100
All	All	1053/1156 (91%)	1025 (97%)	28 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	722/725 (100%)	708 (98%)	14 (2%)	50	65
2	B	116/186 (62%)	116 (100%)	0	100	100
3	E	116/126 (92%)	114 (98%)	2 (2%)	53	68
All	All	954/1037 (92%)	938 (98%)	16 (2%)	53	68

All (16) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	27	SER
1	A	56	VAL
1	A	92	SER
1	A	157	THR
1	A	211	SER
1	A	496	ASN
1	A	515	SER
1	A	522	THR
1	A	530	VAL
1	A	540	SER
1	A	671	GLU
1	A	706	ARG
1	A	736	VAL
1	A	797	VAL
3	E	100	ARG
3	E	104	SER

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (17) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	35	GLN
1	A	58	GLN
1	A	59	GLN
1	A	125	GLN
1	A	478	ASN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	567	ASN
1	A	568	ASN
1	A	578	ASN
1	A	584	GLN
1	A	591	GLN
1	A	618	ASN
1	A	688	HIS
1	A	791	HIS
1	A	795	GLN
2	B	132	GLN
3	E	58	ASN
3	E	68	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

1 non-standard protein/DNA/RNA residue is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
1	SEP	A	19	1	8,9,10	1.63	1 (12%)	8,12,14	1.02	1 (12%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	SEP	A	19	1	-	4/5/8/10	-

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	19	SEP	P-O1P	3.53	1.61	1.50

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	19	SEP	O2P-P-OG	2.63	113.74	106.73

There are no chirality outliers.

All (4) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
1	A	19	SEP	N-CA-CB-OG
1	A	19	SEP	CB-OG-P-O2P
1	A	19	SEP	CB-OG-P-O3P
1	A	19	SEP	CB-OG-P-O1P

There are no ring outliers.

No monomer is involved in short contacts.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 4 ligands modelled in this entry, 1 is monoatomic - leaving 3 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
7	A1JT8	A	904	-	24,25,25	3.79	14 (58%)	29,35,35	3.17	10 (34%)
4	AGS	A	901	6	29,33,33	0.31	0	39,52,52	0.61	1 (2%)
5	GOL	A	902	-	5,5,5	0.28	0	5,5,5	0.43	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
7	A1JT8	A	904	-	-	2/17/18/18	0/3/3/3
4	AGS	A	901	6	-	2/21/38/38	0/3/3/3
5	GOL	A	902	-	-	0/4/4/4	-

All (14) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
7	A	904	A1JT8	C07-C06	8.90	1.52	1.38
7	A	904	A1JT8	C05-C04	7.98	1.53	1.38
7	A	904	A1JT8	C08-C03	6.91	1.52	1.38
7	A	904	A1JT8	S09-N12	5.19	1.69	1.61
7	A	904	A1JT8	C15-N16	4.47	1.40	1.34
7	A	904	A1JT8	C05-C06	-3.97	1.32	1.38
7	A	904	A1JT8	C18-C19	3.85	1.51	1.46
7	A	904	A1JT8	C08-C07	-3.46	1.32	1.38
7	A	904	A1JT8	C06-S09	3.05	1.81	1.76
7	A	904	A1JT8	O10-S09	2.98	1.46	1.43
7	A	904	A1JT8	C21-C20	2.96	1.56	1.42
7	A	904	A1JT8	O11-S09	2.86	1.46	1.43
7	A	904	A1JT8	C04-C03	-2.76	1.33	1.38
7	A	904	A1JT8	C21-C22	2.47	1.43	1.34

All (11) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
7	A	904	A1JT8	O11-S09-O10	-13.86	102.51	119.55
7	A	904	A1JT8	O10-S09-N12	4.32	113.79	107.04
7	A	904	A1JT8	C22-S23-C19	3.90	99.26	92.32
7	A	904	A1JT8	C18-N17-N16	3.55	108.65	104.67
7	A	904	A1JT8	C15-N16-N17	-3.36	107.07	112.43
7	A	904	A1JT8	C18-C19-S23	2.96	126.00	119.91
7	A	904	A1JT8	O11-S09-N12	2.72	111.28	107.04
7	A	904	A1JT8	O11-S09-C06	2.58	111.14	107.97
7	A	904	A1JT8	C01-O02-C03	-2.34	112.44	117.51
7	A	904	A1JT8	C20-C19-C18	-2.12	124.08	128.98
4	A	901	AGS	PA-O3A-PB	-2.10	125.62	132.83

There are no chirality outliers.

All (4) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
7	A	904	A1JT8	N17-C18-C19-S23
4	A	901	AGS	PG-O3B-PB-O2B
4	A	901	AGS	PA-O3A-PB-O2B
7	A	904	A1JT8	N17-C18-C19-C20


There are no ring outliers.

1 monomer is involved in 1 short contact:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	A	902	GOL	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.

Ligand A1JT8 A 904



Ligand AGS A 901



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	806/818 (98%)	0.05	44 (5%) 30 36	15, 41, 81, 105	5 (0%)
2	B	130/204 (63%)	0.68	6 (4%) 37 43	40, 64, 86, 100	0
3	E	122/134 (91%)	0.80	19 (15%) 5 5	18, 55, 99, 112	2 (1%)
All	All	1058/1156 (91%)	0.21	69 (6%) 25 28	15, 46, 87, 112	7 (0%)

All (69) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	18	VAL	8.7
1	A	631[A]	GLU	6.0
3	E	76	VAL	5.9
3	E	87	ILE	5.4
1	A	10	THR	5.4
1	A	633	GLY	5.2
1	A	11	ALA	4.9
3	E	120	THR	4.9
1	A	575	VAL	4.8
3	E	49[A]	ASN	4.8
1	A	375	LEU	4.7
1	A	17	ARG	4.6
3	E	75	ASN	4.2
1	A	632	LYS	4.1
3	E	121	LEU	4.1
2	B	29	SER	3.9
1	A	729	VAL	3.9
1	A	14	ILE	3.7
1	A	705	TYR	3.6
1	A	217	ASP	3.3
1	A	522	THR	3.2
1	A	517	ALA	3.2
2	B	91	ASP	3.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
3	E	74	ASN	3.1
1	A	16	ARG	3.1
1	A	13	LYS	3.1
1	A	714	TYR	3.0
3	E	90	PHE	3.0
3	E	96	ALA	2.9
1	A	369[A]	GLU	2.9
1	A	462	LYS	2.9
2	B	158	GLN	2.8
1	A	376	SER	2.7
1	A	577	SER	2.7
1	A	576	ALA	2.6
1	A	634	LYS	2.6
1	A	12	SER	2.6
1	A	817	ALA	2.6
1	A	415	GLY	2.6
1	A	567	ASN	2.6
3	E	134	ILE	2.6
2	B	33	ILE	2.5
1	A	93	GLN	2.5
1	A	630	ILE	2.5
1	A	521	TYR	2.5
1	A	520	LYS	2.4
3	E	111	GLN	2.4
1	A	725	GLU	2.4
1	A	728	SER	2.4
1	A	477	LYS	2.4
1	A	15	VAL	2.4
3	E	2	ALA	2.3
1	A	26	LYS	2.3
3	E	88	ASP	2.3
1	A	370	LYS	2.3
1	A	741	LEU	2.2
3	E	98	ASP	2.2
1	A	22	GLU	2.2
2	B	156	ILE	2.2
1	A	565	LYS	2.2
3	E	103	GLU	2.1
3	E	95	LYS	2.1
3	E	117	ASN	2.1
1	A	416	GLY	2.1
1	A	730	GLU	2.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
3	E	105	ASP	2.1
1	A	568	ASN	2.0
3	E	89	ASN	2.0
2	B	89	VAL	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
1	SEP	A	19	10/11	0.89	0.14	54,71,82,89	0

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
5	GOL	A	902	6/6	0.87	0.17	45,52,65,69	0
7	A1JT8	A	904	23/23	0.98	0.06	19,33,41,43	0
6	MG	A	903	1/1	0.99	0.02	19,19,19,19	0
4	AGS	A	901	31/31	0.99	0.04	19,23,28,32	0




The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around A1JT8 A 904:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around AGS A 901:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.