



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 19, 2026 – 12:24 pm GMT

PDB ID : 9TI1 / pdb_00009ti1
Title : Crystal structure of the zinc-containing Phosphotriesterase dPTE2-H55
Authors : Manser, B.P.; Deliz Liang, A.
Deposited on : 2025-12-04
Resolution : 1.55 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 1.8.4, CSD as541be (2020)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.48


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.55 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	164625	1935 (1.56-1.56)
Clashscore	180529	2073 (1.56-1.56)
Ramachandran outliers	177936	2037 (1.56-1.56)
Sidechain outliers	177891	2034 (1.56-1.56)
RSRZ outliers	164620	1935 (1.56-1.56)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	342	
1	B	342	

2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 10452 atoms, of which 4986 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Parathion hydrolase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace	
			Total	C	H	N	O				S
1	A	328	5031	1594	2500	447	482	8	0	0	0
1	B	327	5000	1586	2480	443	483	8	0	0	0

There are 58 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	1	SER	-	expression tag	UNP P0A434
A	2	ASN	-	expression tag	UNP P0A434
A	3	ILE	-	expression tag	UNP P0A434
A	4	GLY	-	expression tag	UNP P0A434
A	5	SER	-	expression tag	UNP P0A434
A	6	GLY	-	expression tag	UNP P0A434
A	7	ILE	-	expression tag	UNP P0A434
A	8	THR	-	expression tag	UNP P0A434
A	9	ASN	-	expression tag	UNP P0A434
A	10	SER	-	expression tag	UNP P0A434
A	31	MET	THR	engineered mutation	UNP P0A434
A	54	ASP	LYS	engineered mutation	UNP P0A434
A	88	GLU	SER	engineered mutation	UNP P0A434
A	95	GLU	ARG	engineered mutation	UNP P0A434
A	159	ARG	LEU	engineered mutation	UNP P0A434
A	162	ARG	LYS	engineered mutation	UNP P0A434
A	180	ASP	ALA	engineered mutation	UNP P0A434
A	191	ASP	ALA	engineered mutation	UNP P0A434
A	199	ASP	SER	engineered mutation	UNP P0A434
A	215	ASP	SER	engineered mutation	UNP P0A434
A	246	ALA	SER	engineered mutation	UNP P0A434
A	251	LEU	ILE	engineered mutation	UNP P0A434
A	270	ALA	MET	engineered mutation	UNP P0A434
A	271	ASP	LYS	engineered mutation	UNP P0A434
A	320	ASP	GLN	engineered mutation	UNP P0A434

Continued on next page...


Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
A	324	GLU	ALA	engineered mutation	UNP P0A434
A	325	THR	GLY	engineered mutation	UNP P0A434
A	327	MET	THR	engineered mutation	UNP P0A434
A	329	ASP	THR	engineered mutation	UNP P0A434
B	1	SER	-	expression tag	UNP P0A434
B	2	ASN	-	expression tag	UNP P0A434
B	3	ILE	-	expression tag	UNP P0A434
B	4	GLY	-	expression tag	UNP P0A434
B	5	SER	-	expression tag	UNP P0A434
B	6	GLY	-	expression tag	UNP P0A434
B	7	ILE	-	expression tag	UNP P0A434
B	8	THR	-	expression tag	UNP P0A434
B	9	ASN	-	expression tag	UNP P0A434
B	10	SER	-	expression tag	UNP P0A434
B	31	MET	THR	engineered mutation	UNP P0A434
B	54	ASP	LYS	engineered mutation	UNP P0A434
B	88	GLU	SER	engineered mutation	UNP P0A434
B	95	GLU	ARG	engineered mutation	UNP P0A434
B	159	ARG	LEU	engineered mutation	UNP P0A434
B	162	ARG	LYS	engineered mutation	UNP P0A434
B	180	ASP	ALA	engineered mutation	UNP P0A434
B	191	ASP	ALA	engineered mutation	UNP P0A434
B	199	ASP	SER	engineered mutation	UNP P0A434
B	215	ASP	SER	engineered mutation	UNP P0A434
B	246	ALA	SER	engineered mutation	UNP P0A434
B	251	LEU	ILE	engineered mutation	UNP P0A434
B	270	ALA	MET	engineered mutation	UNP P0A434
B	271	ASP	LYS	engineered mutation	UNP P0A434
B	320	ASP	GLN	engineered mutation	UNP P0A434
B	324	GLU	ALA	engineered mutation	UNP P0A434
B	325	THR	GLY	engineered mutation	UNP P0A434
B	327	MET	THR	engineered mutation	UNP P0A434
B	329	ASP	THR	engineered mutation	UNP P0A434

- Molecule 2 is ZINC ION (CCD ID: ZN) (formula: Zn) (labeled as "Ligand of Interest" by depositor).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	2	Total Zn 2 2	0	0
2	B	2	Total Zn 2 2	0	0

- Molecule 3 is 1,2-ETHANEDIOL (CCD ID: EDO) (formula: $C_2H_6O_2$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	H	O		
3	B	1	10	2	6	2	0	0


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	O		
4	A	215	215	215	0	0
4	B	192	192	192	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Parathion hydrolase



- Molecule 1: Parathion hydrolase



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	63.49Å 69.98Å 154.16Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	19.69 – 1.55 19.69 – 1.55	Depositor EDS
% Data completeness (in resolution range)	76.3 (19.69-1.55) 76.7 (19.69-1.55)	Depositor EDS
R_{merge}	0.28	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.40 (at 1.55Å)	Xtrriage
Refinement program	PHENIX 1.21.2_5419	Depositor
R, R_{free}	0.177 , 0.187 0.177 , 0.187	Depositor DCC
R_{free} test set	3804 reflections (3.77%)	wwPDB-VP
Wilson B-factor (Å ²)	11.9	Xtrriage
Anisotropy	0.049	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.43 , 38.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	10452	wwPDB-VP
Average B, all atoms (Å ²)	16.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.70% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: ZN, KCX, EDO

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.49	0/2566	0.70	0/3486
1	B	0.47	0/2555	0.69	0/3472
All	All	0.48	0/5121	0.69	0/6958

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	18	ARG	Sidechain

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2531	2500	2508	12	0
1	B	2520	2480	2488	13	0
2	A	2	0	0	0	0
2	B	2	0	0	0	0
3	B	4	6	6	0	0
4	A	215	0	0	3	0
4	B	192	0	0	3	0
All	All	5466	4986	5002	25	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (25) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:88:GLU:OE2	4:B:501:HOH:O	2.04	0.76
1:B:278:ASP:OD2	4:B:502:HOH:O	2.06	0.72
1:A:278:ASP:OD2	4:A:501:HOH:O	2.14	0.66
1:B:44:ARG:HG3	4:B:663:HOH:O	2.10	0.52
1:A:264:LEU:HD23	1:A:267:GLN:OE1	2.11	0.50
1:B:34:HIS:HB2	1:B:280:LEU:HB3	1.94	0.50
1:A:310:ILE:HB	1:A:311:PRO:HD3	1.95	0.49
1:B:147:VAL:HG21	1:B:161:LEU:HD23	1.95	0.49
1:A:291:MET:HE3	1:A:291:MET:HA	1.96	0.48
1:B:83:ILE:HD11	1:B:108:TRP:CG	2.49	0.48
1:B:310:ILE:HB	1:B:311:PRO:HD3	1.96	0.48
1:A:330:ASN:HB2	1:A:331:PRO:HD3	1.97	0.47
1:A:34:HIS:O	1:A:280:LEU:HA	2.15	0.45
1:A:58:GLU:HG2	1:A:62:ARG:NH1	2.32	0.45
1:A:279:TRP:CH2	1:A:298:ASN:HB3	2.51	0.45
1:A:83:ILE:HG22	1:A:83:ILE:O	2.17	0.45
1:B:179:THR:HB	1:B:185:ASP:HB2	1.98	0.44
1:B:330:ASN:HB2	1:B:331:PRO:HD3	2.00	0.44
1:B:280:LEU:HD12	1:B:291:MET:HE1	2.01	0.43
1:B:83:ILE:O	1:B:83:ILE:HG22	2.20	0.42
1:A:179:THR:HB	1:A:185:ASP:HB2	2.00	0.42
4:A:654:HOH:O	1:B:112:PRO:HG3	2.18	0.41
1:A:233:PRO:HB3	1:A:304:PHE:CD2	2.56	0.41
1:B:34:HIS:O	1:B:280:LEU:HA	2.20	0.41
1:A:223:ARG:HG3	4:A:561:HOH:O	2.21	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	325/342 (95%)	318 (98%)	7 (2%)	0	100	100
1	B	324/342 (95%)	316 (98%)	8 (2%)	0	100	100
All	All	649/684 (95%)	634 (98%)	15 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	265/275 (96%)	263 (99%)	2 (1%)	79	63
1	B	264/275 (96%)	262 (99%)	2 (1%)	79	63
All	All	529/550 (96%)	525 (99%)	4 (1%)	79	63

All (4) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	276	SER
1	A	280	LEU
1	B	276	SER
1	B	280	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
1	KCX	B	146	2,1	9,11,12	0.70	0	5,12,14	1.66	1 (20%)
1	KCX	A	146	2,1	9,11,12	0.50	0	5,12,14	0.78	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	KCX	B	146	2,1	-	0/9/10/12	-
1	KCX	A	146	2,1	-	0/9/10/12	-

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	146	KCX	CE-NZ-CX	-2.86	117.30	121.89

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 5 ligands modelled in this entry, 4 are monoatomic - leaving 1 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	EDO	B	401	-	3,3,3	0.26	0	2,2,2	0.31	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	EDO	B	401	-	-	0/1/1/1	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	327/342 (95%)	-0.43	1 (0%) 90 93	7, 13, 24, 37	0
1	B	326/342 (95%)	-0.19	2 (0%) 85 90	9, 16, 30, 45	0
All	All	653/684 (95%)	-0.31	3 (0%) 87 91	7, 14, 27, 45	0

All (3) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	338	THR	3.7
1	A	339	LEU	2.8
1	B	251	LEU	2.3

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q < 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
1	KCX	B	146	12/13	0.92	0.08	7,9,13,16	0
1	KCX	A	146	12/13	0.93	0.07	5,7,12,13	0

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	EDO	B	401	4/4	0.87	0.14	21,25,30,33	0
2	ZN	B	402	1/1	0.99	0.02	10,10,10,10	0
2	ZN	A	402	1/1	0.99	0.02	13,13,13,13	0
2	ZN	B	403	1/1	1.00	0.01	13,13,13,13	0
2	ZN	A	401	1/1	1.00	0.01	10,10,10,10	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around ZN B 402:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around ZN A 402:


$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around ZN B 403:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.5 Other polymers [i](#)

There are no such residues in this entry.