Full wwPDB X-ray Structure Validation Report (i)

Oct 26, 2026 – 06:08 pm GMT

PDB ID	:	9TMM / pdb_00009tmm
Title	:	Structure of the TCR-A targeting GPC3 [FLAELAYDL] peptide bound to HLA-A*02:01
Authors	:	Oyon-Olea, L.; Lopez-Sagaseta, J.
Deposited on	:	2025-12-15
Resolution	:	2.30 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org*
A user guide is available at
https://www.wwpdb.org/validation/2017/XrayValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at
http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (i)) were used in the production of this report:


MolProbity	:	4-5-2 with Phenix2.2
Mogul	:	1.8.4, CSD as541be (2020)
Xtriage (Phenix)	:	2.2
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.015 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.51

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:
X-RAY DIFFRACTION

The reported resolution of this entry is 2.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	6319 (2.30-2.30)
Clashscore	190562	6919 (2.30-2.30)
Ramachandran outliers	187476	6854 (2.30-2.30)
Sidechain outliers	187428	6854 (2.30-2.30)
RSRZ outliers	180081	6325 (2.30-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	280	4% 87% 12%
2	B	101	17% 83% 16%
3	C	212	12% 71% 5% 23%
4	D	244	17% 92% 7%
5	P	9	89% 11%

2 Entry composition (i)

There are 9 unique types of molecules in this entry. The entry contains 6169 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called MHC class I.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	277	Total	C	N	O	S	0	0	0
			2214	1386	402	417	9			

- Molecule 2 is a protein called Beta-2-microglobulin.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	100	Total	C	N	O	S	0	0	0
			765	483	133	146	3			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	0	MET	-	initiating methionine	UNP P61769
B	1	GLY	-	expression tag	UNP P61769

- Molecule 3 is a protein called TCR-A alpha chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	C	163	Total	C	N	O	S	0	0	0
			1179	750	195	228	6			


- Molecule 4 is a protein called TCR-A beta chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
4	D	244	Total	C	N	O	S	0	0	0
			1751	1110	308	325	8			

- Molecule 5 is a protein called Glypican-3 beta subunit.


Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
5	P	9	Total	C	N	O	0	0	0
			74	51	9	14			

- Molecule 6 is 1,2-ETHANEDIOL (CCD ID: EDO) (formula: $C_2H_6O_2$).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	A	1	Total C O 4 2 2	0	0
6	A	1	Total C O 4 2 2	0	0
6	A	1	Total C O 4 2 2	0	0
6	A	1	Total C O 4 2 2	0	0
6	A	1	Total C O 4 2 2	0	0
6	A	1	Total C O 4 2 2	0	0
6	A	1	Total C O 4 2 2	0	0
6	B	1	Total C O 4 2 2	0	0
6	C	1	Total C O 4 2 2	0	0
6	C	1	Total C O 4 2 2	0	0

- Molecule 7 is BORIC ACID (CCD ID: BO3) (formula: BH_3O_3).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
7	C	1	Total B O 4 1 3	0	0

- Molecule 8 is SODIUM ION (CCD ID: NA) (formula: Na).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
8	C	1	Total Na 1 1	0	0

- Molecule 9 is water.


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
9	A	70	Total O 70 70	0	0
9	B	14	Total O 14 14	0	0
9	C	29	Total O 29 29	0	0
9	D	26	Total O 26 26	0	0
9	P	2	Total O 2 2	0	0

3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: MHC class I

Chain A:




- Molecule 2: Beta-2-microglobulin

Chain B:




- Molecule 3: TCR-A alpha chain


Chain C:



- Molecule 4: TCR-A beta chain

Chain D:





• Molecule 5: Glypican-3 beta subunit

Chain P:





4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 65 2 2	Depositor
Cell constants a, b, c, α, β, γ	78.64Å 78.64Å 574.49Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	49.42 – 2.30 49.42 – 2.30	Depositor EDS
% Data completeness (in resolution range)	99.7 (49.42-2.30) 99.7 (49.42-2.30)	Depositor EDS
R_{merge}	0.28	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) >$ [1]	2.46 (at 2.29Å)	Xtriage
Refinement program	PHENIX 1.21.2_5419	Depositor
R, R_{free}	0.244 , 0.278 0.244 , 0.278	Depositor DCC
R_{free} test set	2502 reflections (5.13%)	wwPDB-VP
Wilson B-factor (Å^2)	36.4	Xtriage
Anisotropy	0.111	Xtriage
Bulk solvent k_{sol}(e/Å^3), B_{sol}(Å^2)	0.36 , 57.6	EDS
L-test for twinning[2]	$< \|L\| > = 0.40$, $< L^2 > = 0.23$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o,F_c correlation	0.91	EDS
Total number of atoms	6169	wwPDB-VP
Average B, all atoms (Å^2)	49.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.24% of the height of the origin peak. No significant pseudotranslation is detected.*

[1] Intensities estimated from amplitudes.

[2] Theoretical values of $< |L| >$, $< L^2 >$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: NA, BO3, EDO

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\#\|Z\| > 5$	RMSZ	$\#\|Z\| > 5$
1	A	0.10	0/2279	0.26	0/3102
2	B	0.10	0/788	0.28	0/1076
3	C	0.11	0/1208	0.27	0/1654
4	D	0.09	0/1802	0.27	0/2480
5	P	0.12	0/75	0.25	0/101
All	All	0.10	0/6152	0.27	0/8413

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2214	0	2025	24	0
2	B	765	0	646	11	0
3	C	1179	0	1020	7	0
4	D	1751	0	1475	10	0
5	P	74	0	73	1	0
6	A	28	0	42	2	0
6	B	4	0	6	1	0
6	C	8	0	12	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
7	C	4	0	3	0	0
8	C	1	0	0	0	0
9	A	70	0	0	0	0
9	B	14	0	0	0	0
9	C	29	0	0	0	0
9	D	26	0	0	0	0
9	P	2	0	0	0	0
All	All	6169	0	5302	48	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (48) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:C:163:MET:HE1	3:C:166:MET:HE3	1.49	0.92
1:A:97:MET:HE2	6:B:201:EDO:H12	1.78	0.64
1:A:104:SER:H	6:A:307:EDO:H12	1.63	0.62
2:B:97:ASP:HB3	2:B:100:MET:HB2	1.83	0.61
2:B:38:VAL:HG22	2:B:83:VAL:HG22	1.82	0.60
3:C:15:GLU:HG3	3:C:82:HIS:HA	1.84	0.59
1:A:21:PHE:HE1	1:A:73:HIS:HD2	1.50	0.59
1:A:233:ARG:HH11	2:B:9:GLN:HE21	1.50	0.58
1:A:96:ARG:HD3	1:A:115:TYR:CZ	2.38	0.58
1:A:229:LEU:HD11	1:A:242:LYS:HE3	1.87	0.56
1:A:157:ALA:HB2	3:C:52:SER:HB3	1.89	0.54
1:A:34:ARG:HH11	1:A:47:ARG:NH2	2.06	0.54
4:D:19:VAL:HG23	4:D:80:VAL:HG13	1.93	0.51
1:A:13:ARG:HB3	1:A:16:ARG:HB3	1.90	0.51
1:A:200:LEU:HD12	1:A:248:VAL:HG21	1.91	0.51
4:D:156:GLU:HB2	4:D:213:GLN:HB3	1.91	0.51
4:D:123:PRO:HB3	4:D:150:PHE:HB3	1.94	0.50
1:A:76:ASP:HB3	5:P:9:LEU:HD12	1.94	0.49
4:D:48:PHE:CD2	4:D:57:GLU:HB3	2.48	0.48
1:A:21:PHE:CE1	1:A:73:HIS:HD2	2.32	0.47
3:C:163:MET:HE2	3:C:166:MET:HB2	1.95	0.47
1:A:169:ARG:CZ	6:A:302:EDO:H12	2.45	0.47
4:D:35:TRP:NE1	4:D:78:LEU:HB2	2.30	0.47
2:B:25:ASN:HB3	2:B:66:LEU:HD11	1.98	0.45
1:A:43:ARG:HD2	1:A:63:THR:HG21	1.99	0.45
1:A:233:ARG:HH11	2:B:9:GLN:NE2	2.14	0.45

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:C:83:TRP:CD1	3:C:111:PRO:HD3	2.52	0.45
2:B:7:LYS:NZ	2:B:30:GLY:HA3	2.32	0.45
3:C:82:HIS:O	3:C:109:VAL:HG11	2.18	0.44
2:B:42:LYS:HB2	2:B:47:ILE:HD11	1.99	0.44
4:D:217:LEU:O	4:D:231:VAL:HA	2.18	0.43
4:D:154:HIS:HB3	4:D:215:TYR:HB2	2.01	0.43
1:A:7:PHE:HB2	1:A:24:VAL:HG23	2.01	0.43
1:A:126:LYS:HD3	1:A:131:SER:OG	2.19	0.43
2:B:74:THR:HG23	2:B:77:ASP:H	1.84	0.43
1:A:11:VAL:HG22	1:A:93:THR:HG23	2.02	0.42
1:A:233:ARG:HD2	2:B:11:TYR:CE1	2.55	0.42
2:B:71:PHE:HD1	2:B:79:TYR:CZ	2.38	0.41
2:B:42:LYS:HE3	2:B:79:TYR:OH	2.21	0.41
4:D:44:LEU:HD13	4:D:44:LEU:HA	1.92	0.41
1:A:27:VAL:HG11	1:A:178:LEU:HD13	2.02	0.41
1:A:48:ALA:O	1:A:51:ILE:HG22	2.20	0.41
4:D:130:PRO:HG2	4:D:141:ALA:HB1	2.02	0.41
1:A:34:ARG:HG2	1:A:47:ARG:HG2	2.03	0.41
1:A:108:PHE:CZ	1:A:160:GLU:HG2	2.56	0.41
1:A:217:GLN:HG2	1:A:222:ASP:HA	2.02	0.40
3:C:34:PHE:HB2	3:C:92:VAL:HB	2.04	0.40
4:D:229:LYS:HA	4:D:230:PRO:HD3	1.94	0.40

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	275/280 (98%)	271 (98%)	4 (2%)	0	100	100
2	B	98/101 (97%)	97 (99%)	1 (1%)	0	100	100
3	C	157/212 (74%)	153 (98%)	4 (2%)	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
4	D	242/244 (99%)	242 (100%)	0	0	100	100
5	P	7/9 (78%)	6 (86%)	1 (14%)	0	100	100
All	All	779/846 (92%)	769 (99%)	10 (1%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	220/235 (94%)	220 (100%)	0	100	100
2	B	74/95 (78%)	74 (100%)	0	100	100
3	C	112/189 (59%)	109 (97%)	3 (3%)	39	58
4	D	150/212 (71%)	148 (99%)	2 (1%)	61	77
5	P	7/7 (100%)	7 (100%)	0	100	100
All	All	563/738 (76%)	558 (99%)	5 (1%)	70	84

All (5) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
3	C	48	LEU
3	C	109	VAL
3	C	159	CYS
4	D	44	LEU
4	D	80	VAL

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (12) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	73	HIS
2	B	9	GLN
2	B	14	HIS

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
2	B	18	ASN
2	B	43	ASN
2	B	52	HIS
3	C	38	GLN
3	C	45	GLN
3	C	58	GLN
3	C	97	ASN
4	D	38	GLN
4	D	82	ASN

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no oligosaccharides in this entry.

5.6 Ligand geometry (i)

Of 12 ligands modelled in this entry, 1 is monoatomic - leaving 11 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\#\|Z\| > 2$	Counts	RMSZ	$\#\|Z\| > 2$
7	BO3	C	302	-	3,3,3	0.11	0	3,3,3	1.24	0
6	EDO	A	304	-	3,3,3	0.27	0	2,2,2	0.31	0
6	EDO	A	306	-	3,3,3	0.27	0	2,2,2	0.30	0
6	EDO	B	201	-	3,3,3	0.27	0	2,2,2	0.29	0
6	EDO	A	301	-	3,3,3	0.27	0	2,2,2	0.31	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\#\|Z\|>2$	Counts	RMSZ	$\#\|Z\|>2$
6	EDO	A	303	-	3,3,3	0.26	0	2,2,2	0.31	0
6	EDO	C	303	-	3,3,3	0.25	0	2,2,2	0.51	0
6	EDO	A	302	-	3,3,3	0.27	0	2,2,2	0.25	0
6	EDO	A	307	-	3,3,3	0.27	0	2,2,2	0.29	0
6	EDO	A	305	-	3,3,3	0.27	0	2,2,2	0.32	0
6	EDO	C	301	-	3,3,3	0.27	0	2,2,2	0.36	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
6	EDO	A	304	-	-	0/1/1/1	-
6	EDO	A	306	-	-	0/1/1/1	-
6	EDO	B	201	-	-	1/1/1/1	-
6	EDO	A	301	-	-	0/1/1/1	-
6	EDO	A	303	-	-	0/1/1/1	-
6	EDO	C	303	-	-	1/1/1/1	-
6	EDO	A	302	-	-	0/1/1/1	-
6	EDO	A	307	-	-	0/1/1/1	-
6	EDO	A	305	-	-	0/1/1/1	-
6	EDO	C	301	-	-	0/1/1/1	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (2) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
6	B	201	EDO	O1-C1-C2-O2
6	C	303	EDO	O1-C1-C2-O2

There are no ring outliers.

3 monomers are involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
6	B	201	EDO	1	0
6	A	302	EDO	1	0
6	A	307	EDO	1	0

5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95^{th} percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å^2)	Q<0.9
1	A	277/280 (98%)	0.38	10 (3%) 46 48	26, 42, 65, 90	0
2	B	100/101 (99%)	1.13	17 (17%) 4 5	35, 61, 91, 96	0
3	C	163/212 (76%)	0.73	26 (15%) 5 5	25, 41, 80, 97	0
4	D	244/244 (100%)	0.97	42 (17%) 4 4	26, 56, 85, 113	1 (0%)
5	P	9/9 (100%)	0.14	0 100 100	26, 29, 40, 46	0
All	All	793/846 (93%)	0.72	95 (11%) 9 9	25, 47, 82, 113	1 (0%)

All (95) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
4	D	244	ASP	4.9
3	C	129	SER	4.8
4	D	218	SER	4.5
4	D	220	ASN	4.5
3	C	139	PHE	4.5
2	B	72	THR	4.2
4	D	182	ALA	4.2
3	C	83	TRP	4.2
3	C	113	ILE	4.1
4	D	16	GLY	4.1
4	D	217	LEU	4.0
3	C	140	ASP	4.0
3	C	128	SER	3.8
4	D	181	PRO	3.8
3	C	177	SER	3.7
1	A	195	ASP	3.7
4	D	10	TRP	3.6
4	D	183	LEU	3.6
4	D	114	LEU	3.5
4	D	219	GLU	3.5

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
4	D	223	TRP	3.5
2	B	73	PRO	3.5
2	B	21	SER	3.5
1	A	226	ASP	3.5
3	C	127	LYS	3.4
3	C	173	ALA	3.4
3	C	141	SER	3.2
2	B	74	THR	3.2
4	D	178	LYS	3.1
3	C	161	LEU	3.1
4	D	226	ASP	3.1
2	B	20	LYS	3.0
3	C	120	VAL	2.9
1	A	196	HIS	2.9
4	D	215	TYR	2.9
1	A	224	THR	2.9
3	C	168	PHE	2.8
2	B	78	GLU	2.8
2	B	69	THR	2.8
2	B	71	PHE	2.8
4	D	224	THR	2.8
2	B	1	GLY	2.8
2	B	70	GLU	2.8
3	C	119	ALA	2.8
4	D	14	PRO	2.7
4	D	227	ARG	2.7
2	B	41	LEU	2.7
3	C	121	TYR	2.7
3	C	170	SER	2.7
4	D	188	TYR	2.7
4	D	58	VAL	2.7
4	D	138	THR	2.7
4	D	186	SER	2.7
4	D	15	ARG	2.6
2	B	93	ILE	2.6
4	D	243	ALA	2.6
1	A	192	ALA	2.6
3	C	133	VAL	2.6
4	D	232	THR	2.5
2	B	2	ILE	2.5
2	B	79	TYR	2.5
3	C	115	ASN	2.5

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
4	D	13	VAL	2.4
1	A	111	GLY	2.4
4	D	117	LEU	2.4
4	D	17	GLN	2.4
4	D	221	ASP	2.4
4	D	1	VAL	2.4
4	D	19	VAL	2.3
4	D	234	ILE	2.3
4	D	189	ALA	2.3
3	C	172	SER	2.3
4	D	180	GLN	2.3
4	D	185	ASP	2.3
2	B	52	HIS	2.3
4	D	184	GLN	2.2
4	D	119	ASN	2.2
1	A	229	LEU	2.2
3	C	117	ASP	2.2
1	A	275	PRO	2.2
2	B	45	GLU	2.2
3	C	174	VAL	2.1
3	C	136	PHE	2.1
3	C	154	TYR	2.1
4	D	118	LYS	2.1
1	A	193	VAL	2.1
4	D	200	PHE	2.1
3	C	175	ALA	2.1
2	B	75	GLU	2.1
4	D	116	ASP	2.1
1	A	250	SER	2.0
4	D	61	LEU	2.0
4	D	135	ILE	2.0
3	C	163	MET	2.0
3	C	112	ASN	2.0

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

There are no oligosaccharides in this entry.

6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å^2)	Q<0.9
6	EDO	A	301	4/4	0.82	0.21	64,73,78,90	0
6	EDO	A	307	4/4	0.86	0.17	59,65,68,70	0
6	EDO	A	305	4/4	0.87	0.12	45,47,53,53	0
7	BO3	C	302	4/4	0.88	0.23	44,52,53,65	0
6	EDO	B	201	4/4	0.89	0.16	51,53,54,57	0
6	EDO	A	303	4/4	0.89	0.13	44,47,48,56	0
8	NA	C	304	1/1	0.89	0.11	52,52,52,52	0
6	EDO	A	302	4/4	0.90	0.16	49,54,55,62	0
6	EDO	C	303	4/4	0.92	0.12	46,51,51,57	0
6	EDO	A	306	4/4	0.92	0.13	61,64,65,68	0
6	EDO	C	301	4/4	0.92	0.12	36,37,44,57	0
6	EDO	A	304	4/4	0.93	0.15	35,39,45,53	0

6.5 Other polymers (i)

There are no such residues in this entry.