Full wwPDB EM Validation Report (i)

Oct 5, 2026 – 12:16 PM JST

PDB ID	:	9UD0 / pdb_00009ud0
EMDB ID	:	EMD-64057
Title	:	Cryo-EM structure of R2Tg bound to template RNA and target DNA in the target-primed reverse transcription initiation state (state 1, Myb domain focused)
Authors	:	Komatsu, D.; Hiraizumi, M.; Fujimatsu, K.; Yamada, S.; Yamashita, K.; Nishimasu, H.
Deposited on	:	2025-04-06
Resolution	:	3.14 Å(reported)

This is a Full wwPDB EM Validation Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org*
A user guide is available at
https://www.wwpdb.org/validation/2017/EMValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at
http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (i)) were used in the production of this report:

EMDB validation analysis	:	0.0.1.dev133
MolProbity	:	4-5-2 with Phenix2.2
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
EM percentile statistics	:	202505.v01 (Using data in the EMDB archive up until May 2025)
MapQ	:	1.9.13
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.51

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:
ELECTRON MICROSCOPY

The reported resolution of this entry is 3.14 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.





Metric	Whole archive (#Entries)	EM structures (#Entries)	Similar EM resolution (#Entries, resolution range(Å))
Clashscore	229148	23984	-
Ramachandran outliers	224038	23583	-
Sidechain outliers	223484	23102	-
Q-score	-	25397	14483 (2.64 - 3.64)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion < 40%). The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	1287	5% • 95%
2	C	80	21% 5% • 71%
3	D	50	32% 14% 54%

2 Entry composition (i)

There are 3 unique types of molecules in this entry. The entry contains 1517 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Reverse transcriptase-like protein.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
1	A	70	574	352	113	108	1	0	0

- Molecule 2 is a DNA chain called 28S DNA top strand.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	P		
2	C	23	484	226	101	134	23	0	0

- Molecule 3 is a DNA chain called 28S DNA bottom strand, 3' side.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	P		
3	D	23	459	219	75	142	23	0	0

3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and atom inclusion in map density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red diamond above a residue indicates a poor fit to the EM map for this residue (all-atom inclusion < 40%). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Reverse transcriptase-like protein

Chain A: 5% 95%



MET ILE ASP PRO GLY GLU SER GLU LYS TYR LEU GLY LEU GLN PHE ASP PRO TRP ILE GLY ILE ALA ARG SER GLY LEU SER THR LYS LEU ASP PHE TRP LEU GLN ARG ILE ASP GLN ALA PRO LEU LYS PRO LEU GLN LYS THR ASP ILE LEU LYS THR TYR THR ILE PRO ARG LEU ILE

TYR ILE ALA ASP HIS SER GLU VAL LYS THR ALA LEU LEU GLU THR LEU ASP GLN LYS ILE ARG THR ALA VAL LYS GLU TRP LEU HIS LEU PRO PRO CYS THR CYS ASP ALA ILE LEU TYR SER SER THR ARG ASP GLY GLY LEU GLY ILE THR LYS LEU ALA GLY LEU ILE PRO SER VAL

GLN ALA ARG ARG LEU HIS ARG ILE ALA GLN SER SER ASP ASP THR MET LYS CYS PHE MET GLU LYS GLU LYS MET GLU GLN LEU HIS LYS LYS LEU TRP ILE GLN ALA GLY GLY ASP ARG GLU ASN ILE PRO SER ILE TRP GLU ALA PRO PRO SER SER GLU PRO PRO ASN ASN VAL SER

THR ASN SER GLU TRP GLU ALA PRO THR GLN LYS ASP LYS PHE PRO LYS PRO CYS ASN TRP ARG LYS ASN GLU PHE LYS LYS TRP THR LYS LEU ALA SER GLN GLY ARG GLY ILE VAL ASN PHE GLU ARG ASP LYS ILE SER ASN HIS TRP ILE GLN TYR TYR ARG ARG ILE PRO HIS ARG

LYS LEU LEU THR ALA LEU GLN LEU ARG ALA ASN VAL TYR PRO THR ARG GLU PHE LEU ALA ARG GLY ARG GLN ASP GLN TYR ILE LYS ALA CYS ARG HIS CYS ASP ALA ASP ILE GLU SER CYS ALA HIS ILE ILE GLY ASN CYS PRO VAL THR GLN ASP ALA ARG ILE LYS ARG HIS ASN

TYR ILE CYS GLU LEU LEU LEU GLU GLU ALA LYS LYS LYS ASP TRP VAL VAL PHE LYS GLU PRO HIS ILE ARG ASP SER ASN LYS GLU LEU TYR LYS PRO ASP LEU ILE PHE VAL LYS ASP ALA ARG ALA LEU VAL VAL ASP VAL THR VAL ARG TYR GLU ALA ALA LYS SER SER LEU GLU

GLU ALA ALA ALA GLU LYS VAL ARG LYS TYR LYS HIS LEU GLU THR GLU VAL ARG HIS LEU THR ASN ALA LYS ASP VAL THR PHE VAL GLY PHE PRO LEU GLY ALA ARG GLY LYS TRP HIS GLN ASP ASN PHE LYS LEU LEU THR GLU LEU GLY LEU SER LYS SER ARG GLN VAL LYS MET

ALA GLU THR PHE SER THR VAL ALA LEU PHE SER SER VAL ASP ILE VAL HIS MET PHE ALA SER ARG ALA ARG LYS SER MET VAL MET GLY GLY SER ASP TYR LYS ASP HIS ASP GLY ASP TYR LYS ASP HIS ASP ILE ASP TYR LYS ASP ASP ASP ASP LYS

- Molecule 2: 28S DNA top strand

Chain C:

DA DT DT DC DA DA T7 G16 G24 C25 G26 G27 G28 A29 DG DT DA DA DC DT DA DT DG DA DC DT DC DT DC DT DT DA DA DG DG DT DA DG DC DC DA DA DA DT DG DC DC DT DC DG DT DC DA DT DC DT DA DA

DT DT DA DG DT DG DA

- Molecule 3: 28S DNA bottom strand, 3' side

Chain D:

DC DC DT DT DA DA DG DA DG DA DG DT DC DA DT DA DG DT DT DA DC T52 C53 C54 C55 G56 C57 C58 A63 C64 A74 DT DT DG DA DA DT

4 Experimental information (i)

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, Not provided	
Number of particles used	168252	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	PHASE FLIPPING AND AMPLITUDE CORRECTION	Depositor
Microscope	TFS KRIOS	Depositor
Voltage (kV)	300	Depositor
Electron dose ($e^-/Å^2$)	49.6	Depositor
Minimum defocus (nm)	800	Depositor
Maximum defocus (nm)	1800	Depositor
Magnification	Not provided	
Image detector	GATAN K3 (6k x 4k)	Depositor
Maximum map value	0.346	Depositor
Minimum map value	-0.129	Depositor
Average map value	-0.000	Depositor
Map value standard deviation	0.004	Depositor
Recommended contour level	0.05	Depositor
Map size (Å)	298.8, 298.8, 298.8	wwPDB
Map dimensions	300, 300, 300	wwPDB
Map angles (°)	90.0, 90.0, 90.0	wwPDB
Pixel spacing (Å)	0.99599993, 0.99599993, 0.99599993	Depositor

5 Model quality (i)

5.1 Standard geometry (i)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\#\|Z\|>5$	RMSZ	$\#\|Z\|>5$
1	A	0.62	0/579	1.12	1/773 (0.1%)
2	C	0.78	0/546	1.53	8/843 (0.9%)
3	D	0.83	0/510	1.61	10/782 (1.3%)
All	All	0.75	0/1635	1.44	19/2398 (0.8%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand.A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

There are no bond length outliers.

All (19) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})
3	D	52	DT	O3'-P-O5'	-8.16	88.49	104.00
3	D	53	DC	O5'-C5'-C4'	7.37	115.30	109.40
3	D	53	DC	O3'-P-O5'	-7.11	90.49	104.00
2	C	29	DA	O5'-C5'-C4'	6.69	114.76	109.40
3	D	64	DC	O4'-C1'-N1	6.40	112.48	108.00
3	D	53	DC	O4'-C1'-N1	6.23	112.36	108.00
2	C	27	DG	O4'-C1'-N9	6.12	112.28	108.00
3	D	63	DA	O4'-C1'-N9	6.02	112.21	108.00
2	C	28	DG	O4'-C1'-C2'	-5.98	101.11	105.90
2	C	28	DG	O4'-C1'-N9	5.92	112.14	108.00
2	C	27	DG	O3'-P-O5'	-5.77	93.03	104.00
1	A	324	GLN	N-CA-CB	5.76	118.37	110.01
3	D	58	DC	O4'-C1'-C2'	-5.68	101.35	105.90
3	D	64	DC	O4'-C1'-C2'	-5.65	101.38	105.90
3	D	52	DT	C4'-C3'-O3'	5.25	122.83	109.70

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})
2	C	16	DG	O4'-C1'-N9	5.24	111.67	108.00
2	C	24	DG	O4'-C1'-C2'	-5.18	101.75	105.90
2	C	26	DG	O3'-P-O5'	-5.13	94.25	104.00
3	D	52	DT	C2'-C3'-O3'	-5.03	96.82	109.40

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	326	ARG	Sidechain

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	574	0	595	9	0
2	C	484	0	256	1	0
3	D	459	0	260	1	0
All	All	1517	0	1111	11	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (11) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:370:ILE:HG22	1:A:382:ILE:HD13	1.69	0.75
1:A:352:LEU:HD23	1:A:374:ILE:HG12	1.82	0.60
1:A:371:ALA:HA	1:A:382:ILE:HD11	1.92	0.52
1:A:353:LEU:HD11	1:A:382:ILE:HG23	1.94	0.49
1:A:371:ALA:HA	1:A:382:ILE:CD1	2.43	0.49
1:A:370:ILE:CG2	1:A:382:ILE:HD13	2.41	0.48
1:A:326:ARG:O	1:A:327:ILE:C	2.59	0.46
1:A:371:ALA:HB2	1:A:382:ILE:HD12	1.99	0.44
2:C:28:DG:C6	2:C:29:DA:C6	3.07	0.42

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:353:LEU:CD1	1:A:382:ILE:HG23	2.49	0.41
3:D:55:DC:H2"	3:D:56:DG:C8	2.57	0.40

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	68/1287 (5%)	59 (87%)	8 (12%)	1 (2%)	8 29

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	336	SER

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	63/1122 (6%)	61 (97%)	2 (3%)	34 59

All (2) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	357	GLU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	386	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no oligosaccharides in this entry.

5.6 Ligand geometry (i)

There are no ligands in this entry.

5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.

6 Map visualisation (i)

This section contains visualisations of the EMDB entry EMD-64057. These allow visual inspection of the internal detail of the map and identification of artifacts.

Images derived from a raw map, generated by summing the deposited half-maps, are presented below the corresponding image components of the primary map to allow further visual inspection and comparison with those of the primary map.

6.1 Orthogonal projections (i)

6.1.1 Primary map



X



Y



Z

6.1.2 Raw map



X



Y



Z

The images above show the map projected in three orthogonal directions.

6.2 Central slices (i)

6.2.1 Primary map

X Index: 150


Y Index: 150


Z Index: 150

6.2.2 Raw map


X Index: 150


Y Index: 150


Z Index: 150

The images above show central slices of the map in three orthogonal directions.

6.3 Largest variance slices (i)

6.3.1 Primary map

X Index: 149


Y Index: 172


Z Index: 107

6.3.2 Raw map


X Index: 149


Y Index: 171


Z Index: 107

The images above show the largest variance slices of the map in three orthogonal directions.

6.4 Orthogonal standard-deviation projections (False-color) (i)

6.4.1 Primary map


X


Y


Z

6.4.2 Raw map


X


Y


Z

The images above show the map standard deviation projections with false color in three orthogonal directions. Minimum values are shown in green, max in blue, and dark to light orange shades represent small to large values respectively.

6.5 Orthogonal surface views (i)

6.5.1 Primary map

X Y Z

The images above show the 3D surface view of the map at the recommended contour level 0.05. These images, in conjunction with the slice images, may facilitate assessment of whether an appropriate contour level has been provided.

6.5.2 Raw map

X Y Z

These images show the 3D surface of the raw map. The raw map's contour level was selected so that its surface encloses the same volume as the primary map does at its recommended contour level.

6.6 Mask visualisation (i)

This section shows the 3D surface view of the primary map at 50% transparency overlaid with the specified mask at 0% transparency

A mask typically either:

- Encompasses the whole structure
- Separates out a domain, a functional unit, a monomer or an area of interest from a larger structure

6.6.1 emd_64057_msk_1.map (i)



X



Y



Z

7 Map analysis (i)

This section contains the results of statistical analysis of the map.

7.1 Map-value distribution (i)



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.

7.2 Volume estimate (i)



The volume at the recommended contour level is 22 nm^3; this corresponds to an approximate mass of 20 kDa.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.

7.3 Rotationally averaged power spectrum (i)



*Reported resolution corresponds to spatial frequency of 0.318 Å^{-1}

8 Fourier-Shell correlation (i)

Fourier-Shell Correlation (FSC) is the most commonly used method to estimate the resolution of single-particle and subtomogram-averaged maps. The shape of the curve depends on the imposed symmetry, mask and whether or not the two 3D reconstructions used were processed from a common reference. The reported resolution is shown as a black line. A curve is displayed for the half-bit criterion in addition to lines showing the 0.143 gold standard cut-off and 0.5 cut-off.

8.1 FSC (i)



*Reported resolution corresponds to spatial frequency of 0.318 Å^{-1}

8.2 Resolution estimates (i)

Resolution estimate (Å)	Estimation criterion (FSC cut-off)		
	0.143	0.5	Half-bit
Reported by author	3.14	-	-
Author-provided FSC curve	3.14	3.49	3.15
Unmasked-calculated*	3.99	7.22	4.09

*Resolution estimate based on FSC curve calculated by comparison of deposited half-maps. The value from deposited half-maps intersecting FSC 0.143 CUT-OFF 3.99 differs from the reported value 3.14 by more than 10 %

9 Map-model fit (i)

This section contains information regarding the fit between EMDB map EMD-64057 and PDB model 9UD0. Per-residue inclusion information can be found in section 3 on page 4.

9.1 Map-model overlay (i)



X



Y



Z

The images above show the 3D surface view of the map at the recommended contour level 0.05 at 50% transparency in yellow overlaid with a ribbon representation of the model coloured in blue. These images allow for the visual assessment of the quality of fit between the atomic model and the map.

9.2 Q-score mapped to coordinate model (i)



The images above show the model with each residue coloured according its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.

9.3 Atom inclusion mapped to coordinate model (i)



The images above show the model with each residue coloured according to its atom inclusion. This shows to what extent they are inside the map at the recommended contour level (0.05).

9.4 Atom inclusion (i)



At the recommended contour level, 89% of all backbone atoms, 81% of all non-hydrogen atoms, are inside the map.

9.5 Map-model fit summary (i)

The table lists the average atom inclusion at the recommended contour level (0.05) and Q-score for the entire model and for each chain.

Chain	Atom inclusion	Q-score
All	0.8070	0.4060
A	0.6260	0.3680
C	0.9280	0.4240
D	0.9000	0.4340