Full wwPDB EM Validation Report (i)

Oct 19, 2026 – 02:19 PM JST

PDB ID	:	9UTP / pdb_00009utp
EMDB ID	:	EMD-64496
Title	:	Cryo-EM structure of the oxytocin receptor bound to nanobody Nb2B11
Authors	:	Gao, K.; Nie, J.; Zhang, X.; Liu, X.
Deposited on	:	2025-05-05
Resolution	:	3.39 Å(reported)

This is a Full wwPDB EM Validation Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org*
A user guide is available at
https://www.wwpdb.org/validation/2017/EMValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at
http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (i)) were used in the production of this report:

EMDB validation analysis	:	0.0.1.dev133
MolProbity	:	4-5-2 with Phenix2.2
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
EM percentile statistics	:	202505.v01 (Using data in the EMDB archive up until May 2025)
MapQ	:	1.9.13
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.51

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:
ELECTRON MICROSCOPY

The reported resolution of this entry is 3.39 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.





Metric	Whole archive (#Entries)	EM structures (#Entries)	Similar EM resolution (#Entries, resolution range(Å))
Clashscore	229148	23984	-
Ramachandran outliers	224038	23583	-
Sidechain outliers	223484	23102	-
Q-score	-	25397	14220 (2.89 - 3.89)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion < 40%). The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	446	9% 53% 12% • 35%
2	B	126	13% • 87%

2 Entry composition (i)

There are 2 unique types of molecules in this entry. The entry contains 2193 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Oxytocin receptor.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
1	A	292	2082	1383	335	346	18	0	0

There are 66 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-7	ASP	-	expression tag	UNP P30559
A	-6	TYR	-	expression tag	UNP P30559
A	-5	LYS	-	expression tag	UNP P30559
A	-4	ASP	-	expression tag	UNP P30559
A	-3	ASP	-	expression tag	UNP P30559
A	-2	ASP	-	expression tag	UNP P30559
A	-1	ASP	-	expression tag	UNP P30559
A	0	ALA	-	expression tag	UNP P30559
A	65	HIS	ARG	conflict	UNP P30559
A	120	LEU	VAL	conflict	UNP P30559
A	153	ALA	ASP	conflict	UNP P30559
A	167	VAL	ALA	conflict	UNP P30559
A	193	ARG	GLN	conflict	UNP P30559
A	218	THR	ALA	conflict	UNP P30559
A	322	CYS	SER	conflict	UNP P30559
A	325	LYS	ASN	conflict	UNP P30559
A	333	MET	THR	conflict	UNP P30559
A	995	HIS	-	expression tag	UNP P30559
A	996	HIS	-	expression tag	UNP P30559
A	997	HIS	-	expression tag	UNP P30559
A	998	HIS	-	expression tag	UNP P30559
A	999	HIS	-	expression tag	UNP P30559
A	1000	HIS	-	expression tag	UNP P30559
A	1001	ALA	-	expression tag	UNP P30559
A	1002	PHE	-	expression tag	UNP P30559
A	1003	LYS	-	expression tag	UNP P30559
A	1004	ILE	-	expression tag	UNP P30559
A	1005	TRP	-	expression tag	UNP P30559

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
A	1006	GLN	-	expression tag	UNP P30559
A	1007	ASN	-	expression tag	UNP P30559
A	1008	SER	-	expression tag	UNP P30559
A	1009	ALA	-	expression tag	UNP P30559
A	1010	LEU	-	expression tag	UNP P30559
A	1011	GLY	-	expression tag	UNP P30559
A	1012	LYS	-	expression tag	UNP P30559
A	1013	GLU	-	expression tag	UNP P30559
A	1014	LEU	-	expression tag	UNP P30559
A	1015	ALA	-	expression tag	UNP P30559
A	1016	ALA	-	expression tag	UNP P30559
A	1085	GLU	-	expression tag	UNP P30559
A	1086	PRO	-	expression tag	UNP P30559
A	1087	VAL	-	expression tag	UNP P30559
A	1088	ALA	-	expression tag	UNP P30559
A	1089	ILE	-	expression tag	UNP P30559
A	1090	LEU	-	expression tag	UNP P30559
A	1091	THR	-	expression tag	UNP P30559
A	1092	LEU	-	expression tag	UNP P30559
A	1093	SER	-	expression tag	UNP P30559
A	1094	PHE	-	expression tag	UNP P30559
A	1095	LYS	-	expression tag	UNP P30559
A	1119	ALA	-	expression tag	UNP P30559
A	1120	TYR	-	expression tag	UNP P30559
A	1121	LYS	-	expression tag	UNP P30559
A	1122	GLY	-	expression tag	UNP P30559
A	1123	LEU	-	expression tag	UNP P30559
A	1124	GLU	-	expression tag	UNP P30559
A	1125	GLU	-	expression tag	UNP P30559
A	1126	GLY	-	expression tag	UNP P30559
A	1127	SER	-	expression tag	UNP P30559
A	1128	LYS	-	expression tag	UNP P30559
A	1129	ALA	-	expression tag	UNP P30559
A	1130	GLY	-	expression tag	UNP P30559
A	1131	GLU	-	expression tag	UNP P30559
A	1132	ALA	-	expression tag	UNP P30559
A	1133	THR	-	expression tag	UNP P30559
A	1134	MET	-	expression tag	UNP P30559

- Molecule 2 is a protein called Nb2B11.

Mol	Chain	Residues	Atoms				AltConf	Trace
			Total	C	N	O		
2	B	17	111	69	18	24	0	0

3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and atom inclusion in map density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red diamond above a residue indicates a poor fit to the EM map for this residue (all-atom inclusion < 40%). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Oxytocin receptor

Chain A:



- Molecule 2: Nb2B11

Chain B:



4 Experimental information (i)

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, Not provided	
Number of particles used	274430	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	PHASE FLIPPING ONLY	Depositor
Microscope	TFS KRIOS	Depositor
Voltage (kV)	300	Depositor
Electron dose ($e^-/Å^2$)	50	Depositor
Minimum defocus (nm)	1100	Depositor
Maximum defocus (nm)	1800	Depositor
Magnification	Not provided	
Image detector	GATAN K3 (6k x 4k)	Depositor
Maximum map value	4.912	Depositor
Minimum map value	-3.607	Depositor
Average map value	-0.003	Depositor
Map value standard deviation	0.091	Depositor
Recommended contour level	0.6	Depositor
Map size (Å)	216.5, 216.5, 216.5	wwPDB
Map dimensions	200, 200, 200	wwPDB
Map angles (°)	90.0, 90.0, 90.0	wwPDB
Pixel spacing (Å)	1.0825, 1.0825, 1.0825	Depositor

5 Model quality (i)

5.1 Standard geometry (i)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\#\|Z\| >5$	RMSZ	$\#\|Z\| >5$
1	A	0.18	0/2123	0.34	0/2908
2	B	0.13	0/112	0.53	0/151
All	All	0.18	0/2235	0.36	0/3059

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2082	0	2014	42	0
2	B	111	0	97	0	0
All	All	2193	0	2111	42	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 10.

All (42) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:141:ILE:HD12	1:A:1089:ILE:HD13	1.50	0.92
1:A:141:ILE:O	1:A:1086:PRO:HG3	1.77	0.84

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:333:MET:HB2	1:A:336:LEU:HB2	1.62	0.81
1:A:141:ILE:CD1	1:A:1089:ILE:HD13	2.11	0.80
1:A:223:ILE:C	1:A:1001:ALA:N	2.40	0.79
1:A:171:GLN:HA	1:A:174:ILE:HG12	1.66	0.76
1:A:137:ARG:HH21	1:A:273:THR:HG21	1.56	0.71
1:A:273:THR:CG2	1:A:1132:ALA:O	2.45	0.64
1:A:144:PRO:CB	1:A:1086:PRO:HD2	2.28	0.63
1:A:273:THR:HG21	1:A:1132:ALA:O	1.99	0.63
1:A:202:THR:HA	1:A:296:MET:HE1	1.80	0.62
1:A:305:PRO:HB2	1:A:308:ALA:HB2	1.82	0.61
1:A:79:LYS:HG2	1:A:80:HIS:HD2	1.67	0.58
1:A:276:MET:HE1	1:A:333:MET:HE3	1.85	0.58
1:A:221:GLY:O	1:A:1002:PHE:CB	2.52	0.58
1:A:1130:GLY:O	1:A:1134:MET:N	2.38	0.57
1:A:80:HIS:ND1	1:A:157:VAL:HG11	2.20	0.57
1:A:288:TRP:HE1	1:A:321:ASN:HD21	1.53	0.56
1:A:73:ARG:HH12	1:A:140:ALA:HB2	1.72	0.54
1:A:221:GLY:O	1:A:1002:PHE:HB2	2.07	0.54
1:A:120:LEU:HD13	1:A:171:GLN:OE1	2.07	0.54
1:A:274:VAL:HG12	1:A:1134:MET:HA	1.91	0.53
1:A:276:MET:HE1	1:A:333:MET:CE	2.41	0.51
1:A:212:PRO:O	1:A:216:LEU:HB2	2.12	0.50
1:A:128:TYR:HE1	1:A:212:PRO:HG3	1.77	0.49
1:A:125:ALA:O	1:A:129:LEU:HB2	2.13	0.47
1:A:272:ARG:N	1:A:1131:GLU:O	2.48	0.47
1:A:118:LEU:HA	1:A:121:VAL:HB	1.97	0.47
1:A:221:GLY:O	1:A:1002:PHE:HB3	2.15	0.46
1:A:297:TRP:HZ2	1:A:305:PRO:HD2	1.80	0.46
1:A:140:ALA:O	1:A:1085:GLU:CB	2.63	0.46
1:A:108:PRO:HD2	1:A:111:LEU:HB2	1.97	0.44
1:A:141:ILE:C	1:A:1086:PRO:HG3	2.38	0.44
1:A:144:PRO:CB	1:A:1086:PRO:CD	2.93	0.44
1:A:137:ARG:NH2	1:A:273:THR:HG21	2.30	0.44
1:A:57:ASN:HB3	1:A:86:LEU:HG	2.00	0.43
1:A:120:LEU:HD13	1:A:171:GLN:CD	2.44	0.42
1:A:1089:ILE:HG21	1:A:1134:MET:HB3	2.02	0.41
1:A:274:VAL:CG1	1:A:1134:MET:HA	2.50	0.41
1:A:57:ASN:HD21	1:A:85:ASP:HB3	1.86	0.41
1:A:201:ILE:HD12	1:A:201:ILE:HA	1.95	0.41
1:A:282:LEU:HD23	1:A:282:LEU:HA	1.92	0.41

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	282/446 (63%)	272 (96%)	10 (4%)	0	100	100
2	B	15/126 (12%)	10 (67%)	5 (33%)	0	100	100
All	All	297/572 (52%)	282 (95%)	15 (5%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	193/363 (53%)	184 (95%)	9 (5%)	23	50
2	B	10/99 (10%)	9 (90%)	1 (10%)	7	26
All	All	203/462 (44%)	193 (95%)	10 (5%)	24	49

All (10) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	51	LEU
1	A	121	VAL
1	A	141	ILE
1	A	175	PHE
1	A	188	TRP
1	A	295	GLN
1	A	311	PHE
1	A	314	VAL

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	1134	MET
2	B	104	SER

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no oligosaccharides in this entry.

5.6 Ligand geometry (i)

There are no ligands in this entry.

5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

The following chains have linkage breaks:

Mol	Chain	Number of breaks
1	A	2

All chain breaks are listed below:

Model	Chain	Residue-1	Atom-1	Residue-2	Atom-2	Distance (Å)
1	A	1016:ALA	C	1085:GLU	N	24.39
1	A	1095:LYS	C	1119:ALA	N	15.63

6 Map visualisation (i)

This section contains visualisations of the EMDB entry EMD-64496. These allow visual inspection of the internal detail of the map and identification of artifacts.

Images derived from a raw map, generated by summing the deposited half-maps, are presented below the corresponding image components of the primary map to allow further visual inspection and comparison with those of the primary map.

6.1 Orthogonal projections (i)

6.1.1 Primary map



X



Y



Z

6.1.2 Raw map



X



Y



Z

The images above show the map projected in three orthogonal directions.

6.2 Central slices (i)

6.2.1 Primary map


X Index: 100


Y Index: 100


Z Index: 100

6.2.2 Raw map


X Index: 100


Y Index: 100


Z Index: 100

The images above show central slices of the map in three orthogonal directions.

6.3 Largest variance slices (i)

6.3.1 Primary map


X Index: 98


Y Index: 100


Z Index: 97

6.3.2 Raw map


X Index: 98


Y Index: 99


Z Index: 97

The images above show the largest variance slices of the map in three orthogonal directions.

6.4 Orthogonal standard-deviation projections (False-color) (i)

6.4.1 Primary map


X


Y


Z

6.4.2 Raw map


X


Y


Z

The images above show the map standard deviation projections with false color in three orthogonal directions. Minimum values are shown in green, max in blue, and dark to light orange shades represent small to large values respectively.

6.5 Orthogonal surface views (i)

6.5.1 Primary map

X Y Z

The images above show the 3D surface view of the map at the recommended contour level 0.6. These images, in conjunction with the slice images, may facilitate assessment of whether an appropriate contour level has been provided.

6.5.2 Raw map

X Y Z

These images show the 3D surface of the raw map. The raw map's contour level was selected so that its surface encloses the same volume as the primary map does at its recommended contour level.

6.6 Mask visualisation (i)

This section shows the 3D surface view of the primary map at 50% transparency overlaid with the specified mask at 0% transparency

A mask typically either:

- Encompasses the whole structure
- Separates out a domain, a functional unit, a monomer or an area of interest from a larger structure

6.6.1 emd_64496_msk_1.map (i)



X



Y



Z

7 Map analysis (i)

This section contains the results of statistical analysis of the map.

7.1 Map-value distribution (i)



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.

7.2 Volume estimate (i)



The volume at the recommended contour level is 24 nm^3; this corresponds to an approximate mass of 22 kDa.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.

7.3 Rotationally averaged power spectrum (i)



*Reported resolution corresponds to spatial frequency of 0.295 Å^{-1}

8 Fourier-Shell correlation (i)

Fourier-Shell Correlation (FSC) is the most commonly used method to estimate the resolution of single-particle and subtomogram-averaged maps. The shape of the curve depends on the imposed symmetry, mask and whether or not the two 3D reconstructions used were processed from a common reference. The reported resolution is shown as a black line. A curve is displayed for the half-bit criterion in addition to lines showing the 0.143 gold standard cut-off and 0.5 cut-off.

8.1 FSC (i)



*Reported resolution corresponds to spatial frequency of 0.295 Å^{-1}

8.2 Resolution estimates (i)

Resolution estimate (Å)	Estimation criterion (FSC cut-off)		
	0.143	0.5	Half-bit
Reported by author	3.39	-	-
Author-provided FSC curve	-	-	-
Unmasked-calculated*	3.86	4.26	3.94

*Resolution estimate based on FSC curve calculated by comparison of deposited half-maps. The value from deposited half-maps intersecting FSC 0.143 CUT-OFF 3.86 differs from the reported value 3.39 by more than 10 %

9 Map-model fit (i)

This section contains information regarding the fit between EMDB map EMD-64496 and PDB model 9UTP. Per-residue inclusion information can be found in section 3 on page 6.

9.1 Map-model overlay (i)



X



Y



Z

The images above show the 3D surface view of the map at the recommended contour level 0.6 at 50% transparency in yellow overlaid with a ribbon representation of the model coloured in blue. These images allow for the visual assessment of the quality of fit between the atomic model and the map.

9.2 Q-score mapped to coordinate model (i)



X Y Z

The images above show the model with each residue coloured according its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.

9.3 Atom inclusion mapped to coordinate model (i)



X Y Z

The images above show the model with each residue coloured according to its atom inclusion. This shows to what extent they are inside the map at the recommended contour level (0.6).

9.4 Atom inclusion (i)



At the recommended contour level, 73% of all backbone atoms, 74% of all non-hydrogen atoms, are inside the map.

9.5 Map-model fit summary (i)

The table lists the average atom inclusion at the recommended contour level (0.6) and Q-score for the entire model and for each chain.

Chain	Atom inclusion	Q-score
All	0.7350	0.4370
A	0.7330	0.4380
B	0.7710	0.4210